
















Original Research

Comparative Genomics of Two Novel *Bacillus* Strains: Microbiomic Insights into the Sequences, Metabolomics, and Potential Safe Use in the Creation of Biopreparations

Valentina A. Filippova^{1,2}, Georgi Yu. Laptev^{1,2}, Larisa A. Ilina^{1,2},
Elena A. Yildirim^{1,2}, Ekaterina S. Ponomareva², Evgeni A. Brazhnik¹,
Natalia I. Novikova², Daria G. Tiurina², Andrei V. Dubrovin^{2,3}, Alisa S. Dubrovina²,
Kseniya A. Sokolova^{1,2}, Irina A. Klyuchnikova¹, Vasiliy A. Zaikin²,
Darren K. Griffin^{4,5}, Michael N. Romanov^{2,4,6,*}

¹Molecular Genetics and Microbiomics Laboratory, BIOTROF Ltd., 196602 Pushkin, St. Petersburg, Russia

²Federal State Budgetary Educational Institution of Higher Education “St. Petersburg State Agrarian University”, 196605 Pushkin, St. Petersburg, Russia

³Information Technologies, Mechanics and Optics (ITMO) University, 197101 St. Petersburg, Russia

⁴School of Natural Sciences, University of Kent, CT2 7NJ Canterbury, Kent, UK

⁵Animal Genomics and Bioresource Research Unit (AGB Research Unit), Faculty of Science, Kasetsart University, 10900 Chatuchak, Bangkok, Thailand

⁶L. K. Ernst Federal Research Centre for Animal Husbandry, Dubrovitsy, 142132 Moscow Oblast, Russia

*Correspondence: m.romanov@kent.ac.uk (Michael N. Romanov)

Academic Editor: Baohong Zhang

Submitted: 14 November 2024 Revised: 8 January 2025 Accepted: 12 February 2025 Published: 19 June 2025

Abstract

Background: *Bacillus* bacteria are often used in the production of biopreparations. Moreover, these bacteria can be used in agriculture as probiotics or starters for manufacturing fodder preserved by fermentation (silage). The ability of *Bacillus* bacteria to produce many biologically active molecules and metabolites with antimicrobial activity means that these bacteria can stimulate plant growth and restore the balance of the microbiome in the digestive system of certain animals. **Methods:** Using molecular biological analysis, bioinformatic annotation, and metabolic profiling of whole genome sequences, we analyzed two promising candidates for creating biopreparations, i.e., two *Bacillus* strains, namely *B. mucilaginosus* 159 and *B. subtilis* 111. We compared the genomes of these two strains and characterized both their microbiomic and metabolomic features. **Results:** We demonstrated that both strains lacked elements contributing to the formation of toxic and virulent properties; however, both exhibited potential in the biosynthesis of B vitamins and siderophores. Additionally, these strains could synthesize many antimicrobial substances of different natures and spectrums of action. *B. mucilaginosus* 159 could synthesize macrolactin H (an antibiotic from the polyketide group), mersacidin (a class II lanthipeptide), and bacilysin. Meanwhile, *B. subtilis* 111 could synthesize andalusicin (a class III lanthipeptide), bacilysin, macrolactin H, difficidin, bacillaene (a polyene antibiotic), fengycin (a lipopeptide with antifungal activity), and surfactin (another lipopeptide). Further, a unique pathway of intracellular synthesis of the osmoprotectant glycine betaine was identified in *B. subtilis* 111, with the participation of betaine aldehyde dehydrogenase (BetB); this is not widely represented in bacteria of the genus *Bacillus*. These compounds can increase osmotic stability, which may be key for manufacturing biological starters for silage preparation. **Conclusions:** These two *Bacillus* strains are safe for use as probiotic microorganisms or starters in producing preserved fodder. However, *B. subtilis* 111 may be preferable due to a wider spectrum of synthesized antimicrobial substances and vitamins. Our findings exemplify using genomic technologies to describe the microbiomic and metabolomic characteristics of significant bacterial groups such as *Bacillus* species.

Keywords: *Bacillus* spp.; biopreparations; probiotics; silage; whole genome sequencing; functional microbiomics; metabolomics; antimicrobial substances; vitamins; lipopeptides

1. Introduction

Efforts to transition to more environmentally friendly agriculture systems are now commonplace worldwide. The following are desirable aspects of environmentally friendly agriculture: (1) refusal to use particular chemicals in crop production; (2) refusal to use other certain chemicals for feed antibiotics in livestock farming; (3) increasing the effi-

ciency of roughage assimilation; (4) improving the quality of plant feed preparation; (5) reducing the share of compound feed for ruminants and other animals [1–6]. Using microorganisms with various enzymatic and metabolic activities represents one of the main components of the new environmentally friendly approach to agriculture [7–13]. Enzyme, probiotic, prebiotic, and combined enzyme–



probiotic preparations are increasingly used to improve efficiency and stimulate the growth and development of ruminants and other animals. This can enhance non-specific immunity and improve the quality of feed preparation [14–19]. The main purpose of using so-called “biopreparations” in feed production is to facilitate the creation of a microbial community that will lead to rapid acidification of the environment and prevent the development of pathogenic and opportunistic microorganisms [20–23], such as clostridia and enterobacteria [24–27]. Clostridia, which are found on crops and in the soil as spores, grow anaerobically. Clostridia also produce butyric acid and breaks down amino acids, making the silage tasteless and possessing reduced nutritional value. Enterobacteria are non-spore-forming facultative anaerobes that ferment sugars to acetic acid and other products and can break down amino acids [27–31].

The most commonly used microorganisms as probiotic preparations include bacilli, lactobacilli, and bifidobacteria [17,22,32,33]. Moreover, the functional microbiomics of probiotic bacteria and their influence on related and other gut microbes have received particular attention [34–40]. The field of “gut microbiomics” investigates the function of the intestinal microbiota by examining its overall “microbiome”, including the genomes of its individual representatives [39,41]. Microbiomics provides molecular tools for analyzing microbial communities, facilitating their comprehensive characterization. Thus, using such microbiomic approaches can provide more information regarding the function of a certain bacterial group [12,42]. Strategies for developing food and feed preparations take advantage of incorporating genomic and microbiomic studies [38,43–45], which generally rely upon omics and multi-omics approaches [22,46–48]. The term “microbiomic” is currently widely used in the scientific literature [49–54] as an adjective to mean “of, or relating to, the microbiome or to microbiomics” [55]. Paleri [56] defines microbiomic research as “Microbiomic studies cover the entire life systems as the collective genomes of the microbes that are composed of bacteria, bacteriophage, fungi, protozoa and viruses.” Thus, the term “microbiomic” was used in the present study in relation to this accepted terminology.

According to many studies, the genus *Bacillus* has several useful properties, making these bacteria effective biopreparations [57–61]. These properties include (1) high preservation, (2) stability during storage, (3) antagonistic activity to a wide range of pathogenic and opportunistic microorganisms, and (4) high enzymatic activity. Bacteria in the genus *Bacillus* can form endospores, which allow them to remain viable under extreme conditions, i.e., at high or low temperatures, radiation, suboptimal pH, pressure, and in the presence of toxic chemicals that damage vegetative cells [62–64]. The ability to synthesize antibiotics, bacteriocins, cyclic lipopeptides, and lytic enzymes with antimicrobial activity is characteristic of the probiotic activity of bacteria in the genus *Bacillus* [22,23,26,65–68].

The genomes of microorganisms from this taxonomic group have many loci that determine the biosynthesis of antimicrobial compounds [19]. This provides one of their most important functional properties that are considered when choosing suitable microorganisms for drug production [22]. Herewith, the metabolomic data [69,70] suggest that *Bacillus* spp. can perform metabolism via the pentose phosphate pathway, making them effective producers of vitamins, among which the most significant are cobalamin, riboflavin, folic acid, and biotin [71–73]. Whole genome sequencing (WGS) allows us to assess fully the biotechnological capabilities of promising strains [61,74,75]. As a result, we can identify the genetic determinants that are associated with the possibility of biosynthesis of various biologically active substances important for creating biopreparations for agriculture. At the same time, we can confirm the absence of different virulence factors that designate the biological safety of using microorganism strains [22,46].

Among the issues associated with creating probiotic preparations, it is important to reveal the presence of unknown properties in strains, both positive (synthesis of antimicrobial substances and various enzymatic activities) and negative (synthesis of undesirable substances and the presence of pathogenicity factors). The development of new technologies in molecular genetics makes it possible to rapidly determine the complete genome of microorganisms [61,74,75]. The use of next-generation sequencing (NGS) technologies, which include WGS, for the analysis of strains of probiotic microorganisms allows for a complete characterization of a strain under study and predicts its capabilities [61], thus facilitating the creation of an effective and safe probiotic drug [22].

Among studies on the genomes of bacilli strains, Kapse *et al.* [60] performed WGS to examine the *B. coagulans* HS243 strain. This genome study revealed the presence of several marker genes that conform to probiotic features. In particular, genome analysis of HS243 revealed the presence of multi-subunit ATPases, hydrolases, and adhesion proteins necessary for survival and colonization of the intestine. The HS243 genome also contains genes for the biosynthesis of vitamins and essential amino acids, which suggests that HS243 can be used as a food additive. The strain was found to have genes responsible for producing bacteriocins; thus, it is apparent that the *B. coagulans* HS243 strain can potentially prevent diseases. Sulthana *et al.* [61] confirmed the genetic safety of the *B. subtilis* UBBS-14 strain using WGS. It was shown that this strain lacks antibiotic-resistance genes (within its mobile genetic elements) and pathogenicity factors. Therefore, it can be considered biologically safe.

The present study aimed to generate and compare the genomes of two promising strains to create biopreparations: *B. mucilaginosus* 159 and *B. subtilis* 111. A comprehensive characterization and assessment of *B. mucilaginosus* 159 and *B. subtilis* 111 were achieved using molecular biolog-

ical analysis and bioinformatic annotations regarding their biological properties and potential use in antimicrobial feed additives.

2. Materials and Methods

2.1 Strains, DNA Extraction, and Whole Genome Sequencing

The *B. subtilis* 111 and *B. mucilaginosus* 159 strains were obtained from the BIOTROF LLC collection (Pushkin, St. Petersburg, Russia). Total DNA was isolated from the strains using the Genomic DNA Purification kit (Thermo Fisher Scientific, Inc., Waltham, MA, USA). Furthermore, we employed the Nextera DNA Flex Library Prep kit (Illumina, Inc., San Diego, CA, USA) to prepare the DNA library samples for WGS using the Illumina MiSeq platform. The MiSeq Reagent kit v3 (300 cycles; Illumina) was utilized to sequence the resultant libraries [76–78]. The above purification, preparation, and reagent commercial kits were used following the in-house protocols [38,78–80] based on the manufacturers' instructions.

2.2 Bioinformatics Analysis

Sequence quality assessment was performed using the specialized FastQC software [81]. Unreliable sequences and adapters were removed using the Trimmomatic-0.38 program [82]. Paired-end nucleotide sequences, filtered by a length of at least 50 base pairs (bp), were assembled *de novo* using the genome assemblers SPAdes-3.11.1 (for *B. subtilis* 111) and SPAdes v. 3.12.0 (for *B. mucilaginosus* 159) [83]. The genome coverage for *B. subtilis* 111 and *B. mucilaginosus* 159 was 100× and 160×, respectively. The resultant genome assemblies in the form of contigs were validated and deposited using the NCBI BioSample and BioProject databases [84], with the following assigned respective BioSample and BioProject identifiers (accession numbers): SUB14629276 and PRJNA1141124 for *B. subtilis* 111, and SAMN33397894 and PRJNA937299 for *B. mucilaginosus* 159. The contigs assembled and obtained this way were used for subsequent analysis, including annotation for genes (protein-coding genes) using NCBI RefSeq version 6.7 [84]. Additionally, for the independent comparison, a well-documented and annotated genome of the related model microorganism *B. subtilis* subsp. *subtilis* str. 168 [85], with the BioSample and BioProject identifiers SAMEA3138188 and PRJNA76, respectively, was considered.

Functional annotation of genomic sequences was performed using the RAST 2.0 web server (RAST, Rapid Annotations Using Subsystems Technology; [86]). Herewith, the annotation scheme followed Classic RAST, while the FIGfams protein collection version 70 [87] was used to process the genome. An additional annotation was performed using Prokka version 1.13.3 [88] and the Kyoto Encyclopedia of Genes and Genomes (KEGG) Pathway database [89,90]. In particular, the resultant amino acid sequences

were uploaded to the KEGG Automated Annotation Server (KAAS). The sequences were processed using the BBH (bi-directional best hit) method and the BLAST search program [89,90].

To search for gene clusters involved in the biosynthesis of secondary metabolites [46], the antiSMASH version 7.1.0 web service [91] was used in the relaxed detection strictness mode, followed by refinements in the NCBI Protein database [92]. The Norine database [93] was utilized to identify non-ribosomal peptides (NRPs). The search for prophage sequences was performed using the PHASTER web server [94]. The virulence of genomes was analyzed using the VirulenceFinder-2.0 (version 2.0.5), with the parameters set to the %ID threshold of 90% and the minimum sequence length of 60% program, and with the support of the Center for Genomic Epidemiology web service (National Food Institute, Technical University of Denmark, Lyngby, Denmark; [95]). The antibiotic resistance determinants were searched and assessed using the ResFinder program (version 4.6.0), with all parameters specified by default [96].

3. Results and Discussion

3.1 Major Characteristics of the *Bacillus* spp. Strain Genomes

As illustrated in Table 1, the genome sizes of *B. subtilis* 111 and *B. mucilaginosus* 159 were established as similar: 3,949,468 and 3,970,760 bp, respectively. The guanine-cytosine (GC) composition was similar in percentage terms and amounted to 46.6% for *B. subtilis* 111 and 46.3% for *B. mucilaginosus* 159. According to the ResFinder assessment, no antibiotic resistance determinants were found in the genomes of *B. subtilis* 111 and *B. mucilaginosus* 159. A search for genes encoding putative virulence factors by the VirulenceFinder program also showed that these were absent in *B. subtilis* 111 and *B. mucilaginosus* 159.

We identified all the main groups of genes for proteins that characterize the metabolomic profiles in the genomes of the *B. subtilis* 111 and *B. mucilaginosus* 159 strains. These proteins jointly mediate and modulate certain biological processes involved in the functions of amino acid transport and metabolism, as well as transcription, translation, and carbohydrate/protein transport and metabolism. The strains had products that resembled a full set of functional metabolic pathways, including glycolysis, the tricarboxylic acid cycle, and the pentose phosphate pathway.

As presented in Table 2 (data produced using RAST 2.0 [86]) and Fig. 1 (data produced using RAST 2.0 [86] and antiSMASH 7.1.0 [91]), analysis of the genomes of *B. subtilis* 111 and *B. mucilaginosus* 159 in comparison to one another showed that the genomes of the studied bacteria contained a large number of genetic determinants. These define the metabolomic profiles of the strains, including the metabolism of carbohydrates, proteins, and amino acids.

Table 1. The main characteristics of the two probiotic *Bacillus* strain genomes compared to the model microorganism.

| Assembly characteristics | <i>B. subtilis</i> 111 | <i>B. mucilaginosus</i> 159 | Model microorganism <i>B. subtilis</i> subsp. <i>subtilis</i> str. 168 |
|--|------------------------|-----------------------------|--|
| Genome assembly size, bp | 3,941,553 | 3,970,760 | 4,214,814 |
| Contig N50 | 2,109,194 | 500,910 | 4,215,606 |
| Guanine-cytosine (GC) composition, % | 46.6 | 46.3 | 43.5 |
| Number of open reading frames (coding sequences) | 3991 | 4002 | 4114 |
| Genes | 3964 | 3983 | 4536 |
| Protein-coding genes | 3807 | 3817 | 4237 |
| Number of subsystems | 444 | 438 | 515 |
| Number of RNAs | 87 | 73 | 181 |

Table 2. Comparative analysis of the cellular metabolic systems of the two *Bacillus* bacteria as performed using RAST 2.0 (<https://rast.nmpdr.org/>).

| Metabolic systems | Distribution of the number of genes by functions of bacterial metabolic systems | | | | | |
|---|---|------|------------------------|------|---|------|
| | <i>B. mucilaginosus</i> 159 | | <i>B. subtilis</i> 111 | | Model organism <i>B. subtilis</i> subsp. <i>subtilis</i> str. 168 | |
| | Total | % | Total | % | Total | % |
| Protein metabolism | 175 | 6.8 | 190 | 7.3 | 242 | 7.1 |
| Carbohydrate metabolism | 372 | 14.5 | 373 | 14.4 | 535 | 15.8 |
| Metabolism of amino acids and their derivatives | 410 | 16.0 | 401 | 15.4 | 450 | 13.3 |
| Metabolism of nucleosides and nucleotides | 110 | 4.3 | 105 | 4.0 | 145 | 4.3 |
| Metabolism of fatty acids, lipids and isoprenoids | 136 | 5.3 | 134 | 5.2 | 102 | 3.0 |
| Metabolism of aromatic compounds | 11 | 0.4 | 11 | 0.4 | 10 | 0.3 |
| RNA metabolism | 156 | 6.1 | 155 | 6.0 | 198 | 5.8 |
| DNA metabolism | 79 | 3.1 | 84 | 3.2 | 136 | 4.0 |
| Sulfur metabolism | 39 | 1.5 | 39 | 1.5 | 44 | 1.3 |
| Phosphorus metabolism | 33 | 1.3 | 28 | 1.1 | 25 | 0.7 |
| Nitrogen metabolism | 29 | 1.1 | 31 | 1.2 | 30 | 0.9 |
| Systems determining potassium balance | 8 | 0.3 | 8 | 0.3 | 26 | 0.8 |
| Iron entry into the cell | 19 | 0.7 | 28 | 1.1 | 32 | 0.9 |
| Synthesis of cofactors, vitamins, prosthetic groups, and pigments | 202 | 7.9 | 230 | 8.9 | 311 | 9.2 |
| Cell division and cell cycle | 52 | 2.0 | 52 | 2.0 | 48 | 1.4 |
| Motility and chemotaxis | 84 | 3.3 | 83 | 3.2 | 69 | 2.0 |
| Regulation and signaling systems | 63 | 2.5 | 61 | 2.3 | 50 | 1.5 |
| Secondary metabolism | 6 | 0.2 | 5 | 0.2 | 6 | 0.2 |
| Dormancy and sporulation | 114 | 4.4 | 114 | 4.4 | 141 | 4.2 |
| Respiration process | 59 | 2.3 | 59 | 2.3 | 91 | 2.7 |
| Response to stress factors | 102 | 4.0 | 100 | 3.9 | 111 | 3.3 |
| Cell membrane transport systems | 60 | 2.3 | 58 | 2.2 | 84 | 2.5 |
| Cell wall and capsule synthesis | 128 | 5.0 | 125 | 4.8 | 142 | 4.2 |
| Cellular protection | 64 | 2.5 | 63 | 2.4 | 77 | 2.3 |
| Others | 56 | 2.2 | 60 | 2.3 | 290 | 8.5 |
| Total | 2567 | 100 | 2597 | 100 | 3395 | 100 |

Here, the strain *B. subtilis* 111 was found to have a higher proportion of genes involved in protein metabolism, i.e., 190 vs. 175 genes in *B. mucilaginosus* 159. A higher proportion of genes were also involved in the biosynthesis of cofactors, vitamins, prosthetic groups and pigments (230 vs. 202 genes) and more genes were involved in iron entry into the cell (28 vs. 19 genes). In contrast, the strain *B.*

mucilaginosus 159 had more genes involved in amino acid metabolism (410 vs. 401 genes).

Annotation of genes in the KEGG Pathway database [89,90] demonstrated that strain 111 is capable of biosynthesizing B vitamins (thiamine, riboflavin, pantothenic acid, B₁₂), biotin, tetrahydrofolate, lipoic acid, and siderophores (bacillibactin). The siderophore bacillibactin




Fig. 1. Analysis of the cellular metabolic systems of the two *Bacillus* bacteria as performed using RAST 2.0 (<https://rast.nmp.dr.org/>) and antiSMASH 7.1.0 web service (<https://github.com/antismash/antismash/releases>). (A) *B. mucilaginosus* 159. (B) *B. subtilis* 111. To visualize the analysis results, the data were pre-displayed on a computer screen and saved as images using macOS Sequoia version 15 and the screenshot function. The screenshots were then edited using Adobe Photoshop (Adobe Inc., San Jose, CA, USA) version 26.1 to achieve the required quality.

is a catecholic trilactone secreted by various bacilli species for iron absorption during periods of iron limitation. Strain 159 could biosynthesize riboflavin, thiamine, and the siderophore bacillibactin.

In addition, the strains exhibited broad capabilities for the biosynthesis of fatty acids. We identified all the main enzymes responsible for forming fatty acids (C3—C18): FabD, FabF, FabG, FabZ, FabI (FabK, FabL), etc.

The genome of *B. subtilis* 111 contains genes for the biosynthesis of vitamin B₆: thiamine monophosphate kinase [EC:2.7.4.16], thiamine phosphate phos-

phatase [EC:3.1.3.100], and 18 genes involved in this process. Both strains contain genes for enzymes involved in the biosynthesis of riboflavin, as well as its coenzyme derivatives, i.e., flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD): riboflavin synthase [EC:2.5.1.9], riboflavin kinase [EC:2.7.1.26], FAD synthetase [EC:2.7.7.2], FMN reductase [EC:1.5.1.38] and FMN reductase [NAD(P)H] [EC:1.5.1.39], as well as other enzymes. Strain 111 contains a gene for purine nucleoside phosphorylase [EC:2.4.2.1] involved in the biosynthesis of nicotinamide and nicotinic acid. The biological role of pan-

tothenic acid is due to its participation in the biosynthesis of coenzyme A (CoA). The genome of strain 111 contains the enzyme pantoate- β -alanine ligase [EC:6.3.2.1] (catalyzes the biosynthesis of B₅ from pantoic acid and β -alanine) and pantothenate kinase [EC:2.7.1.33] (the first enzyme in the pathway of CoA biosynthesis). Pantothenate kinase phosphorylates pantothenate (vitamin B₅) to form 4'-phosphopantothenate using an adenosine triphosphate (ATP) molecule. Other genes were also identified in the biosynthesis of pantothenic acid from pyruvate, catalyzing its further conversion to CoA. The studied strain also possessed the biotin biosynthesis pathway and the key enzyme biotin synthase [EC:2.8.1.6].

3.2 Synthesis of Antimicrobial Compounds

Fig. 1 shows the metabolic capacities of the two strains to synthesize antimicrobial substances. According to the complete genomic sequence analysis, the *B. mucilaginosus* 159 strain can simultaneously synthesize several antimicrobial compounds of different natures and spectrums of action (Fig. 2A; data produced using antiSMASH 7.1.0 [91] and KAAS [89,90]). This analysis allowed us to identify sequences encoding biosynthesis of macrolactin H (an antibiotic from the polyketide group), mersacidin (a class II lanthipeptide), and bacilysin. The *B. subtilis* 111 strain genome (Fig. 2B) can synthesize andalusicin (class III lanthipeptide), bacilysin, the polyketide antibiotic macrolactin H, difficidin, and bacillaene (a polyene antibiotic).

Both strains could synthesize the antibiotic bacilysin (Fig. 3; data produced using KEGG Pathway database and KAAS [89,90]). The strains possessed all the genes controlling this process (gene cluster *bacABCDE*). Bacilysin is a non-ribosome-synthesized dipeptide antibiotic with an l-alanine residue at the N-terminus and a non-proteinogenic amino acid, l-anticapsin, at the C-terminus. Despite its simple structure, bacilysin is active against a wide range of bacteria and fungi, including *Candida albicans*. The antibacterial action of bacilysin depends mainly on its transport into host cells, its hydrolysis by intracellular peptidases to anticapsin, and inhibition of glucosamine-6-phosphate (GlcN6P) synthase by anticapsin.

3.3 Lipopeptide Synthesis

As follows from Fig. 1B, the *B. subtilis* strain 111 can synthesize fengycin (a lipopeptide with antifungal activity) and surfactin (another lipopeptide).

It was previously established that *Bacillus* spp. lipopeptides are small metabolites with a cyclic structure formed by 7–10 amino acids (including 2–4 D-amino acids) and a β -hydroxybutyric acid with 13–19 C atoms [97]. These lipopeptides have a variety of biological activities, including biofilm interactions and antifungal, anti-inflammatory, antitumor, antiviral, and antiplatelet properties. The multiple activities of lipopeptides have stimulated significant interest in the exploitation of these lipopep-

tides for use as antibiotics, feed additives, antitumor agents, acute thrombolytic therapeutics, and drug delivery systems. Hence, gaining knowledge of the inherent roles played by these structurally varied lipopeptides in *Bacillus* species is essential for effectively producing more potent products and understanding microbial regulatory systems. To increase the efficiency of their biosynthesis for industrial applications, ongoing efforts are required as there is currently a lack of knowledge regarding the direct target of these lipopeptides [97,98].

Surfactin is one of the powerful surfactants of biological origin and has a wide range of potential applications in industry and medicine. The main factor limiting its active usage is the high production cost. Due to the complexity of the structure, the chemical synthesis of surfactin is unprofitable, meaning microbiological synthesis is the main approach to its production [75]. Fengycin is an antifungal lipopeptide first discovered in the *B. subtilis* F-29-3 strain. Fengycin inhibits mycelial fungi but is ineffective against yeast and bacteria [99,100]. Thus, we may conclude that the *B. subtilis* strain 111 has a broader spectrum of activity, including fungicidal properties.

3.4 Osmotic Tolerance

The osmotic stability of strains may be crucial for developing biological starters for silage production. If the dry matter content in the grass is 30% or more, the increased osmotic pressure in plant cells consequently leads to inhibiting the development of lactic acid bacteria and causing a slowdown in the acidification of the grass mass, especially in the first, most crucial stage of its ensiling [101]. However, using osmotolerant bacteria in silage preparations helps overcome this problem [102–105]. Osmoprotectants, also known as compatible solutes, are tiny organic compounds that function as osmolytes to assist organisms in resisting excessive osmotic stress. Osmoprotectants have a neutral charge and are not hazardous at high concentrations. Overall, osmoprotectants can be divided into three chemical classes: (1) betaines and related molecules, (2) sugars and polyols, and (3) amino acids. These molecules accumulate in cells and balance the osmotic difference between the cell environment and the cytoplasm. There are two pathways for the accumulation of osmoprotectants in the cell: transport from the external environment and the biosynthesis of osmoprotectants inside the cell [106].

The uptake of osmotic stress protectants occurs via the Opu family of transporters, a system of central importance for managing osmotic stress in bacteria. The osmolarity of the environment controls opuD-mediated glycine betaine transport activity in *Bacillus* spp. High osmolarity maximizes glycine betaine absorption activity by activating pre-existing OpuD proteins and stimulating *de novo* OpuD synthesis. Strains have different mechanisms of tolerance to high osmotic pressure [107].




Fig. 2. Gene clusters involved in the biosynthesis of secondary metabolites that were determined using the antiSMASH 7.1.0 web service (<https://github.com/antismash/antismash/releases>) and KAAS (KEGG Automated Annotation Server; <https://www.genome.jp/kegg/kaas/>). (A) *B. mucilaginosus* 159. (B) *B. subtilis* 111. To visualize the analysis results, the data were pre-displayed on a computer screen and saved as images using macOS Sequoia version 15 and the screenshot function. The images were then edited using Adobe Photoshop version 26.1 to achieve the required quality.

Previously, bacilli were believed to be mainly capable of synthesizing only one osmoprotective substance within themselves, i.e., proline. However, one of the adaptation mechanisms that was identified in both studied *Bacillus* spp. strains (as shown in Fig. 4; data produced using KAAS

[89,90]) turned out to be the biosynthesis mechanism of glycine betaine, which is a very effective osmoprotectant that accumulates in high concentrations in response to increased osmotic pressure from choline under the action of the choline dehydrogenase (BetA(GbsB)) and betaine alde-




Fig. 3. Biosynthesis of the antibiotic bacilysin by the strains *B. subtilis* 111 and *B. mucilaginosus* 159 (according to the KEGG Pathway database (<https://www.genome.jp/kegg/pathway.html>) and using KAAS (KEGG Automated Annotation Server; <https://www.genome.jp/kegg/kaas/>)). To visualize the analysis results, the data were pre-displayed on a computer screen and saved as images using macOS Sequoia version 15 and the screenshot function. The images were then edited using Adobe Photoshop version 26.1 to achieve the required quality.




Fig. 4. Biosynthesis of glycine betaine by *B. subtilis* 111 strain and *B. mucilaginosus* 159 strain (according to the KEGG Pathway database; <https://www.genome.jp/kegg/pathway.html>). To visualize the analysis results, the data were pre-displayed on a computer screen and saved as images using macOS Sequoia version 15 and the screenshot function. The image was then edited using Adobe Photoshop version 26.1 to achieve the required quality.

hyde dehydrogenase (BetB) enzymes. Glycine betaine is synthesized in cells from choline in a two-step pathway un-

der the action of these two enzymes [108,109]. Among the compatible solutions used, glycine betaine plays a partic-

ularly important role for bacilli since it can be both synthesized and imported through three high-affinity and osmotically induced transport systems [107]. Glycine betaine importation is mediated by the betaine–choline–carnitine transporter (BCCT)-type transporter OpuD and the ABC transporters OpuA and OpuC [110].

Thus, the strains we studied here are characterized not only by their ability to pump osmoprotectants from the environment but also by their ability to synthesize proline and glycine betaine independently (Fig. 4).

Our findings provide comprehensive information about the metabolomic profiles of the two strains and the functional properties of the important proteins they are capable of synthesizing. Among them, we also highlight the genetic determinants for such a crucial metabolic system as response to stress factors (Table 2) [111–113]. The present study was performed in accordance with state-of-the-art investigations aimed at describing and defining a healthy gut based on important characteristics of microbial functionality that may be derived from microbiomics and its research methodologies [13,38,114]; these, in turn, benefit from omics and multi-omics approaches [19,22,46,47,115].

4. Conclusions

Analysis of the genomic sequences of *B. subtilis* 111 and *B. mucilaginosus* 159 demonstrated that these bacteria were similar in terms of their metabolomic profiles based on the sets of proteins they produce. Moreover, these strains did not contain antibiotic resistance determinants or virulence factors. Antimicrobial activity against Gram-positive and Gram-negative bacteria is among the important requirements for promising strains used as silage starters. Both strains can synthesize many antimicrobial substances of various natures and spectrums of action. The *B. mucilaginosus* 159 strain can synthesize macrolactin H (an antibiotic from the polyketide group), mersacidin (a class II lanthipeptide), and bacilysin. The *B. subtilis* 111 strain can synthesize andalusicin (class III lanthipeptide), bacilysin, polyketide antibiotic macrolactin H, difficidin, bacillaene (a polyene antibiotic), as well as fengycin (a lipopeptide with antifungal activity) and surfactin (another lipopeptide). Thus, the spectrum of biosynthesized antimicrobial substances is wider in the *B. subtilis* 111 strain. The osmotic stability of strains can also be a determining property for generating biological starters for silage preparation. The studied strains are characterized by the ability to pump osmoprotectants from the environment and synthesize them independently, both proline and glycine betaine. In general, it can be stated that these strains are safe for use as probiotic microorganisms or as starters for obtaining preserved feed. The *B. subtilis* 111 strain may be preferable due to a wider spectrum of synthesized antimicrobial substances and vitamins. Our study demonstrates the applicability of modern genomic technologies to characterize functional microbiomic and metabolomic features of such important bacte-

rial groups as *Bacillus* spp. and their subsequent use as antimicrobial feed additives.

Availability of Data and Materials

This article contains a presentation of all the experiments and findings from the study. Further details can be obtained upon request from the corresponding author.

Author Contributions

Conceptualization, GYL, LAI, VAF and EAY; methodology, LAI and VAF; software, ESP and EAB; validation, VAF, NIN and DKG; formal analysis, ESP; investigation, IAK, ASD, KAS and VAZ; resources, DGT and NIN; data curation, DGT and MNR; writing—original draft preparation, EAY and MNR; writing—review and editing, EAY, MNR and DKG; visualization, AVD, EAY and MNR; supervision, GYL and DKG; project administration, GYL and NIN; funding acquisition, LAI. All authors have read and agreed to the published version of the manuscript. All authors contributed to editorial changes in the manuscript. All authors have participated sufficiently in the work and agreed to be accountable for all aspects of the work.

Ethics Approval and Consent to Participate

Not applicable.

Acknowledgment

Not applicable.

Funding

The research was supported by the Russian Science Foundation grant No. 23-16-20007 (<https://rscf.ru/project/23-16-20007/>) and by the St. Petersburg Science Foundation grant No. 23-16-20007.

Conflict of Interest

The authors declare no conflicts of interest. Authors Valentina A. Filippova, Georgi Yu. Laptev, Larisa A. Ilina, Elena A. Yildirim, Evgeni A. Brazhnik, Kseniya A. Sokolova, and Irina A. Klyuchnikova are affiliated with BIOTROF Ltd., which sponsored this study. The study design, data analysis, interpretation, and manuscript preparation were carried out independently and were not influenced by the sponsor.

References

- [1] Boschiero M, De Laurentis V, Caldeira C, Sala S. Comparison of organic and conventional cropping systems: A systematic review of life cycle assessment studies. *Environmental Impact Assessment Review*. 2023; 102: 107187. <https://doi.org/10.1016/j.eiar.2023.107187>.
- [2] Callaway TR, Lillehoj H, Chuanchuen R, Gay CG. Alternatives to Antibiotics: A Symposium on the Challenges and Solutions

- for Animal Health and Production. *Antibiotics*. 2021; 10: 471. <https://doi.org/10.3390/antibiotics10050471>.
- [3] Bratishko NI, Pritulenko OV, Gaviley EV, Polyakova LL, Tereshchenko AV, Zhukorsky OM. Ratio ω -6 to ω -3 in PUFAs of compound feeds. *Ptitsevodstvo*. 2014; 9: 24–27. (In Russian with English summary). Available at: <https://www.researchgate.net/publication/342987874>. (Accessed: 10 November 2024).
 - [4] Bratishko NI, Polyakova LL, Prytulenko OV, Tereshchenko OV, Gaviley OV. The effect of an increased level of individual amino acids in compound feed with triticale on the metabolism in the body of chickens. *Ptahivnictvo*. 2011; 67: 37–43. (In Ukrainian). Available at: <https://www.researchgate.net/publication/342816297>. (Accessed: 10 November 2024).
 - [5] Zhukorskyi O, Prytulenko O, Tereshchenko O, Bratishko N. The presence of soybean, triticale and sorghum in compound feed for poultry is fully justified. *Zerno i khlib*. 2012; 2: 14–16. (In Ukrainian). Available at: <https://www.researchgate.net/publication/342832208>. (Accessed: 10 November 2024).
 - [6] Laptev G, Tyurina D, Yildirim E, Gorfunkel E, Ilina L, Filippova V, et al. Effects of glyphosate and antibiotics on the expression of genes related to performance, antioxidant protection and histological barrier in the cecum of broilers. In Ronzhin A, Bakach M, Kostyaev A (eds.) *Smart Innovation, Systems and Technologies, Proceedings of the Fourth International Conference on Agriculture Digitalization and Organic Production (ADOP 2024)*, Minsk, Belarus, June 05–08, 2024 (pp. 337–345). Springer Nature Singapore Pte Ltd. Singapore. 2024. https://doi.org/10.1007/978-981-97-4410-7_28.
 - [7] Iqbal B, Li G, Alabbosh KF, Hussain H, Khan I, Tariq M, et al. Advancing environmental sustainability through microbial reprogramming in growth improvement, stress alleviation, and phytoremediation. *Plant Stress*. 2023; 10: 100283. <https://doi.org/10.1016/j.stress.2023.100283>.
 - [8] Nadarajah K, Abdul Rahman NS. The microbial connection to sustainable agriculture. *Plants*. 2023; 12: 2307. <https://doi.org/10.3390/plants12122307>.
 - [9] Jagadeesan Y, Meenakshisundaram S, Pichaimuthu S, Balaiah A. A scientific version of understanding “Why did the chickens cross the road”? – A guided journey through *Bacillus* spp. towards sustainable agriculture, circular economy and biofortification. *Environmental Research*. 2024; 244: 117907. <https://doi.org/10.1016/j.envres.2023.117907>.
 - [10] Wallace RJ, Newbold CJ. Probiotics for ruminants. In Fuller R. (ed.) *The Scientific Basis* (pp. 317–353). Springer, Dordrecht: UK. 1992. https://doi.org/10.1007/978-94-011-2364-8_12.
 - [11] McCann JC, Elolimy AA, Loor JJ. Rumen microbiome, probiotics, and fermentation additives. *Veterinary Clinics: Food Animal Practice*. 2017; 33: 539–553. <https://doi.org/10.1016/j.cvfa.2017.06.009>.
 - [12] Artursson V. Bacterial-Fungal Interactions Highlighted using Microbiomics: Potential Application for Plant Growth Enhancement. Doctoral Dissertation. Swedish University of Agricultural Sciences: Uppsala. 2005. Available at: <http://pub.epsilon.slu.se/id/file/1572>. (Accessed: 10 November 2024).
 - [13] Sommer MOA. Functional Microbiomics: Uncovering Novel Reservoirs of Microbial Machinery for Antibiotic and Biomass Inhibitor Processing. Harvard University: Cambridge, MA, USA. 2008. Available at: <https://search.proquest.com/openview/9cf12267cd61ec063a36e22ed142a917/1?pq-origsite=gscholar&cbl=18750&diss=y>. (Accessed: 10 November 2024).
 - [14] El Jeni R, Villot C, Koyun OY, Osorio-Doblado A, Baloyi JJ, Lourenco JM, et al. *Invited review*: “Probiotic” approaches to improving dairy production: Reassessing “magic foo-foo dust”. *Journal of Dairy Science*. 2024; 107: 1832–1856. <https://doi.org/10.3168/jds.2023-23831>.
 - [15] Kholif AE, Anele A, Anele UY. Microbial feed additives in ruminant feeding. *AIMS Microbiology*. 2024; 10: 542–571. <https://doi.org/10.3934/microbiol.2024026>.
 - [16] Pavlov DS, Egorov IA, Nekrasov RV, Laktionov KS, Kravtsova LZ, Pravdin VG, et al. The use of feed additives produced by solid state microbial fermentation, to increase nutritional value of feeds and level of meals and cakes in concentrates. *Problemy biologii produktivnyh zivotnyh*. 2011; 1: 89–92. (In Russian with English summary). Available at: <http://archive.today/2025.03.13-100316/https://www.elibrary.ru/item.asp?id=16092172>. (Accessed: 10 November 2024).
 - [17] Seo JK, Kim SW, Kim MH, Upadhaya SD, Kam DK, Ha JK. Direct-fed microbials for ruminant animals. *Asian-Australasian Journal of Animal Sciences*. 2010; 23: 1657–1667. <https://doi.org/10.5713/ajas.2010.r.08>.
 - [18] Tarakanov BV, Nikolicheva TA, Aleshin VV. Probiotiki. Dostizheniya i perspektivy ispol'zovaniya v zhivotnovodstve. Proshloye, nastoyashcheye i budushcheye zootekhnicheskoy nauki: Nauchnyye trudy VIZha. 2004; 3: 69–73. (In Russian). Available at: <https://scholar.google.com/scholar?hl=en&q=Probiotics.+Achievements+and+prospects+of+use+in+animal+husbandry>. (Accessed: 10 November 2024).
 - [19] Susanti D, Volland A, Tawari N, Baxter N, Gangaiah D, Plata G, et al. Multi-Omics characterization of host-derived *Bacillus* spp. probiotics for improved growth performance in poultry. *Frontiers in Microbiology*. 2021; 12: 747845. <https://doi.org/10.3389/fmicb.2021.747845>.
 - [20] Vlasatikova L, Zeman M, Crhanova M, Matiasovicova J, Karasova D, Faldynova M, et al. Colonization of chickens with competitive exclusion products results in extensive differences in metabolite composition in cecal digesta. *Poultry Science*. 2023; 103: 103217. <https://doi.org/10.1016/j.psj.2023.103217>.
 - [21] Kayal A, Stanley D, Radovanovic A, Horyanto D, Van TT, Bajagai YS. Controlled intestinal microbiota colonisation in broilers under the industrial production system. *Animals*. 2022; 12: 3296. <https://doi.org/10.3390/ani12233296>.
 - [22] Wang R, Lin F, Ye C, Aihemaitijiang S, Halimulati M, Huang X, et al. Multi-omics analysis reveals therapeutic effects of *Bacillus subtilis*-fermented *Astragalus membranaceus* in hyperuricemia via modulation of gut microbiota. *Food Chemistry*. 2023; 399: 133993. <https://doi.org/10.1016/j.foodchem.2022.133993>.
 - [23] Kochish II, Smolensky VI, Laptev GY, Romanov MN, Nikonov IN, Panin AN, et al. [Methodical Recommendations on the Implementation of the Developed System for the Prevention of Pathogen Bacteria by Correcting Diets in Laying Hens and Using Antimicrobial Additives]. *Sel'skokhozyaistvennyye tekhnologii: Moscow*. 2018. (In Russian). Available at: <https://kar.kent.ac.uk/75283>. (Accessed: 10 November 2024).
 - [24] Fathima S, Hakeem WG, Shanmugasundaram R, Selvaraj RK. Necrotic enteritis in broiler chickens: a review on the pathogen, pathogenesis, and prevention. *Microorganisms*. 2022; 10: 1958. <https://doi.org/10.3390/microorganisms10101958>.
 - [25] Kim GB, Seo YM, Kim CH, Paik IK. Effect of dietary prebiotic supplementation on the performance, intestinal microflora, and immune response of broilers. *Poultry Science*. 2011; 90: 75–82. <https://doi.org/10.3382/ps.2010-00732>.
 - [26] Huyghebaert G, Ducatelle R, Van Immerseel F. An update on alternatives to antimicrobial growth promoters for broilers. *The Veterinary Journal*. 2011; 187: 182–188. <https://doi.org/10.1016/j.tvjl.2010.03.003>.
 - [27] Yildirim E, Ilina L, Laptev G, Tyurina D, Filippova V, Dubrovin A, et al. The search for sources of enterobacteria and clostridia endotoxins in Russian dairy farms: possible transfer of endotoxins through the feed-cow-milk chain. In Ronzhin A, Bakach M, Kostyaev A (eds.) *Smart Innovation, Systems and Technologies*,

- Proceedings of the Fourth International Conference on Agriculture Digitalization and Organic Production (ADOP 2024), Minsk, Belarus, June 05-08, 2024 (pp. 325–335). Springer Nature Singapore Pte Ltd. Singapore. 2024. https://doi.org/10.1007/978-981-97-4410-7_27.
- [28] Singh J, Sharma S, Nara S. Evaluation of gold nanoparticle based lateral flow assays for diagnosis of enterobacteriaceae members in food and water. *Food Chemistry*. 2015; 170: 470–483. <https://doi.org/10.1016/j.foodchem.2014.08.092>.
- [29] Maifreni M, Frigo F, Bartolomeoli I, Innocente N, Biasutti M, Marino M. Identification of the Enterobacteriaceae in Montasio cheese and assessment of their amino acid decarboxylase activity. *Journal of Dairy Research*. 2013; 80: 122–127. <https://doi.org/10.1017/S002202991200074X>.
- [30] Bover-Cid S, Hugas M, Izquierdo-Pulido M, Vidal-Carou MC. Amino acid-decarboxylase activity of bacteria isolated from fermented pork sausages. *International Journal of Food Microbiology*. 2001; 66: 185–189. [https://doi.org/10.1016/S0168-1605\(00\)00526-2](https://doi.org/10.1016/S0168-1605(00)00526-2).
- [31] Rooke JA, Hatfield RD. Biochemistry of ensiling. In Buxton DR, Muck RE, Harrison JH (eds.) *Silage Science and Technology, Agronomy Monographs* (pp. 95–139). American Society of Agronomy, Inc., Crop Science Society of America, Inc., Soil Science Society of America, Inc., John Wiley & Sons, Inc.: Hoboken, NJ, USA. 2003. <https://doi.org/10.2134/agronmonogr42.c3>.
- [32] Ciorba MA. A gastroenterologist’s guide to probiotics. *Clinical gastroenterology and hepatology: the Official Clinical Practice Journal of the American Gastroenterological Association*. 2012; 10: 960–968. <https://doi.org/10.1016/j.cgh.2012.03.024>.
- [33] National Institutes of Health. Probiotics: fact sheet for health professionals. National Institutes of Health: Bethesda, MD, USA. 2023. Available at: <https://ods.od.nih.gov/factsheets/Probiotics-HealthProfessional/>. (Accessed: 10 November 2024).
- [34] Romanov MN, Grozina AA, Ilina LA, Laptev GY, Yildirim EA, Filippova VA, *et al.* From feed regulation to regulated feeding: intestinal microbiome and performance optimization in broiler chickens in response to antibiotic and probiotic treatment. In *Life of Genomes 2022: Abstracts of the International Conference, Kazan, Russia, 23–24 November 2022* (pp. 44–45). Research Center “Regulatory Genomics”, Institute of Fundamental Medicine and Biology, Kazan (Volga Region) Federal University: Kazan, Russia. 2022. Available at: <http://lifeofgenomes.r-genomics.com/wp-content/uploads/2022/12/Abstracts.pdf#page=45>. (Accessed: 10 November 2024).
- [35] Chandrasekaran P, Weiskirchen S, Weiskirchen R. Effects of probiotics on gut microbiota: An overview. *International Journal of Molecular Sciences*. 2024; 25: 6022. <https://doi.org/10.3390/ijms25116022>.
- [36] Grozina AA, Ilina LA, Laptev GY, Yildirim EA, Ponomareva ES, Filippova VA, *et al.* Probiotics as an alternative to antibiotics in modulating the intestinal microbiota and performance of broiler chickens. *Journal of Applied Microbiology*. 2023; 134: lxad213. <https://doi.org/10.1093/jambio/lxad213>.
- [37] Xiao L, Zhang C, Zhang X, Zhao X, Mahsa GC, Ma K, *et al.* Effects of *Lactocaseibacillus paracasei* SNB-derived postbiotic components on intestinal barrier dysfunction and composition of gut microbiota. *Food Research International*. 2024; 175: 113773. <https://doi.org/10.1016/j.foodres.2023.113773>.
- [38] Laptev G, Turina D, Yildirim E, Ilina L, Gorfunkel E, Filippova V, *et al.* Analysis of changes in broiler microbiome biodiversity parameters due to intake of glyphosate and probiotic *Bacillus* sp. GL-8 using next-generation sequencing. In Ronzhin A, Kostyaev A (eds.) *Smart Innovation, Systems and Technologies, Proceedings of the Third International Conference on Agriculture Digitalization and Organic Production (ADOP 2023)*, St. Petersburg, Russia, 5–7 June 2023 (pp. 161–170). Springer Nature Singapore Pte Ltd. Singapore. 2023. https://doi.org/10.1007/978-981-99-4165-0_15.
- [39] De Vos WM. Microbiomics of probiotic and other intestinal bacteria. *Annals of Nutrition and Metabolism*. 2007. Available at: <https://www.jstor.org/stable/48507901>. (Accessed: 10 November 2024).
- [40] Laptev G, Yildirim E, Ilina L, Ponomareva E, Kalitkina K, Turina D, *et al.* Effect of a probiotic strain administration in different feeding phases on α - and β -diversity and gene expression of the rumen microbiome in lactating cows. In Ronzhin A, Kostyaev A (eds.) *Smart Innovation, Systems and Technologies, Proceedings of the Third International Conference on Agriculture Digitalization and Organic Production (ADOP 2023)*, St. Petersburg, Russia, 5–7 June 2023 (pp. 181–191). Springer Nature Singapore Pte Ltd. Singapore. 2023. https://doi.org/10.1007/978-981-99-4165-0_17.
- [41] Rajendhran J, Gunasekaran P. Human microbiomics. *Indian Journal of Microbiology*. 2010; 50: 109–112. <https://doi.org/10.1007/s12088-010-0034-9>.
- [42] Jernberg C. Use of Microbiomics to Study Human Impacts on Complex Microbial Communities. Doctoral Dissertation. Karolinska institutet: Stockholm. 2006. Available at: <https://www.diva-portal.org/smash/record.jsf?pid=diva2:1072771>. (Accessed: 10 November 2024).
- [43] Jagadeesan B, Gerner-Smidt P, Allard MW, Leuillet S, Winkler A, Xiao Y, *et al.* The use of next generation sequencing for improving food safety: Translation into practice. *Food Microbiology*. 2019; 79: 96–115. <https://doi.org/10.1016/j.fm.2018.11.005>.
- [44] Billington C, Kingsbury JM, Rivas L. Metagenomics approaches for improving food safety: a review. *Journal of Food Protection*. 2022; 85: 448–464. <https://doi.org/10.4315/JFP-21-301>.
- [45] Rajendhran J, Gunasekaran P. Human genomics and microbiomics: the postgenomics scenario. *Current Science*. 2009. Available at: <https://www.researchgate.net/publication/287626551>. (Accessed: 10 November 2024).
- [46] Palazzotto E, Weber T. Omics and multi-omics approaches to study the biosynthesis of secondary metabolites in microorganisms. *Current Opinion in Microbiology*. 2018; 45: 109–116. <https://doi.org/10.1016/j.mib.2018.03.004>.
- [47] Singh D, Geat N, Mehriya M, Rajawat MV, Prasanna R, Kumar A, *et al.* Omics (genomics, proteomics, metabolomics, etc.) tools to study the environmental microbiome and bioremediation. In Kashyap BK, Solanki MK, Kamboj DV, Pandey AK (eds.) *Waste to Energy: Prospects and Applications* (pp. 235–260). Springer: Singapore. 2020. Available at: https://doi.org/10.1007/978-981-33-4347-4_10. (Accessed: 10 November 2024).
- [48] Chai R, Huo R, Tao C, Qiu H, Shui X, Yin H, *et al.* A multi-omics approach reveals the changes in the gut microbiome and metabolism of large yellow croaker (*Larimichthys crocea*) by dietary supplementation with *Bacillus*. *Aquaculture*. 2025; 595: 741711. <https://doi.org/10.1016/j.aquaculture.2024.741711>.
- [49] Saladié M, Caparrós-Martín JA, Agudelo-Romero P, Wark PAB, Stick SM, O’Gara F. Microbiomic analysis on low abundant respiratory biomass samples; improved recovery of microbial DNA from bronchoalveolar lavage fluid. *Frontiers in Microbiology*. 2020; 11: 572504. <https://doi.org/10.3389/fmicb.2020.572504>.
- [50] Chang L, Zhao T, Zhao C, Zhu W, Xu L, Liu J, *et al.* Microbiomic and transcriptomic insight into the pathogenesis of meningitis-like disease in cultured *Pelophylax nigromaculatus*. *Aquaculture*. 2021; 530: 735736. <https://doi.org/10.1016/j.aquaculture.2020.735736>

- [51] Broderick D, Marsh R, Waite D, Pillarisetti N, Chang AB, Taylor MW. Realising respiratory microbiome meta-analyses: time for a standardised framework. *Microbiome*. 2023; 11: 57. <https://doi.org/10.1186/s40168-023-01499-w>.
- [52] Park J, Jung SY, Kim HY, Lee KE, Go YJ, Kim HS, *et al.* Microbiomic association between the saliva and salivary stone in patients with sialolithiasis. *Scientific Reports*. 2024; 14: 9184. <https://doi.org/10.1038/s41598-024-59546-x>.
- [53] Wang Y, Su W, Liu Z, Wang Y, Li L, Xu H, *et al.* The microbiomic signature of hemorrhoids and comparison with associated microbiomes. *Frontiers in Microbiology*. 2024; 15: 1329976. <https://doi.org/10.3389/fmicb.2024.1329976>.
- [54] Guan H, Zhao S, Tan Y, Fang X, Zhang Y, Zhang Y, *et al.* Microbiomic insights into the oral microbiome's role in type 2 diabetes mellitus: standardizing approaches for future advancements. *Frontiers in Endocrinology*. 2024; 15: 1416611. <https://doi.org/10.3389/fendo.2024.1416611>.
- [55] Dictionary.com. Microbiomic. Dictionary.com, LLC. 2024. Available at: <https://www.dictionary.com/browse/microbiomic> (Accessed: 10 November 2024).
- [56] Paleri P. Microbiomic Security (Microbiomicsec)(m_{s2}). Revisiting National Security: Prospecting Governance for Human Well-Being (pp. 989–1015). Springer Nature: Singapore. 2022. https://doi.org/10.1007/978-981-16-8293-3_24.
- [57] Hoa NT, Baccigalupi L, Huxham A, Smertenko A, Van PH, Ammendola S, *et al.* Characterization of *Bacillus* species used for oral bacteriotherapy and bacteriophylaxis of gastrointestinal disorders. *Applied and Environmental Microbiology*. 2000; 66: 5241–5247. <https://doi.org/10.1128/AEM.66.12.5241-5247.2000>.
- [58] Urdaci MC, Bressollier P, Pinchuk I. *Bacillus clausii* probiotic strains: antimicrobial and immunomodulatory activities. *Journal of Clinical Gastroenterology*. 2004; 38: S86–S90. <https://doi.org/10.1097/01.mcg.0000128925.06662.69>.
- [59] Lefevre M, Racedo SM, Ripert G, Housez B, Cazaubiel M, Maudet C, *et al.* Probiotic strain *Bacillus subtilis* CU1 stimulates immune system of elderly during common infectious disease period: a randomized, double-blind placebo-controlled study. *Immunity & Ageing: I & A*. 2015; 12: 24. <https://doi.org/10.1186/s12979-015-0051-y>.
- [60] Kapse NG, Engineer AS, Gowdaman V, Wagh S, Dhakephalkar PK. Functional annotation of the genome unravels probiotic potential of *Bacillus coagulans* HS243. *Genomics*. 2019; 111: 921–929. <https://doi.org/10.1016/j.ygeno.2018.05.022>.
- [61] Sulthana A, Lakshmi SG, Madempudi RS. Genome sequencing and annotation of *Bacillus subtilis* UBBS-14 to ensure probiotic safety. *Journal of Genomics*. 2019; 7: 14–17. <https://doi.org/10.7150/jgen.31170>.
- [62] Oh JK, Pajarillo EA, Chae JP, Kim IH, Yang DS, Kang DK. Effects of *Bacillus subtilis* CSL2 on the composition and functional diversity of the faecal microbiota of broiler chickens challenged with *Salmonella* Gallinarum. *Journal of Animal Science and Biotechnology*. 2017; 8: 1. <https://doi.org/10.1186/s40104-016-0130-8>.
- [63] Trufanov O, Kotyk A, Trufanova V, Tereshchenko O, Zhukorskiy O. Detection of antibiotics, active against *Bacillus subtilis*, in grain and feed. *Agricultural Science and Practice*. 2015; 2: 60–66. <https://doi.org/10.15407/agrisp.2.01.060>.
- [64] Bernardeau M, Lehtinen MJ, Forssten SD, Nurminen P. Importance of the gastrointestinal life cycle of *Bacillus* for probiotic functionality. *Journal of Food Science and Technology*. 2017; 54: 2570–2584. <https://doi.org/10.1007/s13197-017-2688-3>.
- [65] Shleeva MO, Kondratieva DA, Kaprelyants AS. *Bacillus licheniformis*: a producer of antimicrobial substances, including Antimycobacterials, which are feasible for medical applications. *Pharmaceutics*. 2023; 15: 1893. <https://doi.org/10.3390/pharma15071893>.
- [66] Blanco Crivelli X, Cundon C, Bonino MP, Sanin MS, Bentancor A. The complex and changing genus *Bacillus*: A diverse bacterial powerhouse for many applications. *Bacteria*. 2024; 3: 256–270. <https://doi.org/10.3390/bacteria3030017>.
- [67] Markelova N, Chumak A. Antimicrobial activity of *Bacillus* cyclic lipopeptides and their role in the host adaptive response to changes in environmental conditions. *International Journal of Molecular Sciences*. 2025; 26: 336. <https://doi.org/10.3390/ijms26010336>.
- [68] Sorokulova I. Modern status and perspectives of *Bacillus* bacteria as probiotics. *Journal of Probiotics & Health*. 2013; 1: e106. <https://doi.org/10.4172/2329-8901.1000e106>.
- [69] Nishioka T, Matsuda K, Fujita Y. Combined analysis of metabolome and transcriptome: catabolism in *Bacillus subtilis*. In Tomita M, Nishioka T (eds.) *Metabolomics: The Frontier of Systems Biology* (pp. 127–140). Springer: Tokyo. 2005. https://doi.org/10.1007/4-431-28055-3_9.
- [70] Perez-Fons L, Bramley PM, Fraser PD. The optimisation and application of a metabolite profiling procedure for the metabolic phenotyping of *Bacillus* species. *Metabolomics*. 2014; 10: 77–90. <https://doi.org/10.1007/s11306-013-0553-6>.
- [71] Sauer U, Cameron DC, Bailey JE. Metabolic capacity of *Bacillus subtilis* for the production of purine nucleosides, riboflavin, and folic acid. *Biotechnology and Bioengineering*. 1998; 59: 227–238. [https://doi.org/10.1002/\(SICI\)1097-0290\(19980720\)59:2<227::AID-BIT10>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1097-0290(19980720)59:2<227::AID-BIT10>3.0.CO;2-B).
- [72] Gu Q, Li P. Biosynthesis of vitamins by probiotic bacteria. In Rao V, Rao LG (eds.) *Probiotics and Prebiotics in Human Nutrition and Health* (pp. 135–148). IntechOpen: London. 2016. <https://doi.org/10.5772/63117>.
- [73] Zhang M, Zhao X, Chen X, Li M, Wang X. Enhancement of riboflavin production in *Bacillus subtilis* via in vitro and in vivo metabolic engineering of pentose phosphate pathway. *Biotechnology Letters*. 2021; 43: 2209–2216. <https://doi.org/10.1007/s10529-021-03190-2>.
- [74] Senchenkov VY, Lyakhovchenko NS, Nikishin IA, Myagkov DA, Chepurina AA, Polivtseva VN, *et al.* Whole-genome sequencing and biotechnological potential assessment of two bacterial strains isolated from poultry farms in Belgorod, Russia. *Microorganisms*. 2023; 11: 2235. <https://doi.org/10.3390/microrganisms11092235>.
- [75] Trefilov VS, Labanov VA, Khrenova MG, Panova TV, Rodin VA, Savitskaya VY, *et al.* Genomic characterization of *Bacillus subtilis* PY79 and NCIB 3610 as potential producers of surfactin. *Applied Biochemistry and Microbiology*. 2024; 60: 1543–1550. <https://doi.org/10.1134/S0003683824700145>.
- [76] Illumina. Specification sheet: MiSeq System™ – Speed and simplicity for targeted resequencing and small-genome sequencing. (M-GL-00006 v4.0). Illumina, Inc., San Diego, CA, USA. 2021. Available at: <https://www.illumina.com/content/dam/illumina/gcs/assembled-assets/marketing-literature/miseq-system-data-sheet-m-gl-00006/miseq-data-sheet-m-gl-00006.pdf> (Accessed: 10 November 2024).
- [77] Shanmuganandam S, Schwessinger B, Hall R. Library preparation protocol to sequence V3-V4 region of 16S rRNA to run in Illumina MiSeq platform. Protocol Integer ID: 26943. protocols.io. 2019. Available at: <https://www.protocols.io/view/library-preparation-protocol-to-sequence-v3-v4-reg-6i7hchn.html> (Accessed: 10 November 2024).
- [78] Lapev GY, Yildirim EA, Ilina LA, Filippova VA, Kochish II, Gorfunkel EP, Dubrovin AV, *et al.* Effects of essential oils-based supplement and *Salmonella* infection on gene expression, blood parameters, cecal microbiome, and egg production in laying hens. *Animals*. 2021; 11: 360. <https://doi.org/10.3390/ani11020360>.

- [79] Kai S, Matsuo Y, Nakagawa S, Kryukov K, Matsukawa S, Tanaka H, *et al.* Rapid bacterial identification by direct PCR amplification of 16S rRNA genes using the MinION™ nanopore sequencer. *FEBS Open Bio.* 2019; 9: 548–557. <https://doi.org/10.1002/2211-5463.12590>.
- [80] Baer M, Höppe L, Seel W, Lipski A. Impact of DNA extraction, PCR amplification, sequencing, and bioinformatic analysis on food-associated mock communities using PacBio long-read amplicon sequencing. *BMC Microbiology.* 2024; 24: 521. <https://doi.org/10.1186/s12866-024-03677-8>.
- [81] Andrews S. FastQC: A Quality Control Tool for High throughput Sequence Data, Version 0.10.1. Bioinformatics Group, Babraham Institute: Cambridge, UK. 2012. Available at: <http://www.bioinformatics.babraham.ac.uk/projects/fastqc> (Accessed: 10 November 2024).
- [82] Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics.* 2014; 30: 2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- [83] Nurk S, Bankevich A, Antipov D, Gurevich AA, Korobeynikov A, Lapidus A, *et al.* Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *Journal of Computational Biology.* 2013; 20: 714–737. <https://doi.org/10.1089/cmb.2013.0084>.
- [84] Pruitt K, Brown G, Tatusova T, Maglott D. The Reference Sequence (RefSeq) Database. 2002 Oct 9 [Updated 2012 Apr 6]. In McEntyre J, Ostell J (eds.) *The NCBI Handbook*. National Center for Biotechnology Information: Bethesda, MD, USA, 2002. Chapter 18. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK21091/> (Accessed: 10 November 2024).
- [85] Kunst F, Ogasawara N, Moszer I, Albertini AM, Alloni G, Azevedo V, *et al.* The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature.* 1997; 390: 249–256. <https://doi.org/10.1038/36786>.
- [86] Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, *et al.* The RAST Server: rapid annotations using subsystems technology. *BMC Genomics.* 2008; 9: 75. <https://doi.org/10.1186/1471-2164-9-75>.
- [87] Meyer F, Overbeek R, Rodriguez A. FIGfams: yet another set of protein families. *Nucleic Acids Research.* 2009; 37: 6643–6654. <https://doi.org/10.1093/nar/gkp698>.
- [88] Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics.* 2014; 30: 2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.
- [89] Ogata H, Goto S, Fujibuchi W, Kanehisa M. Computation with the KEGG pathway database. *Biosystems.* 1998; 47: 119–128. [https://doi.org/10.1016/S0303-2647\(98\)00017-3](https://doi.org/10.1016/S0303-2647(98)00017-3).
- [90] Kanehisa Laboratories. KEGG: Kyoto Encyclopedia of Genes and Genomes. Kanehisa Laboratories: Fukuoka—Kyoto—Tokyo, Japan. 1995–2024. Available at: <https://www.genome.jp/kegg/> (Accessed: 10 November 2024).
- [91] Blin K, Shaw S, Kloosterman AM, Charlop-Powers Z, Van Wezel GP, Medema MH, *et al.* antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic Acids Research.* 2021; 49: W29–W35. <https://doi.org/10.1093/nar/gkab335>.
- [92] Kim S, Cheng T, He S, Thiessen PA, Li Q, Gindulyte A, *et al.* PubChem protein, gene, pathway, and taxonomy data collections: bridging biology and chemistry through target-centric views of PubChem data. *Journal of Molecular Biology.* 2022; 434: 167514. <https://doi.org/10.1016/j.jmb.2022.167514>.
- [93] Fliissi A, Dufresne Y, Michalik J, Tonon L, Janot S, Noé L, *et al.* Norine, the knowledgebase dedicated to non-ribosomal peptides, is now open to crowdsourcing. *Nucleic Acids Research.* 2016; 44: D1113–D1118. <https://doi.org/10.1093/nar/gkv1143>.
- [94] Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, *et al.* PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Research.* 2016; 44: W16–W21. <https://doi.org/10.1093/nar/gkw387>.
- [95] Kleinheinz KA, Joensen KG, Larsen MV. Applying the ResFinder and VirulenceFinder web-services for easy identification of acquired antibiotic resistance and *E. coli* virulence genes in bacteriophage and prophage nucleotide sequences. *Bacteriophage.* 2014; 4: e27943. <https://doi.org/10.4161/bact.27943>.
- [96] Bortolaia V, Kaas RS, Ruppe E, Roberts MC, Schwarz S, Couto V, *et al.* ResFinder 4.0 for predictions of phenotypes from genotypes. *Journal of Antimicrobial Chemotherapy.* 2020; 75: 3491–3500. <https://doi.org/10.1093/jac/dkaa345>.
- [97] Zhao H, Shao D, Jiang C, Shi J, Li Q, Huang Q, *et al.* Biological activity of lipopeptides from *Bacillus*. *Applied Microbiology and Biotechnology.* 2017; 101: 5951–5960. <https://doi.org/10.1007/s00253-017-8396-0>.
- [98] Saiyam D, Dubey A, Malla MA, Kumar A. Lipopeptides from *Bacillus*: unveiling biotechnological prospects—sources, properties, and diverse applications. *Brazilian Journal of Microbiology.* 2024; 55: 281–295. <https://doi.org/10.1007/s42770-023-01228-3>.
- [99] Vanittanakom N, Loeffler W, Koch U, Jung G. Fengycin—a novel antifungal lipopeptide antibiotic produced by *Bacillus subtilis* F-29-3. *The Journal of Antibiotics.* 1986; 39: 888–901. <https://doi.org/10.7164/antibiotics.39.888>.
- [100] Sur S, Romo TD, Grossfield A. Selectivity and mechanism of fengycin, an antimicrobial lipopeptide, from molecular dynamics. *The Journal of Physical Chemistry B.* 2018; 122: 2219–2226. <https://doi.org/10.1021/acs.jpcc.7b11889>.
- [101] Pobednov YuA, Kosolapov VM. Biological bases of grass silage and haylage making (review). *Sel'skokhozyaistvennaya biologiya.* 2014; 2: 31–41. (In Russian with English summary). <https://doi.org/10.15389/agrobiol.2014.2.31eng>.
- [102] Rooke JA. Lactic acid bacteria and silage fermentations. *Journal of Chemical Technology & Biotechnology.* 1991; 51: 560–562. <https://doi.org/10.1002/jctb.280510418>.
- [103] Elferink SJWHO, Driehuis F, Gottschal JC, Spoelstra SF. Silage fermentation processes and their manipulation. *FAO Plant Production and Protection Paper.* 2000. Available at: <http://www.fao.org/4/x8486e/x8486e09.htm#m9>. (Accessed: 10 November 2024).
- [104] Gaucher F, Bonnassie S, Rabah H, Leverrier P, Pottier S, Jardin J, *et al.* Benefits and drawbacks of osmotic adjustment in *Propionibacterium freudenreichii*. *Journal of Proteomics.* 2019; 204: 103400. <https://doi.org/10.1016/j.jprot.2019.103400>.
- [105] Romanov MN, Bato RV, Yokoyama MT, Rust SR. PCR detection and 16S rRNA sequence-based phylogeny of a novel *Propionibacterium acidipropionici* applicable for enhanced fermentation of high moisture corn. *Journal of Applied Microbiology.* 2004; 97: 38–47. <https://doi.org/10.1111/j.1365-2672.2004.02282.x>.
- [106] Hoffmann T, Wensing A, Brosius M, Steil L, Völker U, Bremer E. Osmotic control of *opuA* expression in *Bacillus subtilis* and its modulation in response to intracellular glycine betaine and proline pools. *Journal of Bacteriology.* 2013; 195: 510–522. <https://doi.org/10.1128/jb.01505-12>.
- [107] Nau-Wagner G, Opper D, Rolbetzki A, Boch J, Kempf B, Hoffmann T, *et al.* Genetic control of osmoadaptive glycine betaine synthesis in *Bacillus subtilis* through the choline-sensing and glycine betaine-responsive GbsR repressor. *Journal of Bacteriology.* 2012; 194: 2703–2714. <https://doi.org/10.1128/jb.06642-11>.
- [108] Boch J, Kempf B, Schmid R, Bremer E. Synthesis of the osmoprotectant glycine betaine in *Bacillus subtilis*: characterization of the gbsAB genes. *Journal of Bacteriology.* 1996; 178: 5121–5129. <https://doi.org/10.1128/jb.178.17.5121-5129.1996>.
- [109] Rath H, Sappa PK, Hoffmann T, Gesell Salazar M, Reder A,

- Steil L, *et al.* Impact of high salinity and the compatible solute glycine betaine on gene expression of *Bacillus subtilis*. *Environmental Microbiology*. 2020; 22: 3266–3286. <https://doi.org/10.1111/1462-2920.15087>.
- [110] Kappes RM, Kempf B, Kneip S, Boch J, Gade J, Meier-Wagner J, *et al.* Two evolutionarily closely related ABC transporters mediate the uptake of choline for synthesis of the osmoprotectant glycine betaine in *Bacillus subtilis*. *Molecular Microbiology*. 1999; 32: 203–216. <https://doi.org/10.1046/j.1365-2958.1999.01354.x>.
- [111] Chebotar IV, Emelyanova MA, Bocharova JA, Mayansky NA, Kopantseva EE, Mikhailovich VM. The classification of bacterial survival strategies in the presence of antimicrobials. *Microbial Pathogenesis*. 2021; 155: 104901. <https://doi.org/10.1016/j.micpath.2021.104901>.
- [112] Amarnath K, Narla AV, Pontrelli S, Dong J, Reddan J, Taylor BR, *et al.* Stress-induced metabolic exchanges between complementary bacterial types underly a dynamic mechanism of inter-species stress resistance. *Nature Communications*. 2023; 14: 3165. <https://doi.org/10.1038/s41467-023-38913-8>.
- [113] Kochish II, Romanov MN, Myasnikova OV, Nikonov IN, Selina MV, Korenyuga MV, *et al.* [Methodical Recommendations on the Use of Antimicrobial Feed Additive for the Prevention of Stress in Industrial Crosses of Laying Hens]. *Sel'skokhozyaistvennyye tekhnologii: Moscow*. 2021. (In Russian). Available at: <https://kar.kent.ac.uk/92646> (Accessed: 10 November 2024).
- [114] Egert M, de Graaf AA, Smidt H, de Vos WM, Venema K. Beyond diversity: Functional microbiomics of the human colon. *Trends in Microbiology*. 2006; 14: 86–91. <https://doi.org/10.1016/j.tim.2005.12.007>.
- [115] Ma T, Shen X, Shi X, Sakandar HA, Quan K, Li Y, *et al.* Targeting gut microbiota and metabolism as the major probiotic mechanism - An evidence-based review. *Trends in Food Science & Technology*. 2023; 138: 178–198. <https://doi.org/10.1016/j.tifs.2023.06.013>.