



## Short Communications (Research Advances)

## Zircon U-Pb age evidence of the Mesoarchean (2.9–3.2 Ga) crustal remnant in the Southern Dabie Orogen, South China

Da-liang Xu<sup>a, b</sup>, Lian-hong Peng<sup>b, \*</sup>, Xin Deng<sup>a, b</sup>, Chao-ran Liu<sup>b, c</sup>, Le Wan<sup>d</sup>, Xiao-ming Zhao<sup>e</sup><sup>a</sup> Research Center for Petrogenesis and Mineralization of Granitoid Rocks, China Geological Survey, Wuhan 430205, China<sup>b</sup> Wuhan Center of China Geological Survey (China Central South China Innovation Center for Geosciences), Ministry of Natural Resources, Wuhan 430205, China<sup>c</sup> China University of Geosciences, Beijing 100083, China<sup>d</sup> School of Earth Science, China University of Geoscience, Wuhan 430074, China<sup>e</sup> Institute of Karst Geology, CAGS, Guilin 541004, China

## 1. Objective

Unlike the North China Plate where Archean and Paleoproterozoic crustal rocks are widely distributed, Early Precambrian basement rocks in the Yangtze Block of South China are locally exposed (Fig. 1a), such as in the Kongling Complex, the Zhongxiang Complex, and the Douling Complex (Zhao GC and Cawood PA, 2012). The Dabie Orogen is the eastward extension of the Qinling Orogen, which formed during the northward subduction of the Yangtze Block beneath the North China Plate in the Triassic. Archean rocks are only locally exposed in the Northern Dabie Orogen, such as the 2.77 Ga Huangtuling granulites within a granitic migmatitic gneiss (Wu YB et al., 2008) and the 2.85 Ga Tuanfeng migmatites (Qiu XF et al., 2020), suggesting the presence of ancient crustal rocks in the Northern Dabie Orogen. In this study, the authors present the results of a combined study of zircon U-Pb ages, trace elements, and Lu-Hf isotope compositions for a newly discovered Paleoproterozoic granitic gneiss with Mesoarchean remnants from the Qichun area in the Southern Dabie Orogen. The research results will provide a new constraint on the Precambrian crustal evolution of the Northern Yangtze Block.

## 2. Methods

Zircons from the granitic gneiss sample (D966–1) were

separated by standard heavy-liquid and magnetic techniques, followed by handpicking under a binocular microscope. After polishing and carbon coating, the internal structure of the zircon grains was evaluated by cathode luminescence (CL) imaging conducted at the Nanjing Hongchuang Geological Exploration Technology Service Co. Ltd. Zircon U-Pb dating and trace element analyses were conducted using an icapQ ICP-MS connected to a RESolution laser-ablation system at the Laboratory of Isotope Geochemistry, Wuhan Center of China Geological Survey. Zircon 91500 was used as a U-Pb standard and NIST610 was used to calibrate U, Th, and Pb contents. A laser spot diameter of 29 μm was used with a repetition frequency of 4 Hz. *In-situ* Lu-Hf isotope analyses of zircons were measured at the same laboratory using a Neptune Plus MC-ICP-MS combined with a RESolution laser-ablation system. The laser spot was 44 μm in diameter with an ablation frequency of 8 Hz. The detailed analytical conditions and procedures are the same as those described by Qiu XF et al. (2020).

## 3. Results


The Dabie complex in the Southern Dabie Orogen is mainly composed of metamorphosed supracrustal rocks and granitic gneisses. A sample (D966–1) was collected from an outcrop of granitic gneiss which is over thrust onto a mica schist (Fig. 1c). They predominantly consist of quartz, plagioclase, and biotite, with minor zircon and apatite as accessory minerals, and show a gneissic texture (Fig. 1d). Zircon grains from the granitic gneiss sample are subhedral to euhedral in morphology, 70–120 μm in length with aspect ratios of 1 : 1 to 3 : 1, and commonly show clear oscillatory zoning (Fig. 2a), typical of an igneous origin. A total of 25 zircon domains were analyzed for U-Pb dating and/or Lu-Hf

First author: E-mail address: [xdl2003geo@163.com](mailto:xdl2003geo@163.com) (Da-liang Xu).\* Corresponding author: E-mail address: [245737309@qq.com](mailto:245737309@qq.com) (Lian-hong Peng).


Literary editor: Li-qiong Jia

doi:10.31035/cg2022056

2096-5192/© 2023 China Geology Editorial Office.



**Fig. 1.** Geological map of South China showing the distribution of Early Precambrian rocks in the Yangtze Block (a, modified from Zhao GC and Cawood PA, 2012), simplified geological map (b), field outcrop (c), and microscope photograph (d) of the discovered granitic gneiss in the Qichun area. Qtz–quartz, Pl–plagioclase.



**Fig. 2.** CL images (a), chondrite-normalized REE patterns (b), U-Pb age diagrams (c), and U-Pb age versus  $\epsilon_{Hf}(t)$  values (d) of representative zircons from granitic gneiss (D966-1) in the Qichun area.

isotope (Supplementary Tables S1–S2), and they yielded apparent  $^{207}\text{Pb}/^{206}\text{Pb}$  ages that varied from  $3276 \pm 93$  Ma to  $2000 \pm 17$  Ma. Based on CL images, trace element compositions, and U-Pb ages, the zircon domains can be categorized into three groups (Fig. 2), as follows.

Group I zircon domains yielded similar U ( $72 \times 10^{-6}$ – $114 \times 10^{-6}$ ) and Th ( $89 \times 10^{-6}$ – $148 \times 10^{-6}$ ), higher Th/U ratios (1.02–1.54), negative Eu anomalies ( $\text{Eu}^* = 0.18$ – $0.26$ ), and steep HREE patterns with small ranges in  $(\text{Yb}/\text{Gd})_{\text{N}}$  ratios of 6.2–11.2. They show apparent  $^{207}\text{Pb}/^{206}\text{Pb}$  ages of  $2000 \pm 17$  Ma to  $2015 \pm 26$  Ma (Supplementary Table S1), with a weighted mean of  $2007 \pm 18$  Ma (MSWD=0.16), indicative of the emplacement age of the magmatic protolith. The zircon domains ( $n=5$ ) have  $^{176}\text{Hf}/^{177}\text{Hf}$  ratios of 0.281225–0.281245 and  $^{176}\text{Lu}/^{177}\text{Hf}$  ratios of 0.000304–0.000500. Their  $\varepsilon\text{Hf}(t)$  values range from  $-9.7$  to  $-10.3$  with two-stage depleted mantle Hf model ages ( $T_{\text{DM2}}$ ) of 3222–3258 Ma (Supplementary Table S2).

Group II zircon domains are relict magmatic zircons. They yielded variable U ( $31 \times 10^{-6}$ – $373 \times 10^{-6}$ ) and Th ( $15 \times 10^{-6}$ – $224 \times 10^{-6}$ ) contents, lower Th/U ratios (0.22–0.74), negative Eu anomalies ( $\text{Eu}^* = 0.59$ – $0.82$ ), and steeper HREE patterns with  $(\text{Yb}/\text{Gd})_{\text{N}}$  ratios of 4.4–13.3. The  $^{207}\text{Pb}/^{206}\text{Pb}$  ages for these domains ranged from  $2945 \pm 13$  Ma to  $2828 \pm 16$  Ma (Supplementary Table S1), with an upper intercept of  $2919 \pm 21$  Ma (MSWD=5.0) and a weighted mean of  $2922 \pm 14$  Ma (MSWD=2.6) (Fig. 2c). Group II domains ( $n=10$ ) exhibit lower  $^{176}\text{Hf}/^{177}\text{Hf}$  ratios of 0.280838–0.280944 and variable  $^{176}\text{Lu}/^{177}\text{Hf}$  ratios of 0.000214–0.001472. Their  $\varepsilon\text{Hf}(t)$  values range from  $-0.6$  to  $-3.2$  with  $T_{\text{DM2}}$  ages of 3377–3534 Ma (Supplementary Table S2).

Group III zircon domains are also relict magmatic zircons. They are discordant, likely due to later metamorphic or alteration events, and have apparent  $^{207}\text{Pb}/^{206}\text{Pb}$  ages of  $3276 \pm 93$  to  $3208 \pm 13$  Ma (Supplementary Table S1). Four zircons define an upper intercept ages of  $3236 \pm 48$  Ma (MSWD=2.9) (Fig. 2c). These domains show variable contents of U ( $248 \times 10^{-6}$ – $704 \times 10^{-6}$ ) and Th ( $133 \times 10^{-6}$ – $189 \times 10^{-6}$ ), with Th/U ratios of 0.19–0.77. They exhibit negative Eu anomalies with  $\text{Eu}^* = 0.53$ – $0.68$  and steep HREE patterns with small ranges in  $(\text{Yb}/\text{Gd})_{\text{N}}$  ratios of 11.7–16.7. These domains ( $n=2$ ) have low  $^{176}\text{Hf}/^{177}\text{Hf}$  ratios of 0.280637–0.280700 and  $^{176}\text{Lu}/^{177}\text{Hf}$  ratios of 0.000942–0.001011. Their  $\varepsilon\text{Hf}(t)$  values are  $-2.0$  to  $-4.7$  with  $T_{\text{DM2}}$  ages 3705–3848 Ma (Supplementary Table S2).

#### 4. Conclusion

Relict magmatic zircons with Mesoarchean U-Pb ages of about 2.9 Ga and about 3.2 Ga were discovered for the first time from a granitic gneiss in the Qichun area in the Southern

Dabie Orogen. They show negative  $\varepsilon\text{Hf}(t)$  values ranging from  $-0.6$  to  $-3.2$  and  $-2.0$  to  $-4.7$ , with two-stage depleted mantle Hf model ages ( $T_{\text{DM2}}$ ) of 3.3–3.5 Ga and 3.7–3.8 Ga, respectively, indicating the existence of Mesoarchean crustal remnants in the Southern Dabie Orogen and the growth of juvenile crust during the Eoarchean to Paleoarchean. The final emplacement age for the protolith of the studied granitic gneiss is about 2007 Ma, which is consistent with the Paleoproterozoic magmatic events in the Kongling and Zhongxiang complexes of the Yangtze Block, with a possible link to the assembly of the supercontinent Columbia (Zhao GC and Cawood PA, 2012). Their two-stage depleted Hf model ages of 3.2 Ga suggest derivation through the episodic reworking of the Archean crust. This study demonstrates the presence of Mesoarchean crustal remnants in the Southern Dabie Orogen and the reworking of Archean crust during the Paleoproterozoic.

#### CRedit authorship contribution statement

Da-liang Xu, Lian-hong Peng and Xin Deng conceived of the presented idea. Da-liang Xu and Chao-ran Liu carried out the experiment. All authors discussed the results and contributed to the final manuscript.

#### Declaration of competing interest

The authors declare no conflicts of interest.

#### Acknowledgment

This work was financially supported by the Open Fund of the Research Center for Petrogenesis and Mineralization of Granitoid Rocks (PMGR202014) and the project of the China Geological Survey (DD20190050).

#### Supplementary dataset

Supplementary data (Supplementary Tables S1–S2) to this article can be found online at doi: [10.31035/cg2022056](https://doi.org/10.31035/cg2022056).

#### References

- Qiu XF, Jiang T, Wu NW, Zhao XM, Xu Q. 2020. Neoproterozoic crustal rocks and Paleoproterozoic migmatization in the Dabie Orogen: Evidence from zircon U-Pb age and Hf isotope. *Acta Geologica Sinica*, 94(3), 729–738 (in Chinese with English abstract).
- Wu YB, Zheng YF, Gao S, Jiao WF, Liu YS. 2008. Zircon U-Pb age and trace element evidence for Paleoproterozoic granulite-facies metamorphism and Archean crustal rocks in the Dabie Orogen. *Lithos*, 101, 308–322. doi: [10.1016/j.lithos.2007.07.008](https://doi.org/10.1016/j.lithos.2007.07.008).
- Zhao GC, Cawood PA. 2012. Precambrian geology of China. *Precambrian Research*, 222, 13–54. doi: [10.1016/j.precamres.2012.09.017](https://doi.org/10.1016/j.precamres.2012.09.017).