

# H<sub>2</sub>O<sub>2</sub> accumulation shapes the differential recovery from nitrogen and magnesium deficiency in cucumber

Qi ZHOU, Kehao CHEN, Ying WANG, Xusheng MENG, Shiyu WANG (✉), Shiwei GUO

Key Lab of Organic-based Fertilizers of China and Jiangsu Provincial Key Lab for Solid Organic Waste Utilization, Nanjing Agricultural University, Nanjing 210095, China.


## KEYWORDS

Nitrogen deficiency, magnesium deficiency, cucumber, antioxidant system, photosynthesis, H<sub>2</sub>O<sub>2</sub>

## HIGHLIGHTS

- Nitrogen and magnesium deficiencies differentially impair cucumber growth, with Mg deficiency causing more severe damage and only partial recovery upon nutrient resupply.
- It was discovered that H<sub>2</sub>O<sub>2</sub> plays a crucial role in the recovery of cucumbers from nitrogen and magnesium deficiency.
- The excessive accumulation of H<sub>2</sub>O<sub>2</sub> under magnesium deficiency surpasses the antioxidant capacity, resulting in irreversible cellular damage. This highlights the significance of magnesium fertilizer in modern agricultural production.
- Application of the H<sub>2</sub>O<sub>2</sub> scavenger DMTU alleviates photoinhibition and partially restores photosynthetic function, providing direct evidence that H<sub>2</sub>O<sub>2</sub> underlies the differential recovery from N and Mg deficiency.

## GRAPHICAL ABSTRACT



## ABSTRACT

Nitrogen (N) and magnesium (Mg) deficiencies involve distinct underlying damage mechanisms and recovery responses. In this study, the physiological and biochemical responses of hydroponically grown cucumber plants to N or Mg deficiency and their recovery after nutrient resupply were evaluated. Considering that these deficiencies induce oxidative stress, we evaluated alterations in the antioxidant system and the accumulation of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). The results revealed that both N deficiency and Mg deficiency inhibited cucumber growth and biomass production, with more pronounced adverse effects observed in leaves under Mg-deficient conditions. N resupply effectively restored leaf morphology, gas exchange parameters, and photosynthetic efficiency. In contrast, Mg resupply resulted in only partial recovery. This limited recovery can be attributed to irreversible damage to the photosynthetic system caused by the inability of antioxidant enzymes to effectively scavenge excessive amounts of reactive oxygen species (ROS), indicating that Mg deficiency triggers H<sub>2</sub>O<sub>2</sub>-induced cellular damage.

Received December 12, 2025;

Accepted March 11, 2026.

Correspondence: minwang@njau.edu.cn

Special Issue: Magnesium Nutrition in Soil–Plant Systems

Furthermore, the application of the H<sub>2</sub>O<sub>2</sub> scavenger DMTU (N,N'-dimethylthiourea) significantly reduced H<sub>2</sub>O<sub>2</sub> levels under both deficiency conditions, particularly under Mg-deficient conditions. These findings demonstrate that N deficiency and Mg deficiency have distinct effects on antioxidant responses and underscore the pivotal role of H<sub>2</sub>O<sub>2</sub> in plant responses to nutrient deficiency stress and subsequent recovery. These findings suggest that targeted modulation of H<sub>2</sub>O<sub>2</sub> levels may serve as a promising strategy to increase crop resilience, offering a clear direction for future research.

© The Author(s) 2027. Published by Higher Education Press. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0>)

## 1 Introduction

Nitrogen (N) and magnesium (Mg) are essential macronutrients that play critical roles in the physiological and biochemical processes of plants<sup>[1,2]</sup>. Typically, N constitutes approximately 1% to 5% of the total plant dry matter, while Mg constitutes approximately 0.15% to 0.35%<sup>[3]</sup>. Most previous studies have focused on the role of N in plant physiological processes, particularly its involvement in photosynthesis through the regulation of chlorophyll content, carbon fixation enzyme activity, and the photorespiration rate<sup>[4]</sup>. Moreover, N serves as a structural component of nucleic acids, proteins, phytohormones, and secondary metabolites; therefore, N deficiency often results in an increased root-to-shoot ratio<sup>[5]</sup>. Like N, Mg plays a crucial role in photosynthetic processes and is also essential for carbon partitioning. As both a structural component of the chlorophyll molecule and an activator of various photosynthesis-related enzymes—such as carboxylases—Mg contributes significantly to plant metabolic functions<sup>[6]</sup>. In addition, sucrose phloem loading is closely associated with the proton gradient across the plasma membranes of phloem cells; consequently, any reduction in Mg-ATP levels may significantly impair the function of H<sup>+</sup>-ATPase<sup>[7,8]</sup>.

N-deficient plants typically exhibit growth inhibition and develop characteristic yellowing or reduced leaf expansion. In response to limited N availability in the soil, plants have evolved specific adaptive mechanisms to maintain growth. For example, root growth is often promoted under N deficiency to increase N uptake<sup>[9]</sup>. Plants deficient in Mg typically exhibit interveinal chlorosis in their older leaves, along with a significant reduction in the root-to-shoot ratio. The pattern of change in the root-to-shoot ratio under Mg deficiency is

opposite to that observed under N deficiency, likely because of the nonsubstitutable role of Mg in assimilate transport and physiological function<sup>[10]</sup>. A reduction in the photosynthetic rate is a commonly observed response in plants under N or Mg deficiency, which can be attributed to two primary factors. First, the leaf chlorophyll content decreases because of reduced uptake of N or Mg. Both types of nutrients are highly mobile within plants and are preferentially reallocated to younger leaves to support active growth under conditions of nutrient limitation<sup>[11,12]</sup>. Second, N or Mg starvation induces a decrease in the activity of photosynthesis-related enzymes, including ribulose-1,5-bisphosphate carboxylase (Rubisco), a key rate-limiting enzyme in CO<sub>2</sub> assimilation<sup>[13,14]</sup>. In general, impaired CO<sub>2</sub> fixation leads to overreduction of the photosynthetic electron transport chain, resulting in excessive accumulation of reactive oxygen species (ROS) and subsequent damage to chloroplasts and cellular structures<sup>[15]</sup>. Although the mechanisms underlying leaf senescence under N deficiency and Mg deficiency are similar, a notable distinction in agricultural practices is that N deficiency is typically reversible upon resupply, whereas Mg deficiency often results in irreversible crop damage<sup>[10,16]</sup>. The fundamental cause of this differential recovery remains unclear.

Previous studies have demonstrated that N deficiency results in a significant reduction in the activities of superoxide dismutase (SOD), guaiacol peroxidase, and catalase (CAT) in wheat stems during the mid-grain filling stage, accompanied by a marked increase in ROS concentrations<sup>[17]</sup>. Mg deficiency results in substantial carbohydrate accumulation in leaves, a response that differs markedly from that associated with N deficiency<sup>[18]</sup>. Moreover, Mg deficiency-induced accumulation of sugars leads to feedback inhibition of photosynthesis, which may increase the production of ROS. Consequently, compared with plants

under N-deficient conditions, plants under Mg-deficient conditions may experience greater ROS-mediated damage, potentially resulting in irreversible cellular injury. Under stress conditions, plants generate various types of ROS, including hydrogen peroxide ( $\text{H}_2\text{O}_2$ ), superoxide radical ( $\text{O}_2^-$ ), hydroxyl radical ( $\cdot\text{OH}$ ), and singlet oxygen ( $^1\text{O}_2$ )<sup>[19]</sup>.  $\text{H}_2\text{O}_2$ , the most abundant and stable ROS, plays a crucial role in various key plant resistance mechanisms<sup>[20]</sup>.  $\text{H}_2\text{O}_2$  is proposed to function as a signaling molecule involved in the regulation of various stress responses<sup>[21]</sup>. Although elevated levels of ROS can induce oxidative damage to cellular structures, previous studies have focused primarily on plant responses to nutrient deficiency stress. Researchers have paid little attention to the specific function of reactive ROS, especially  $\text{H}_2\text{O}_2$ , in this process. It remains uncertain whether recovery from N or Mg deficiency is influenced by reactive oxygen species metabolism.

Owing to the differences in the physiological functions of N and Mg, as well as the distinct responses of plants to N deficiency and Mg deficiency, in contrast to N deficiency, Mg deficiency leads to irreversible damage once visible symptoms appear. A previous study reported that leaf senescence induced by N deficiency was reversed through N resupply, as N-sensitive regulatory pathways effectively and coordinately regulate plant growth<sup>[22,23]</sup>. In contrast, Mg resupply had limited effectiveness in reducing leaf damage, as sucrose phloem export had already been severely impaired prior to the onset of leaf senescence. Previous studies have focused primarily on plant responses to nutrient deficiency stresses, with limited attention given to the recovery mechanisms following nutrient resupply. The potential involvement of ROS, particularly  $\text{H}_2\text{O}_2$ , in N deficiency and Mg deficiency and the related differential recovery responses has not yet been fully elucidated.

To investigate the physiological responses to nutrient deficiency and subsequent resupply, cucumber plants were cultivated hydroponically under five different treatment conditions: a control group with full nutrient supply, N deficiency, Mg deficiency, N resupply following N deficiency, and Mg resupply following Mg deficiency. Parameters such as plant growth, chlorosis, biomass, photosynthesis, elemental concentration, antioxidant system activity, and  $\text{H}_2\text{O}_2$  production were systematically evaluated. This study was designed to (1) assess the physiological effects of N deficiency or Mg deficiency on cucumber plants; (2) elucidate the distinct recovery patterns following N resupply and Mg resupply; and

(3) determine the specific role of  $\text{H}_2\text{O}_2$  in mediating the recovery processes from these nutrient deficiencies. This facilitates a more comprehensive understanding of the disparities in the physiological indicators of plants under N deficiency and Mg deficiency, as well as the disparities in the recovery processes of plants from these two types of nutrient deficiency. Comprehending these disparities offers a methodology for alleviating nutrient deficiency stress and resuming production in agricultural practices.

## 2 Materials and methods

### 2.1 Plant materials and experimental design

In this study, cucumber (*Cucumis sativus* L.) cultivar “Jinchun 4” seeds were soaked in distilled water for 1 h and germinated on moistened gauze at 25 °C in the dark for 48 h. Following germination, the seeds were sown in quartz sand and cultivated in a greenhouse under controlled conditions. The greenhouse was maintained under controlled environmental conditions, including a stable photosynthetic photon flux density (PPFD) of 500  $\mu\text{mol photons m}^{-2}\cdot\text{s}^{-1}$ , a 14-h photoperiod, a constant relative humidity of approximately 50%, and a day/night temperature of 28 °C/20 °C. When the seedlings reached the three-leaf stage, uniform individuals were transplanted into 1 L plastic pots filled with one-quarter strength nutrient solution. Three days after transplantation, the nutrient solution concentration was increased to half the maximum strength; another three days later, it was replaced with full-strength nutrient solution, which was maintained until the conclusion of the experiment. The nutrient solution was continuously aerated throughout the growth period. The full-strength nutrient solution contained the following essential elements: 1.25  $\text{mmol}\cdot\text{L}^{-1}$   $(\text{NH}_4)_2\text{SO}_4$ , 1.25  $\text{mmol}\cdot\text{L}^{-1}$   $\text{Ca}(\text{NO}_3)_2$ , 2.5  $\text{mmol}\cdot\text{L}^{-1}$   $\text{K}_2\text{SO}_4$ , 1.0  $\text{mmol}\cdot\text{L}^{-1}$   $\text{KH}_2\text{PO}_4$ , 2.0  $\text{mmol}\cdot\text{L}^{-1}$   $\text{MgSO}_4$ , 35.8  $\mu\text{mol}\cdot\text{L}^{-1}$  Fe-EDTA, 57.8  $\mu\text{mol}\cdot\text{L}^{-1}$   $\text{H}_3\text{BO}_3$ , 11.4  $\mu\text{mol}\cdot\text{L}^{-1}$   $\text{MnCl}_2$ , 0.96  $\mu\text{mol}\cdot\text{L}^{-1}$   $\text{ZnSO}_4$ , 0.4  $\mu\text{mol}\cdot\text{L}^{-1}$   $\text{CuSO}_4$ , and 0.48  $\mu\text{mol}\cdot\text{L}^{-1}$   $\text{H}_2\text{MoO}_4$ . The nutrient solution was renewed every three days, and the pH was adjusted daily to  $6.0 \pm 0.1$  using 0.1  $\text{mol}\cdot\text{L}^{-1}$  HCl or 0.1  $\text{mol}\cdot\text{L}^{-1}$  NaOH.

Experiment 1: Seedlings were subjected to different nutrient treatments upon full expansion of the fourth leaf (d0). Nutrient stress was imposed by removing either Mg or N from the nutrient solution. The experimental groups included a Mg-deficient group (-Mg), a Mg-deficient group for 9 days

followed by a Mg resupply for 5 days (–Mg+), a N-deficient group (–N), a N-deficient group for 9 days followed by a N resupply for 5 days (–N+), and a control group with a full supply of both Mg and N (CK). Nutrient deprivation was initiated on Day 0 (d0). On Day 9 (d9), two additional treatments were introduced: half of the Mg-deficient and N-deficient cucumber plants were selected and supplemented with Mg or N, respectively, and then cultivated for an additional 5 days, referred to as [(d9) + 5]. On Day 0 (d0), the seedlings were supplied with a standard nutrient solution, a Mg-deficient solution (0 mmol·L<sup>-1</sup> MgSO<sub>4</sub>), or a N-deficient solution [0 mmol·L<sup>-1</sup> (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and 0 mmol·L<sup>-1</sup> Ca(NO<sub>3</sub>)<sub>2</sub>] for a duration of 9 days (d9). In the N deficiency treatment, calcium was supplied as CaCl<sub>2</sub> (1.25 mmol·L<sup>-1</sup>) to replace Ca(NO<sub>3</sub>)<sub>2</sub>, ensuring equivalent Ca<sup>2+</sup> availability. On Day 9 (d9), half of the Mg-deficient and N-deficient plants were resupplied with 2 mmol·L<sup>-1</sup> Mg or 5 mmol·L<sup>-1</sup> N, respectively.

Experiment 2: When the fourth leaves fully expanded, 2.5 μmol·L<sup>-1</sup> N,N'-dimethylthiourea (DMTU) was applied via foliar spraying. After a three-day interval, normal nutrient supply, Mg-deficient, and N-deficient treatments were initiated. The seedlings were supplied with nutrient solutions according to the protocol described in Experiment 1.

## 2.2 Thermal imaging and microscopic observation

Nine days after the initiation of Experiment 1 (d9), thermal images of the entire plant were acquired using an infrared camera (SC620, FLIR Systems, Inc., Wilsonville, OR, USA). Imaging was conducted between 10:00 a.m. and 11:00 a.m., during which time the stomatal openings were confirmed to be fully active. Digital thermograms were analyzed using Thermal CAM Researcher Professional 2.9 software (FLIR Systems, Inc.).

The leaf abaxial epidermal microstructure was examined under a light microscope (Olympus IX71; Olympus Optical Co., Ltd., Tokyo, Japan) at a magnification of 400×.

## 2.3 Photosynthesis and chlorophyll fluorescence measurements

Five days after the resupply of Mg or N [(d9) + 5], gas exchange and photosynthetic efficiency parameters were measured on the upper fully expanded leaves using a portable

photosynthesis system equipped with a fluorescence leaf chamber (LI-6400; LI-COR Biosciences, Lincoln, NE, USA). To measure the net photosynthetic rate (P<sub>n</sub>), stomatal conductance (g<sub>s</sub>), intercellular CO<sub>2</sub> concentration (C<sub>i</sub>), and transpiration rate (T<sub>r</sub>), the leaves were maintained at a temperature of 28 °C and a relative humidity of approximately 50% under a PPFD of 1200 μmol photons m<sup>-2</sup>·s<sup>-1</sup>. Data were recorded after the system reached steady-state equilibrium, typically within approximately 15 min. Following the gas exchange measurements, the minimal fluorescence (F<sub>o</sub>) and maximal fluorescence (F<sub>m</sub>) of leaves were measured under dark-adapted conditions between 2:00 a.m. and 3:00 a.m. The variable fluorescence (F<sub>v</sub>) is determined by calculating the difference between the maximal fluorescence (F<sub>m</sub>) and the minimal fluorescence (F<sub>o</sub>).

The maximum quantum efficiency of photosystem II (F<sub>v</sub>/F<sub>m</sub>) was determined as follows:

$$F_v/F_m = \frac{F_m - F_o}{F_m} \quad (1)$$

## 2.4 Determination of biomass, element concentrations and chlorophyll content

Following the measurement of photosynthesis, five plants from each treatment were randomly selected for the determination of plant height and root length. Each plant was subsequently separated into three distinct fractions: leaves, sheaths, and roots. The dry matter of each fraction was determined by oven drying at 105 °C for 30 min, followed by drying at 70 °C until a constant weight was achieved. Dried leaf samples were predigested overnight using a HNO<sub>3</sub>-HClO<sub>4</sub> mixture (4:1, v/v) and then subjected to digestion on a heating plate with the temperature gradually increasing from 60 to 190 °C over a period of 4 h. The concentrations of various ions (P, K, Ca, Mg, Fe, Mn, Cu, Zn, B, and Mo) in the resulting digestate were analyzed using ICP-MS (NexION 300X, PerkinElmer Inc., Waltham, MA, USA). A separate portion of the dried plant material was digested with a H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O<sub>2</sub> mixture at 270 °C for 1 h, and the N concentration in the digest was quantified using an AutoAnalyzer 3 Digital Colorimeter (Bran + Luebbe Inc., Norderstedt, Germany).

## 2.5 Quantitative determination of hydrogen peroxide and antioxidant enzyme levels

The H<sub>2</sub>O<sub>2</sub> content was determined using a commercially

available assay kit (Nanjing Jiancheng Bioengineering Institute). Fresh leaf tissue (0.2 g) was rapidly frozen in liquid nitrogen and homogenized in 2 mL of ice-cold 0.1% trichloroacetic acid (TCA). The homogenate was subsequently centrifuged at  $19000 \times g$  for 20 min at 4 °C. Afterward, 0.5 mL of the supernatant was mixed with 0.5 mL of potassium phosphate buffer ( $100 \text{ mmol}\cdot\text{L}^{-1}$ , pH 7.8) and 2 mL of potassium iodide solution ( $1 \text{ mol}\cdot\text{L}^{-1}$ ). The reaction mixture was incubated in the dark for 1 h to ensure complete development. The absorbances of both the samples and the calibration standards were measured at 390 nm using a spectrophotometer. The  $\text{H}_2\text{O}_2$  concentration was quantified on the basis of a calibration curve generated from a series of standard solutions provided with the kit.

The activities of SOD and peroxidase (POD) were determined spectrophotometrically by measuring changes in absorbance. Fresh leaf tissue (0.2 g) was rapidly frozen in liquid nitrogen and then homogenized in 2 mL of ice-cold phosphate buffer ( $50 \text{ mmol}\cdot\text{L}^{-1}$ , pH 7.8) containing 1% insoluble polyvinylpyrrolidone (PVPP) and  $0.2 \text{ mmol}\cdot\text{L}^{-1}$  ethylenediaminetetraacetic acid (EDTA). The homogenate was centrifuged at  $10000 \times g$  for 15 min at 4 °C, and 0.5 mL of the resulting supernatant was collected for enzymatic activity assays. SOD activity was assayed by monitoring the photochemical reduction of nitroblue tetrazolium (NBT) at 560 nm according to the method described by Stewart and

Bewley<sup>[24]</sup>. One unit of SOD activity was defined as the 50% inhibition of NBT reduction. POD activity was determined spectrophotometrically by measuring the oxidation of guaiacol at 470 nm, following the method described by Wang et al<sup>[25]</sup>.


## 2.6 Statistical analysis

Statistical analyses were conducted using SPSS statistical software. One-way analysis of variance (ANOVA) was employed to assess differences in the measured parameters across various treatment groups, with Duncan's multiple comparison test applied to identify significant differences among group means. A *P* value of less than 0.05 was considered to indicate statistical significance.

## 3 Results

### 3.1 Effect of Mg or N deficiency on cucumber plant growth

Following a 9-day period of nutrient deficiency cultivation (d9), cucumber seedlings grown in nutrient solutions lacking either Mg or N presented visible foliage deficiency symptoms (Fig. 1(a)). Under Mg deficiency, interveinal chlorosis developed, which is a characteristic symptom of Mg deficiency. This symptom intensified over time, particularly in older



**Fig. 1** Effects of Mg or N deficiency on cucumber growth (d9). (a) Growth status of cucumber plants; (b) dry weight of cucumber plants. Cucumber plants were cultivated under three treatments: normal nutrient solution (CK), Mg-deficient conditions (-Mg), and N-deficient conditions (-N) for a period of 9 days. The data are presented as the means and standard deviations from five independent replicates. Different lowercase letters indicate statistically significant differences ( $P < 0.05$ ) according to Duncan's multiple range test, with comparisons made between treatments for the same plant organ.


leaves, as the duration of Mg deficiency increased. Similarly, N deficiency led to leaf chlorosis and inhibited overall plant growth (Fig. 1(a)). N-deficient plants displayed shorter shoots and reduced shoot growth rates, whereas root growth remained largely unaffected. In contrast, compared with that in the control plants, shoot growth in the Mg-deficient plants was comparable to that in the control, but compared with that in the control plants, root growth in the former decreased. When deficiency symptoms became apparent, total dry matter accumulation was not significantly affected by Mg deficiency; however, N deficiency resulted in an average reduction in whole-plant dry matter of approximately 16% (Fig. 1(b)).

Thermal imaging and microscopic observations revealed

distinct effects of Mg deficiency and N deficiency on leaf physiology. N deficiency led to a significant increase in leaf temperature, as shown in Fig. 2. In contrast, Mg deficiency resulted in elevated temperatures primarily between the veins of middle leaves. Microscopic analysis further demonstrated that visible cell necrosis occurred exclusively under Mg-deficient conditions (Fig. 2(a)).

### 3.2 Effects of Mg or N resupply on the performance and photosynthetic parameters of cucumber plants

Five days after nutrient supplementation [(d9) + 5], the Mg resupply increased plant height and root and stem biomass (Table 1) and partially alleviated deficiency symptoms in the



**Fig. 2** Effects of Mg or N deficiency on the leaf temperature of cucumber plants (d9). (a) Digital color reflectance images, thermal images, and microscopy images of cucumber leaves following treatment with a standard nutrient solution (CK), Mg-deficient solution (-Mg), and N-deficient solution (-N) for 9 days; (b) leaf temperature measurements in cucumber plants. The data are presented as the means and standard deviations from five independent replicates. Different lowercase letters indicate statistically significant differences ( $P < 0.05$ ) according to Duncan's multiple range test.

**Table 1** Effects of Mg or N resupply on the growth of Mg or N deficient cucumber plants [(d9) + 5]

Treatment	Plant height (cm)	Root length (cm)	Dry weight (g)				Root/Shoot
			Total	Root	Stem	Leaf	
CK	65.3 ± 5.1a	40.5 ± 2.1a	3.25 ± 0.06a	0.32 ± 0.03a	1.34 ± 0.06a	1.58 ± 0.05a	0.11 ± 0.01b
-Mg	55.5 ± 1.3b	37.7 ± 2.6a	2.46 ± 0.10c	0.18 ± 0.01c	0.82 ± 0.06cd	1.46 ± 0.04b	0.08 ± 0.01c
-Mg+	63.3 ± 2.5a	39.8 ± 7.6a	2.72 ± 0.23b	0.25 ± 0.03b	1.04 ± 0.14b	1.43 ± 0.08b	0.10 ± 0.01b
-N	34.0 ± 3.0d	42.5 ± 7.3a	2.12 ± 0.08d	0.26 ± 0.03b	0.80 ± 0.06d	1.06 ± 0.04d	0.14 ± 0.01a
-N+	44.7 ± 1.5c	41.0 ± 5.1a	2.41 ± 0.14c	0.25 ± 0.03b	0.97 ± 0.06bc	1.19 ± 0.06c	0.12 ± 0.01b

Note: The treatments were as follows: CK, complete supply of Mg and N; -Mg, Mg deficiency. -N, N deficiency; the cultivation duration was 14 days. -Mg+, Mg deprivation for 9 days followed by Mg resupply for 5 days; -N+, N deprivation for 9 days followed by N resupply for 5 days. The data are presented as the means and standard deviations from five replicates. Different lowercase letters indicate statistically significant differences ( $P < 0.05$ ) according to Duncan's multiple range test.

upper leaves but not in the middle leaves (Fig. 3). In contrast, under N starvation, N resupply not only significantly affected dry weight but also increased plant height (Table 1). Following N resupply, the chlorosis observed in the cucumber plants was markedly alleviated (Fig. 3).

Both Mg deficiency and N deficiency significantly decreased the photosynthetic rate in plants; however, these inhibitory effects were partially alleviated following the resupply of Mg or N. N resupply markedly increased key gas exchange parameters, including the net photosynthetic rate ( $P_n$ ), stomatal conductance ( $g_s$ ), and transpiration rate ( $T_r$ ), while it concomitantly reduced the intercellular  $CO_2$  concentration ( $C_i$ ) in cucumber (Fig. 4). Under N deficiency, all measured photosynthetic parameters were restored to control levels after N resupply. In contrast, although Mg resupply increased the photosynthetic rate, it remained below control levels. While Mg resupply had no significant effect on the maximum quantum efficiency of PSII ( $F_v/F_m$ ) in the middle leaves (Fig. 4(e)), it effectively mitigated leaf necrosis and supported the maintenance of normal growth in the upper leaves (Fig. 3).

### 3.3 Effects of Mg or N resupply on element concentrations and antioxidant enzyme activity in cucumber plants


Compared with the control (CK), Mg deficiency significantly

reduced the Mg concentration in plant leaves but markedly increased the concentrations of Mn and Mo by 73.3% and 72.8%, respectively (Fig. 5(a)). Additionally, the concentrations of Ca, Zn, and Cu increased by 53.2%, 46.9%, and 28.2%, respectively. N deficiency severely inhibited leaf growth, leading to a general decline in the concentrations of most measured elements, with the exception of P and Ca; among all the elements examined, the concentration of N decreased the most under N-deficient conditions (Fig. 5(b)). Following Mg resupply, the concentrations of most elements increased and remained at elevated levels, although the leaf Mg concentration remained substantially low. In contrast, N resupply increased the concentrations of the studied elements (excluding P and Ca), with the concentrations of all the elements eventually recovering to levels comparable to those of the control plants.

Compared with the control treatment, the Mg deficiency and N deficiency treatments significantly increased SOD activity by 185% and 91% in the lower leaves, 284% and 105% in the middle leaves, and 98% and 26% in the upper leaves, respectively (Fig. 6(a)). POD activity exhibited differential responses under the various deficiency conditions. In the Mg-deficient plants, POD activity in the lower and middle leaves was 7.47-fold and 36.85-fold greater, respectively, than that in the control plants, whereas no significant difference was detected between the N-deficient and control plants (Fig. 6(b)).



**Fig. 3** Cucumber leaves from different canopy positions under Mg or N deficiency and subsequent recovery conditions. Color images of representative leaves were captured after 5 days of nutrient resupply [(d9) + 5], with leaf samples arranged from lower to upper canopy positions from left to right. CK: plants supplied with sufficient amounts of Mg and N; -Mg: Mg deficiency treatment; -N: N deficiency treatment; the duration of cultivation was 14 days. -Mg+: Mg was withheld for 9 days, followed by 5 days of Mg resupply; -N+: N was withheld for 9 days, followed by 5 days of N resupply.




**Fig. 4** Effects of Mg or N deficiency and resupply on the photosynthetic performance and chlorophyll fluorescence characteristics of plants at 5 days after treatment initiation [(d9) + 5]. (a) Net photosynthetic rate ( $P_n$ ); (b) stomatal conductance ( $g_s$ ); (c) intercellular  $\text{CO}_2$  concentration ( $C_i$ ); (d) transpiration rate ( $T_r$ ); (e) maximum quantum efficiency of photosystem II ( $F_v/F_m$ ). The data are presented as the means and standard deviations of five biological replicates. Different letters indicate statistically significant differences ( $P < 0.05$ ) according to Duncan’s multiple range test.

The resupply of Mg had a minimal effect on enzyme activity, whereas N resupply decreased SOD activity in the lower leaves but increased POD activity in the middle leaves. Compared with the control treatment, Mg deficiency significantly increased the  $\text{H}_2\text{O}_2$  content by 130% in the middle leaves, and Mg resupply significantly reduced  $\text{H}_2\text{O}_2$  levels (Fig. 6(c)). In the N-deficient plants, the  $\text{H}_2\text{O}_2$  content in leaves at different positions was comparable to that in the control. Upon N resupply, the  $\text{H}_2\text{O}_2$  content in leaves at all positions decreased significantly in the N-deficient plants.

### 3.4 Effects of spraying DMTU on leaf $\text{H}_2\text{O}_2$ production and $F_v/F_m$

Following the application of the  $\text{H}_2\text{O}_2$  scavenger DMTU, the  $\text{H}_2\text{O}_2$  content the leaves at different positions of Mg-deficient plants was comparable to that in the control (CK) and N+DMTU-treated plants (Fig. 7(a)). In the N-deficient plants,  $\text{H}_2\text{O}_2$  levels in the middle leaves remained slightly elevated relative to those in the middle leaves of control plants after DMTU application. Although the  $F_v/F_m$  values in the Mg deficiency and Mg resupply groups were significantly lower



**Fig. 5** Effects of Mg (a) and N (b) deficiency and subsequent resupply on the concentrations of elements in leaves at 5 days after treatment initiation [(d9) + 5]. Changes in the elemental concentrations relative to those in the control group (CK), which received the standard nutrient solution, were calculated as follows: (elemental concentration in treatment – elemental concentration in CK)/(elemental concentration in CK). Asterisks denote statistically significant differences compared with the control group (\* $P < 0.05$ , \*\* $P \leq 0.01$ , \*\*\* $P \leq 0.001$ ), as determined by two-tailed Student's  $t$  test.

than those in the control group (Fig. 5), DMTU application substantially alleviated the extent of photoinhibition (Fig. 7(b)), which corresponded to reduced  $H_2O_2$  accumulation in leaves (Fig. 7(a)).


## 4 Discussion

### 4.1 Divergent stress effects: distinct physiological signatures of N deficiency versus Mg deficiency

Mg or N deficiency suppressed overall growth; however, different mechanisms were involved in their effects on biomass allocation. Deficiencies in Mg or N inhibited cucumber growth and biomass accumulation (Fig. 1). The observed symptoms were similar to those reported in previous studies<sup>[4,26]</sup>. The results revealed that the shoot/root ratios of the plants differed significantly among the –Mg, –N, and control treatment groups. Mg deficiency more strongly inhibited root growth than shoot growth; this increase in the shoot/root ratio under Mg deficiency has been previously reported in other plant species<sup>[27,28]</sup>. In contrast, N deficiency significantly reduced shoot growth, whereas root growth was less strongly affected, suggesting that the mechanisms through which Mg deficiency and N deficiency influence plant growth differ<sup>[29]</sup>. Previous

studies have suggested that Mg deficiency may impair phloem loading, thereby limiting the translocation of carbohydrates from source leaves to sink tissues, such as roots<sup>[30,31]</sup>. The reduced shoot/root ratio under N deficiency can be attributed to a biphasic response: suppression of the leaf elongation rate coupled with the maintenance or stimulation of root growth through the preferential allocation of assimilated carbon to the roots.

Compared with N deficiency, Mg deficiency resulted in more severe structural and functional impairments, as evidenced by leaf chlorosis and confirmed at the cellular level. More pronounced negative effects were observed in plants under Mg deficiency, particularly in the middle and lower leaves (Fig. 1, Fig. 2(a)). Mg deficiency resulted in the development of typical interveinal chlorosis, a phenomenon consistent with the findings of Cai et al<sup>[2]</sup>. Leaves under N deficiency exhibited a pale green appearance, with no necrotic lesions observed. In contrast, light microscopic examination confirmed structural damage in leaves under Mg deficiency, whereas leaves under N deficiency maintained an intact cellular architecture. Further evidence indicating greater cellular damage in leaves under Mg deficiency was provided by thermal imaging, which, through digital infrared thermography, clearly displayed the leaf temperature and water status on the monitor, offering direct visualization of physiological stress<sup>[32,33]</sup>. Both Mg deficiency



**Fig. 6** Effects of Mg or N deficiency and resupply on antioxidant enzyme activity and H<sub>2</sub>O<sub>2</sub> content in leaves at 5 days after treatment initiation [(d9) + 5]. (a) SOD enzyme activity; (b) POD enzyme activity; (c) H<sub>2</sub>O<sub>2</sub> content. The data represent the means ± standard deviations of five biological replicates. Different lowercase letters indicate statistically significant differences ( $P < 0.05$ ) determined by Duncan’s multiple range test among treatments within the same leaf position. U·g<sup>-1</sup> FW, units per gram fresh weight; mmol·(g prot)<sup>-1</sup>, millimoles per gram of protein.


and N deficiency negatively affected the  $T_r$ , whereas the leaf temperature increased exclusively under N deficiency (Fig. 2). Leaf temperature consistently increases because of the inhibition of  $T_r$  caused by water loss from cellular damage resulting from Mg deficiency. Cakmak and Kirkby proposed that Mg deficiency-induced leaf necrosis may be attributed to impaired photosynthetic CO<sub>2</sub> fixation and the subsequent increase in ROS production<sup>[8]</sup>. Unlike Mg deficiency, N deficiency reduces photosynthetic activity, thereby leading to excess electron flow and promoting the overproduction of ROS<sup>[34]</sup>.

N and Mg are essential elements for plant photosynthesis. Both Mg deficiency and N deficiency significantly reduced gas exchange parameters, including  $P_n$ ,  $g_s$ , and  $T_r$ , and increased  $C_i$  (Fig. 4); these observations are consistent with findings

reported in previous studies<sup>[4,35,36]</sup>. Rubisco is the key rate-limiting enzyme in CO<sub>2</sub> assimilation, and its activity has been demonstrated to be positively correlated with Mg and N concentrations, as Mg and N serve as activators or enzymatic activators or structural components<sup>[13,37]</sup>. Moreover, both Mg and N are critical components of chlorophyll; therefore, the decline in photosynthetic performance under deficient conditions can be attributed not only to reduced carbon fixation but also to impaired light-dependent reactions.

#### 4.2 Differential recovery efficacy: the effects of N resupply versus Mg resupply in on alleviating deficiency symptoms

Nine days after the initiation of treatment, the plants were



**Fig. 7** Effects of the foliar application of DMTU on the hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) content and maximum quantum efficiency of photosystem II ( $F_v/F_m$ ) in leaves. (a) H<sub>2</sub>O<sub>2</sub> content across leaves in different positions; statistical analysis was conducted to compare the H<sub>2</sub>O<sub>2</sub> concentrations at the same leaf positions under different treatments; (b)  $F_v/F_m$  values in the middle leaves. The data are presented as the means  $\pm$  standard deviations of five biological replicates. Different lowercase letters indicate statistically significant differences ( $P < 0.05$ ) determined by Duncan's multiple range test among treatments within the same leaf position. mmol·(g prot)<sup>-1</sup>, millimoles per gram of protein.

resupplied with Mg or N to further investigate the distinct physiological responses under Mg deficiency and N deficiency. Five days later, leaves subjected to N deficiency exhibited signs of recovery, whereas necrotic symptoms persisted in leaves subjected to Mg deficiency (Fig. 3). These observations can be attributed to the more severe and detrimental effects of Mg deficiency on cucumber plants. Upon N resupply, the photosynthetic rate was restored to normal levels; however, although the improvement in photosynthetic capacity by Mg resupply remained significantly lower than that in the control treatment. This discrepancy may be explained by the disruption of the photosynthetic electron transport chain under Mg deficiency. Consequently, Mg deficiency likely induced irreversible damage to the leaf tissues.

Photosystem II (PSII) activity plays a crucial role in mediating the photosynthetic response to environmental stress; previous studies have indicated that the  $F_v/F_m$  ratio is a key indicator of photoinhibition in PSII<sup>[36]</sup>. Therefore, we assessed the  $F_v/F_m$  ratios under various treatment conditions to evaluate and compare the extent of leaf damage associated with Mg deficiency and N deficiency. Similarly, Mg deficiency significantly reduced the  $F_v/F_m$  ratio, whereas N deficiency did not significantly affect this ratio (Fig. 4). Previous studies have demonstrated that Mg deficiency reduces the  $F_m$  and the  $F_v/F_m$  while increasing the minimum fluorescence ( $F_o$ ) in sugar beet<sup>[38]</sup>. Furthermore, additional studies have demonstrated that Mg deficiency reduces photosynthetic efficiency,  $F_v/F_m$ , and the photochemical quenching coefficient (qP) but increases nonphotochemical quenching (NPQ), with no

significant effect on  $F_o$ <sup>[39]</sup>. Photoinhibition under Mg deficiency may be attributed to the absorption of photon energy by light-harvesting pigments exceeding the capacity of the photosynthetic electron transport chain to utilize it; the resulting excess excitation energy may lead to the formation of ROS<sup>[31,40]</sup>. A significant accumulation of ROS occurs in leaf cells, particularly within chloroplasts, which may lead to chlorosis and cellular death<sup>[41]</sup>. This explains the limited recovery of photosynthetic capacity following Mg resupply.

Nutrient deficiency can trigger the generation of ROS, resulting in elevated levels of H<sub>2</sub>O<sub>2</sub> in plant cells<sup>[42]</sup>. A previous study demonstrated that elevated accumulation of H<sub>2</sub>O<sub>2</sub> can inhibit photosynthetic efficiency. The removal of H<sub>2</sub>O<sub>2</sub> may therefore be a critical factor in alleviating photoinhibition<sup>[43]</sup>. Therefore, the differences in damage caused by N deficiency and Mg deficiency may be attributed to variations in H<sub>2</sub>O<sub>2</sub> accumulation levels. Furthermore, the differential recovery of photosynthetic capacity in cucumber leaves following N or Mg supplementation is closely associated with the respective efficiencies in scavenging H<sub>2</sub>O<sub>2</sub>.

### 4.3 The recovery of N-deficient and Mg-deficient plants is dependent on the H<sub>2</sub>O<sub>2</sub> concentration

H<sub>2</sub>O<sub>2</sub> is a component of oxidative metabolism and plays important roles in biochemical and physiological processes<sup>[44,45]</sup>. Therefore, we further investigated the H<sub>2</sub>O<sub>2</sub> content in leaves under various treatment conditions. The -Mg

treatment resulted in the most severe deficiency symptoms, with  $H_2O_2$  levels reaching twice those of the control (Fig. 6(c)).  $H_2O_2$  is an inevitable byproduct of the photosynthetic electron transport chain, and its excessive accumulation can lead to damage to photosystems<sup>[46]</sup>. ROS are continuously generated at a basal level, and a balance is maintained between their generation and elimination. They play crucial roles in plants. Nevertheless, once the endogenous ROS balance is disrupted by biotic and abiotic environmental stresses, they can also lead to extensive cellular damage. In the event of Mg deficiency, the export of sucrose from the phloem is severely impeded, which subsequently inhibits  $CO_2$  fixation. Excessive absorbed light energy is subsequently transferred to molecular oxygen, resulting in the generation of ROS<sup>[8,15]</sup>. In the case of N deficiency, the transfer of nitrogen nutrients to the cell wall to resist stress-induced damage weakens nitrogen allocation from the cytoplasm to photosynthetic components. The activity of photosynthesis-related enzymes in chloroplasts decreases notably, leading to the accumulation of  $CO_2$  and a substantial increase in the concentration of ROS<sup>[17]</sup>. As previously noted, nutrient deficiency may contribute to increased concentrations of ROS, such as  $H_2O_2$ . The main site where ROS are produced is the chloroplasts. Therefore, we believe that the primary factor underlying the differential effects of N deficiency and Mg deficiency on photosynthetic damage is  $H_2O_2$ .

Plants possess diverse antioxidant enzymes, such as SOD, CAT, POD, and APX, to eliminate ROS and protect cells against oxidative stress<sup>[2]</sup>. The Mg deficiency-induced increase in antioxidant enzyme activity is a common phenomenon in the leaves of various plant species, including beans, maize, rice, and citrus<sup>[2,47,48]</sup>. In addition to Mg deficiency, N deficiency also increases ROS production, thereby inducing DNA strand breaks, lipid peroxidation, and ultimately cell death<sup>[49]</sup>. Previous studies have demonstrated that the application of ammonium nitrogen enhances antioxidant enzyme activity in blackberries<sup>[50]</sup>, and under heat stress, the application of Mg significantly increased the activities of SOD, POD, and CAT in broad beans while reducing the accumulation of malondialdehyde (MDA) and  $H_2O_2$ <sup>[51]</sup>. An increase in the activities of SOD and POD, which can remove free radicals and maintain the steady-state cellular level of ROS, is likely associated with oxidative stress in plants<sup>[52]</sup>.

The activities of SOD and POD exhibited distinct patterns under N or Mg deficiency and during the subsequent resupply. The variation in enzyme activities among the different

treatments might be associated with the differences in leaf ion concentrations. Mg deficiency significantly altered the levels of Mn and Zn in cucumber leaves (Fig. 5). Ye et al.<sup>[36]</sup> reported antagonistic interactions between Mg and Mn, as well as Zn. These findings suggest that Mg deficiency may increase the uptake of Mn and Mo. Similar results have been reported for potato, wheat, maize, and oil rape<sup>[53–56]</sup>. Mn has been reported to act as an activator of numerous enzymes, particularly those involved in the antioxidant system. In plants, the oxygen-evolving complex (OEC) in PSII, manganese superoxide dismutase (MnSOD), and oxalate oxidase have been demonstrated to exclusively require Mn<sup>[57]</sup>. Moreover, the activities of antioxidant system enzymes, including POD, SOD, and APX, increase with increasing manganese concentration<sup>[58]</sup>. In the present study, Mg deficiency led to a substantial increase in the Mn absorption rate, exceeding 50%, which remained elevated even after Mg resupply. In contrast, compared with the control treatment, N deficiency did not significantly affect the Mn concentration. Thus, alterations in ion uptake, specifically that of Mn, may constitute a component of the adaptive response of plants to alleviate the accumulation of ROS under nutrient stress<sup>[59]</sup>.

In addition to Mn, elevated Cu concentrations are also observed under Mg deficiency and during Mg resupply. Excessive Cu is toxic to plant cells and can adversely affect various biochemical and physiological processes, particularly by disrupting photosynthesis and N metabolism<sup>[60]</sup>. Compared with N deficiency and resupply treatments, minimal differences in ion concentrations were observed following Mg resupply, suggesting that the irreversible effects of Mg deficiency may be attributed to cellular damage caused by  $H_2O_2$  rather than ion imbalance. Furthermore, ion concentrations were restored to normal levels upon N resupply. Given the low levels of  $H_2O_2$  in leaves under N deficiency or N resupply conditions, it is reasonable to conclude that N supplementation can gradually mitigate the effects of reactive oxygen species (ROS), enabling leaves to recover to a normal physiological state.

To further validate our hypothesis, we attempted to protect the leaves by scavenging  $H_2O_2$ . Dimethylthiourea (DMTU), a widely used scavenger of  $H_2O_2$ , has been applied exogenously to leaves in previous studies to eliminate hydrogen peroxide and alleviate photooxidative damage<sup>[61,62]</sup>. Our studies demonstrated that the application of DMTU effectively reduced  $H_2O_2$  accumulation in nutrient-deficient plants,

restoring it to the same level as that of the control group, particularly in Mg-deficient plants (Fig. 7(a)). Crucially, this reduction in oxidative load was correlated with a significant alleviation of photoinhibition, as evidenced by the partial recovery of  $F_v/F_m$  in Mg-deficient leaves (Fig. 7(b)). The direct correlation between  $H_2O_2$  scavenging and the increase in a critical photosynthetic efficiency parameter strongly supports our hypothesis that the accumulation of  $H_2O_2$  is an important limiting factor for recovery from nutrient deficiency. The finding that DMTU was unable to fully restore the maximum quantum efficiency of  $F_v/F_m$  to the control levels in Mg-deficient plants is consistent with the observed partial recovery of plant growth only after Mg resupply. The potential reason is that under Mg deficiency conditions, an excessive quantity of  $H_2O_2$  causes irreversible damage to the leaf cells of plants. In contrast, the lower  $H_2O_2$  burden under N deficiency conditions enables the recovery of photosynthesis and initiation of growth after N supplementation or DMTU treatment. These results further emphasize the crucial role that the  $H_2O_2$  concentration plays in determining the degree of recovery from nutritional stress.

## 5 Conclusions

The results of the present study demonstrated that compared with N deficiency, Mg deficiency caused more severe and irreversible damage to leaf structure and photosynthetic function. However, N supplementation effectively alleviated symptoms of N deficiency, and Mg supplementation resulted in a weaker ability to alleviate symptoms of Mg deficiency. This difference can be attributed to the varying accumulation levels of  $H_2O_2$  under nutrient deficiency stress. Recovery from N or Mg deficiency is dependent on the  $H_2O_2$  concentration; excessive accumulation of  $H_2O_2$  can disrupt cellular structure and ultimately lead to irreversible leaf damage. In agricultural production, N deficiency can be mitigated through timely N application, whereas Mg deficiency causes irreversible damage. Collectively, these findings enhance our understanding of the adverse effects of Mg deficiency and provide insights into mitigating the damage caused by Mg deficiency stress. Since plants become considerably more challenging to recover once visible symptoms of Mg deficiency manifest, early diagnosis and supplementation with magnesium fertilizer are of particular significance in modern agriculture.

### Acknowledgements

This paper was funded by the National Key Research and Development Program of China (2023YFD1901101) and the National Natural Science Foundation of China (42577134).

### Compliance with ethics guidelines

Qi Zhou, Kehao Chen, Ying Wang, Xusheng Meng, Shiyu Wang, Min Wang, and Shiwei Guo declare that they have no conflicts of interest or financial conflicts to disclose. This article does not contain any studies with human or animal subjects performed by any of the authors.

## REFERENCES

1. Batool S, Hussain Z, Bano A, Batool A, Iqbal P. Determination of macronutrients concentration in dry foliage of *sophora mollis* to use green lumen as organic fertilizer. *Insights-Journal of Health and Rehabilitation*, 2024, 2(2): 513–521
2. Cai Y T, Zhang H, Qi Y P, Ye X, Huang Z R, Guo J X, Chen L S, Yang L T. Responses of reactive oxygen species and methylglyoxal metabolisms to magnesium-deficiency differ greatly among the roots, upper and lower leaves of *Citrus sinensis*. *BMC Plant Biology*, 2019, 19(1): 76
3. De Bang T C, Husted S, Laursen K H, Persson D P, Schjoerring J K. The molecular-physiological functions of mineral macronutrients and their consequences for deficiency symptoms in plants. *New Phytologist*, 2021, 229(5): 2446–2469
4. Ye B Z, Shi S L, Zhang R Z, Nie P Q, Tang X Q, Wang K C. Effects of nitrogen deficiency and nitrogen recovery treatments on growth and some physiological and biochemical indexes of *Isatis indigotica* seedlings. *Journal of Plant Resources and Environment*, 2015, 24(4): 83–88 (in Chinese)
5. Hawkesford M, Horst W, Kichey T, Lambers H, Schjoerring J, Møller I S, White P. Functions of macronutrients. In: Marschner P, ed. *Marschner's Mineral Nutrition of Higher Plants*. 3rd ed. Waltham: Academic Press, 2012, 135–189

6. Farhat N, Elkhouni A, Zorrig W, Smaoui A, Abdely C, Rabhi M. Effects of magnesium deficiency on photosynthesis and carbohydrate partitioning. *Acta Physiologiae Plantarum*, 2016, **38**(6): 145
7. Zhao R, Dielen V, Kinet J M, Boutry M. Cosuppression of a plasma membrane H<sup>+</sup>-ATPase isoform impairs sucrose translocation, stomatal opening, plant growth, and male fertility. *Plant Cell*, 2000, **12**(4): 535–546
8. Cakmak I, Kirkby E A. Role of magnesium in carbon partitioning and alleviating photooxidative damage. *Physiologia Plantarum*, 2008, **133**(4): 692–704
9. Kiba T, Krapp A. Plant nitrogen acquisition under low availability: regulation of uptake and root architecture. *Plant and Cell Physiology*, 2016, **57**(4): 707–714
10. Chen L R, Liu M Y, Cai Y J, Wu L Q, Zhang Q F. Magnesium deficiency differentiated effects on the roots growth and shoot metabolism by regulating the distribution of photosynthetic products in tea plants. *Scientia Horticulturae*, 2024, **338**: 113748
11. Tewari R K, Kumar P, Sharma P N. Oxidative stress and antioxidant responses in young leaves of mulberry plants grown under nitrogen, phosphorus or potassium deficiency. *Journal of Integrative Plant Biology*, 2007, **49**(3): 313–322
12. Peng Y Y, Liao L L, Liu S, Nie M M, Li J, Zhang L D, Ma J F, Chen Z C. Magnesium deficiency triggers SGR-mediated chlorophyll degradation for magnesium remobilization. *Plant Physiology*, 2019, **181**(1): 262–275
13. Warren C R. The photosynthetic limitation posed by internal conductance to CO<sub>2</sub> movement is increased by nutrient supply. *Journal of Experimental Botany*, 2004, **55**(406): 2313–2321
14. Stec B. Structural mechanism of RuBisCo activation by carbamylation of the active site lysine. *Proceedings of the National Academy of Sciences of the United States of America*, 2012, **109**(46): 18785–18790
15. Foyer C H, Hanke G. ROS production and signalling in chloroplasts: cornerstones and evolving concepts. *The Plant Journal*, 2022, **111**(3): 642–661
16. Ruan L, Wei K, Li J W, He M D, Wu L Y, Aktar S, Wang L Y, Cheng H. Responses of tea plants (*Camellia sinensis*) with different low-nitrogen tolerances during recovery from nitrogen deficiency. *Journal of the Science of Food and Agriculture*, 2021, **102**(4): 1405–1414
17. Kong L G, Wang F H, Si J S, Feng B, Zhang B, Li S D, Wang Z. Increasing in ROS levels and callose deposition in peduncle vascular bundles of wheat (*Triticum aestivum* L.) grown under nitrogen deficiency. *Journal of Plant Interactions*, 2013, **8**(2): 109–116
18. Marschner H, Cakmak I. High light intensity enhances chlorosis and necrosis in leaves of zinc, potassium, and magnesium deficient bean (*Phaseolus vulgaris*) plants. *Journal of Plant Physiology*, 1989, **134**(3): 308–315
19. Da Silva D M, De Souza K R D, Vilas Boas L V, Santos Alves Y, Alves J D. The effect of magnesium nutrition on the antioxidant response of coffee seedlings under heat stress. *Scientia Horticulturae*, 2017, **224**: 115–125
20. Chen H F, Zhang Q, Cai H M, Zhou W, Xu F S. H<sub>2</sub>O<sub>2</sub> mediates nitrate-induced iron chlorosis by regulating iron homeostasis in rice. *Plant, Cell & Environment*, 2018, **41**(4): 767–781
21. Saxena I, Srikanth S, Chen Z. Cross talk between H<sub>2</sub>O<sub>2</sub> and interacting signal molecules under plant stress response. *Frontiers in Plant Science*, 2016, **7**: 570
22. Balazadeh S, Schildhauer J, Araújo W L, Munné-Bosch S, Fernie A R, Proost S, Humbeck K, Mueller-Roeber B. Reversal of senescence by N resupply to N-starved *Arabidopsis thaliana*: transcriptomic and metabolomic consequences. *Journal of Experimental Botany*, 2014, **65**(14): 3975–3992
23. Fataftah N, Mohr C, Hajirezaei M R, Von Wirén N, Humbeck K. Changes in nitrogen availability lead to a reprogramming of pyruvate metabolism. *BMC Plant Biology*, 2018, **18**(1): 77
24. Stewart R R C, Bewley J D. Lipid peroxidation associated with accelerated aging of soybean axes. *Plant Physiology*, 1980, **65**(2): 245–248
25. Wang R R, Huang J, Liang A C, Wang Y, Mur L A J, Wang M, Guo S W. Zinc and copper enhance cucumber tolerance to fusaric acid by mediating its distribution and toxicity and modifying the antioxidant system. *International Journal of Molecular Sciences*, 2020, **21**(9): 3370
26. Campos C N S, Teixeira G C M, De Mello Prado R, Caione G, da Silva G B Jr, De David C H O, Sales A C, Roque C G, Teodoro P E. Macronutrient deficiency in cucumber plants: impacts in nutrition, growth and symptoms. *Journal of Plant Nutrition*, 2021, **44**(17): 2609–2626
27. Cakmak I, Hengeler C, Marschner H. Partitioning of shoot and root dry matter and carbohydrates in bean plants suffering from phosphorus, potassium and magnesium deficiency. *Journal of Experimental Botany*, 1994, **45**(278): 1245–1250
28. Mengutay M, Ceylan Y, Kutman U B, Cakmak I. Adequate magnesium nutrition mitigates adverse effects of heat stress on maize and wheat. *Plant and Soil*, 2013, **368**(1–2): 57–72
29. Richard-Molard C, Krapp A, Brun F, Ney B, Daniel-Vedele F, Chaillou S. Plant response to nitrate starvation is determined by N storage capacity matched by nitrate uptake capacity in two *Arabidopsis* genotypes. *Journal of Experimental Botany*, 2008, **59**(4): 779–791
30. Cakmak I, Hengeler C, Marschner H. Changes in phloem export of sucrose in leaves in response to phosphorus, potassium and magnesium deficiency in bean plants. *Journal of Experimental Botany*, 1994, **45**(278): 1251–1257
31. Hermans C, Verbruggen N. Physiological characterization of Mg deficiency in *Arabidopsis thaliana*. *Journal of Experimental Botany*, 2005, **56**(418): 2153–2161
32. Wang M, Ling N, Dong X, Zhu Y Y, Shen Q R, Guo S W.

- Thermographic visualization of leaf response in cucumber plants infected with the soil-borne pathogen *Fusarium oxysporum* f. sp. *cucumerinum*. *Plant Physiology and Biochemistry*, 2012, **61**: 153–161
33. Sun Y M, Wang M, Li Y R, Gu Z C, Ling N, Shen Q R, Guo S W. Wilted cucumber plants infected by *Fusarium oxysporum* f. sp. *cucumerinum* do not suffer from water shortage. *Annals of Botany*, 2017, **120**(3): 427–436
  34. Xiang F, Zhou L Y, Liu H Y, Li W. Improving tea quality by balancing ROS and antioxidant system through appropriate ammonium nitrogen application. *Sustainability*, 2022, **14**(15): 9354
  35. Yang G H, Yang L T, Jiang H X, Li Y, Wang P, Chen L S. Physiological impacts of magnesium-deficiency in *Citrus* seedlings: photosynthesis, antioxidant system and carbohydrates. *Trees*, 2012, **26**(4): 1237–1250
  36. Ye X, Chen X F, Deng C L, Yang L T, Lai N W, Guo J X, Chen L S. Magnesium-deficiency effects on pigments, photosynthesis and photosynthetic electron transport of leaves, and nutrients of leaf blades and veins in *Citrus sinensis* seedlings. *Plants*, 2019, **8**(10): 389
  37. Li J, Yokosho K, Liu S, Cao H R, Yamaji N, Zhu X G, Liao H, Ma J F, Chen Z C. Diel magnesium fluctuations in chloroplasts contribute to photosynthesis in rice. *Nature Plants*, 2020, **6**(7): 848–859
  38. Hermans C, Johnson G N, Strasser R J, Verbruggen N. Physiological characterisation of magnesium deficiency in sugar beet: acclimation to low magnesium differentially affects photosystems I and II. *Planta*, 2004, **220**(2): 344–355
  39. Laing W, Greer D, Sun O, Beets P, Lowe A, Payn T. Physiological impacts of Mg deficiency in *Pinus radiata*: growth and photosynthesis. *New Phytologist*, 2000, **146**(1): 47–57
  40. Chen L S, Cheng L L. Both xanthophyll cycle-dependent thermal dissipation and the antioxidant system are up-regulated in grape (*Vitis labrusca* L. cv. Concord) leaves in response to N limitation. *Journal of Experimental Botany*, 2003, **54**(390): 2165–2175
  41. Choudhury F K, Rivero R M, Blumwald E, Mittler R. Reactive oxygen species, abiotic stress and stress combination. *The Plant Journal*, 2017, **90**(5): 856–867
  42. Nieves-Cordones M, López-Delacalle M, Ródenas R, Martínez V, Rubio F, Rivero R M. Critical responses to nutrient deprivation: a comprehensive review on the role of ROS and RNS. *Environmental and Experimental Botany*, 2019, **161**: 74–85
  43. Kumagai E, Araki T, Ueno O. Effect of nitrogen-deficiency on midday photoinhibition in flag leaves of different rice (*Oryza sativa* L.) cultivars. *Photosynthetica*, 2009, **47**(2): 241–246
  44. Yao X H, Mu Y J, Zhang L Y, Chen L, Zou S S, Chen X C, Lu K, Dong H S. AtPIP1;4 and AtPIP2;4 cooperatively mediate H<sub>2</sub>O<sub>2</sub> transport to regulate plant growth and disease resistance. *Plants*, 2024, **13**(7): 1018
  45. Zhu Z L, Dai D D, Zheng M Z, Shi Y L, Siddique S, Wang F F, Zhang S R, Xie C S, Bo D X, Hu B Y, Chen Y Y, Peng D H, Sun M, Zheng J S. Root-knot nematodes exploit the catalase-like effector to manipulate plant reactive oxygen species levels by directly degrading H<sub>2</sub>O<sub>2</sub>. *Molecular Plant Pathology*, 2024, **25**(9): e70000
  46. Zhang H H, Li X, Xu Z S, Wang Y, Teng Z Y, An M J, Zhang Y H, Zhu W X, Xu N, Sun G Y. Toxic effects of heavy metals Pb and Cd on mulberry (*Morus alba* L.) seedling leaves: photosynthetic function and reactive oxygen species (ROS) metabolism responses. *Ecotoxicology and Environmental Safety*, 2020, **195**: 110469
  47. He Z Q, Shang X, Jin X Z, Wang X K, Xing Y Y. Calcium and magnesium regulation of kernel sugar content in maize: role of endogenous hormones and antioxidant enzymes. *International Journal of Molecular Sciences*, 2025, **26**(1): 200
  48. Zhou H M, Peng J, Zhao W L, Zeng Y J, Xie K L, Huang G J. Leaf diffusional capacity largely contributes to the reduced photosynthesis in rice plants under magnesium deficiency. *Plant Physiology and Biochemistry*, 2024, **209**: 108565
  49. Zhou Y Z, He Y P, Sun J Y, Zhang J W, Zhan Y G. Effects of nitrogen deficiency on physiology and growth of *Fraxinus mandshurica*. *Pakistan Journal of Botany*, 2018, **50**(1): 179–187
  50. Duan Y K, Yang H Y, Yang H, Wei Z W, Che J L, Wu W L, Lyu L F, Li W L. Physiological and morphological responses of blackberry seedlings to different nitrogen forms. *Plants*, 2023, **12**(7): 1480
  51. Siddiqui M H, Alamri S A, Al-Khaishany M Y Y, Al-Qutami M A, Ali H M, Al-Wahaibi M H, Al-Wahaibi M S, Alharby H F. Mitigation of adverse effects of heat stress on *Vicia faba* by exogenous application of magnesium. *Saudi Journal of Biological Sciences*, 2018, **25**(7): 1393–1401
  52. Han M M, Wang B Y, Song G L, Shi S Q. Comparative study of alleviation effects of DMTU and PCIB on root growth inhibition in two tall fescue varieties under cadmium stress. *Ecotoxicology and Environmental Safety*, 2020, **196**: 110528
  53. Cao W X, Tibbitts T W. Growth, carbon dioxide exchange and mineral accumulation in potatoes grown at different magnesium concentrations. *Journal of Plant Nutrition*, 1992, **15**(9): 1359–1371
  54. Chatterjee C, Nautiyal N, Agarwala S C. Influence of changes in manganese and magnesium supply on some aspects of wheat physiology. *Soil Science and Plant Nutrition*, 1994, **40**(2): 191–197
  55. Jezek M, Geilfus C M, Bayer A, Mühling K H. Photosynthetic capacity, nutrient status, and growth of maize (*Zea mays* L.) upon MgSO<sub>4</sub> leaf-application. *Frontiers in Plant Science*, 2015, **5**: 781
  56. Billard V, Maillard A, Coquet L, Jouenne T, Cruz F, Garcia-

- Mina J M, Yvin J C, Ourry A, Etienne P. Mg deficiency affects leaf Mg remobilization and the proteome in *Brassica napus*. *Plant Physiology and Biochemistry*, 2016, **107**: 337–343
57. Alejandro S, Höller S, Meier B, Peiter E. Manganese in plants: from acquisition to subcellular allocation. *Frontiers in Plant Science*, 2020, **11**: 300
58. Pan Y H, Shi J N, Li J Y, Zhang R, Xue Y B, Liu Y. Regulatory mechanism through which old soybean leaves respond to Mn toxicity stress. *International Journal of Molecular Sciences*, 2024, **25**(10): 5341
59. Zhang Y, Liang Y, Zhao X, Jin X, Hou L P, Shi Y, Ahammed G J. Silicon compensates phosphorus deficit-induced growth inhibition by improving photosynthetic capacity, antioxidant potential, and nutrient homeostasis in tomato. *Agronomy*, 2019, **9**(11): 733
60. Xing M L, Li W M, Yu H, Wang Y Z, Wu F, Wu M, Xu J. Physiological and transcriptome analyses reveal copper toxicity responses in tomato plants. *Environmental and Experimental Botany*, 2024, **224**: 105819
61. Wu C J, Chen J H, Zhang J Z, Gao R, Li S L, Li Y M, Li Y, Cui D D, Wu G X. Exogenous H<sub>2</sub>O<sub>2</sub> improves the cold tolerance of tomato grafted seedlings by alleviating oxidative damage. *Chinese Journal of Biochemistry and Molecular Biology*, 2023, **39**(8): 1122–1132 (in Chinese)
62. Li W Q, Li J Y, Zhang Y F, Luo W Q, Dou Y, Yu S. Effect of reactive oxygen scavenger N,N'-dimethylthiourea (DMTU) on seed germination and radicle elongation of maize. *International Journal of Molecular Sciences*, 2023, **24**(21): 15557