

The uncertainty analysis of the MODIS GPP product in global maize croplands

Xiaojuan HUANG^{1,2}, Mingguo MA (✉)^{1,2}, Xufeng WANG³, Xuguang TANG^{1,2}, Hong YANG⁴

1 Chongqing Engineering Research Center for Remote Sensing Big Data Application, Southwest University, Chongqing 400715, China
2 Research Base of Karst Eco-environments at Nanchuan in Chongqing, Ministry of Natural Resources, School of Geographical Sciences, Southwest University, Chongqing 400715, China
3 Northwest Institute of Eco-environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China
4 Department of Geography and Environmental Science, University of Reading, Reading RG6 6AB, UK

© Higher Education Press and Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract Gross primary productivity (GPP) is very important in the global carbon cycle. Currently, the newly released estimates of 8-day GPP at 500 m spatial resolution (Collection 6) are provided by the Moderate Resolution Imaging Spectroradiometer (MODIS) Land Science Team for the global land surface via the improved light use efficiency (LUE) model. However, few studies have evaluated its performance. In this study, the MODIS GPP products (GPP_{MOD}) were compared with the observed GPP (GPP_{EC}) values from site-level eddy covariance measurements over seven maize flux sites in different areas around the world. The results indicate that the annual GPP_{MOD} was underestimated by 6%–58% across sites. Nevertheless, after incorporating the parameters of the calibrated LUE, the measurements of meteorological variables and the reconstructed Fractional Photosynthetic Active Radiation (FPAR) into the GPP_{MOD} algorithm in steps, the accuracies of GPP_{MOD} estimates were improved greatly, albeit to varying degrees. The differences between the GPP_{MOD} and the GPP_{EC} were primarily due to the magnitude of LUE and FPAR. The underestimate of maize cropland LUE was a widespread problem which exerted the largest impact on the GPP_{MOD} algorithm. In American and European sites, the performance of the FPAR exhibited distinct differences in capturing vegetation GPP during the growing season due to the canopy heterogeneity. In addition, at the DE-Kli site, the GPP_{MOD} abruptly produced extreme low values during the growing season because of the contaminated FPAR from a continuous rainy season. After correcting the noise of the FPAR, the accuracy of the GPP_{MOD} was improved by approximately 14%. Therefore, it is crucial to further

improve the accuracy of global GPP_{MOD} , especially for the maize crop ecosystem, to maintain food security and better understand global carbon cycle.

Keywords MODIS GPP, eddy covariance, maize cropland, validation, improvement

1 Introduction

Gross primary productivity (GPP) is important for the global carbon cycle between the biosphere and other systems (Lai et al., 2016). It is still a big challenge to accurately quantify the global terrestrial GPP at high spatial and temporal resolutions (Zhang et al., 2015). At the ecosystem level, the eddy covariance technique has been widely used to measure the exchanges of CO_2 , water, and energy between the atmosphere and the land surface (Jung et al., 2015). The light use efficiency (LUE), first proposed by Monteith (1972), is one of approaches to estimate GPP. With the development of geographic information system (GIS) technology, it is increasingly easy to use remote sensing (RS) technology to observe the earth and provide continuous data for the LUE model.

Since 2000, the Moderate Resolution Imaging Spectroradiometer has provided a new way to monitor GPP regularly from space with a spatial resolution of 500 m and the temporal resolution of 8 days (Running et al., 2004), and provided datasets from Collection 4 (C4) to Collection 6 (C6). Compared with C6, there are two main problems with the C4 MOD17A2H dataset. Firstly, in some regions with higher frequencies of cloud cover, the 8-day Maximum Value Composite (MVC) is still contaminated by clouds, yielding incorrect 8-day GPP values. Secondly, the C4 MOD17A2H dataset fails to account in the algorithm for the mismatched spatial resolution between

a 1-km Moderate Resolution Imaging Spectroradiometer (MODIS) pixel and the corresponding $1^{\circ} \times 1.25^{\circ}$ meteorological data from the Data Assimilation Office (DAO). The C4 MOD17A2H data were then improved for Collection 5 (C5) based on the development of the MODIS fraction absorbed photosynthetically active radiation (FPAR) and plant maintenance respiration by National Aeronautics and Space Administration (NASA) in 2007. At present, the C6 of MOD17A2 GPP products have higher quality than the previous collections with a spatial resolution from 500 m to 1 km and improvement in the maximum LUE of the crop and other parameters in the algorithm.

It is necessary to validate the C6 GPP_{MOD} products with local observation (Fu et al., 2012). It is a challenging task to analyze the uncertainty of GPP_{MOD} due to the difficulty of direct measurement of GPP (Chen et al., 2014). Eddy covariance (EC) flux towers have been increasingly used to measure GPP indirectly by partitioning the net ecosystem exchange (NEE) into the ecosystem respiration (ER) during daylight periods. A growing number of flux sites have been used to validate the MODIS product (Turner et al., 2003; Fu et al., 2012; Wang et al., 2013; Tang et al., 2015). Additionally, the eddy covariance technique has made the calibration process of the LUE more feasible than ever before.

GPP_{MOD} has been validated in forest (Gebremichael and Barros, 2006), grassland, and cropland (Zhang et al., 2008; Fu et al., 2012; Wang et al., 2013) with different biomes across different climate zones, indicating that GPP_{MOD} products were overestimated at low productivity sites because of the overestimate of FPAR and underestimated at higher productivity sites due to the underestimate of LUE (Fu et al., 2012). In other words, there is a large underestimation of the GPP_{MOD} due to the uncertainty of the maximum LUE and FPAR in some areas. In the GPP_{MOD} algorithm, the parameter of LUE, which depends on the simple look-up TABLE approach, is the most uncertain component (Running et al., 1999; Wang et al., 2013). In addition, the MODIS FPAR is subject to uncertainty because of atmospheric conditions during the satellite overpass, view angle geometry, and canopy heterogeneity (Cohen et al., 2003; Fensholt et al., 2004).

Crops account for approximately 24% of the earth's land

surface (Peng and Gitelson, 2011). Maize is one of the primary foods for human consumption and an essential fodder for animals. Therefore, it is very important to accurately estimate the maize GPP (Gitelson et al., 2008). However, few works have validated the maize GPP. Wang et al. (2013) validated the GPP_{MOD} product at 4 maize sites in northern China and found that the LUE was the primary reason for the underestimate of the GPP.

This study focuses on the validation of the GPP_{MOD} performance on global maize cropland and aim to make marked improvement of the accuracy. To acquire the estimate of GPP_{MOD} , this study relied on the GPP_{MOD} algorithm driven by the local meteorological data and the LUE calibrated by the seven eddy covariance flux towers and reconstructed FPAR. The objectives are (i) to assess the performance of the MODIS GPP products in maize crops at seven sites around the world; and (ii) to identify the parameters influencing the regional GPP_{MOD} .

2 Data and methods

2.1 Flux sites

The FLUXNET2015 Dataset provide the GPP product using eddy covariance flux tower measurement. In this study, GPP of seven maize sites from the FLUXNET2015 Dataset around the world were selected (Table 1, Fig. 1). More detailed descriptions of these sites can be obtained by the websites for Fluxdata and ChinaFLUX.

American sites including US_Ne1, US_Ne2, and US_Ne3 are large production fields. US_Ne1 and US_Ne2 sites are equipped with center pivot system for irrigation while the US_Ne3 site relies on rainfall. The irrigated sites (US_Ne1 and US_Ne2) have a long history of more than 10-years of maize-soybean rotation and no-till practice. The rain-fed site (US_Ne3) has a variable cultivation history with wheat, soybean, oat, and maize.

European site of DE-Kli is located 4 km south of the Tharandt Forest in Germany. This site has functioned solely as cropland since 1975. The eddy covariance measurements started in May 2004. The crop rotation was followed by rapeseed (2004/2005), winter wheat (2005/2006), maize (2007), and spring barley (2008).

Table 1 Characteristics of the study sites

Site	Site name	Country	Latitude	Longitude	Data period	Reference
US_Ne1	Mead-irrigated continuous maize site	USA	41.1651	96.4766 W	2001–2011	(Verma et al., 2005)
US_Ne2	Mead-irrigated maize-soybean rotation site	USA	41.1649	96.4701 W	2001–2011	(Verma et al., 2005)
US_Ne3	Mead-rainfed maize-soybean rotation site	USA	41.1797	96.4397 W	2001–2011	(Verma et al., 2005)
DE_Kli	Klingenberg	Germany	50.8929	13.5225 E	2007	(Gilmanov et al., 2010)
FR_Gri	Grignon	France	48.8442	1.9519 E	2005	(Lehuger et al., 2010)
CN_YC	Yucheng	China	36.8333	116.5667 E	2012–2013	(Sun et al., 2006)
CN_DM	Daman	China	38.8556	100.3722 E	2013–2014	(Wang et al., 2013)




Fig. 1 Locations of seven maize flux tower sites. The global land cover classification data were produced by the AVHRR (Hansen et al., 2000).

European site of FR-Gri lies in a large cropland field in a plateau situation close to a farm with cattle. The crop rotation here was followed by maize (2005), winter wheat (2005/2006), and barley (2007).

The Asian site of CN_DM is located in typical irrigated farmland in Daman village, Gansu Province, Northwest China, with a primary crop of maize (Tang et al., 2017). The precipitation in this site is about 100–250 mm every year with continental arid climate: dry and hot in summer and cold in winter. The Asian site of CN_YC lies in Yucheng County, Shangdong Province, North China, with a crop rotation of wheat and maize over one year. The annual mean temperature is about 13°C and the annual precipitation is approximately 528 mm.

2.2 MODIS data

MODIS, the main instrument aboard the Terra Earth Observing System (EOS) satellite for monitoring the seasonality of global terrestrial vegetation, was launched on 18 December 1999. Terra MODIS observes the entire Earth's surface with a period cycle of 1 to 2 days, obtaining data with 36 spectral bands. Beginning in 2000, GPP products were provided by the NASA EOS with a temporal resolution of 8 days at 1 km spatial resolution.

To evaluate the MOD17-GPP product with eddy covariance flux, the MOD15A2 and MOD17A2 products were obtained from the EOSDIS. The MOD15A2 data product is the 8-day composites of leaf area index (LAI) and FPAR, and the MOD17A2 is the 8-day composites of GPP and net primary productivity (NPP). The current

version of the above two products, the Collection 6 data at a spatial resolution of 500 m, were used in this study.

2.3 MOD17 algorithm

The MOD17A2 products are available by summing up the 8-day GPP. The description of the MODIS GPP algorithm was described by Running. The algorithm relies on the light use efficiency (ϵ) (Heinsch et al., 2006) linearly relating GPP to the absorbed photosynthetic radiation (APAR) (Monteith, 1972).

$$\text{GPP} = \text{PAR} \times \text{FPAR} \times \epsilon, \quad (1)$$

where PAR is the photosynthetically active radiation and FPAR is the fraction of the photosynthetic active radiation absorbed by vegetation. Estimates of 8-day mean daily FPAR with the spatial resolution of 500 m were provided by the MODIS team. The parameter of ϵ is the LUE for GPP.

$$\epsilon = \epsilon_{\max} \times T_{\min\text{scalar}} \times \text{VPD}_{\text{scalar}}, \quad (2)$$

where ϵ_{\max} is the maximum LUE from the look-up TABLE, relying on vegetation types. $T_{\min\text{scalar}}$ and $\text{VPD}_{\text{scalar}}$ are the scalars for the effects of the minimum temperature and vapor pressure deficit on LUE of vegetation, respectively. The parameters of VPD_{\max} and VPD_{\min} , $T_{\min\text{min}}$, $T_{\min\text{max}}$, and ϵ_{\max} can be acquired by the biome parameters look-up TABLE (BPLUT) in the user guide of MODIS17.

$$f(T_{\min}) = \begin{cases} 0, & T_{\min} < T_{\min_{\min}} \\ \frac{T_{\min} - T_{\min_{\min}}}{T_{\min_{\max}} - T_{\min_{\min}}}, & T_{\min_{\min}} < T_{\min} < T_{\min_{\max}} \\ 1, & T_{\min} > T_{\min_{\max}} \end{cases} \quad (3)$$

where, T_{\min} is the minimum daily temperature ($^{\circ}\text{C}$). $T_{\min_{\min}}$ is the lower limit of the daily minimum air temperature; and $T_{\min_{\max}}$ is the upper limit of the daily minimum air temperature.

$$f(\text{VPD}) = \begin{cases} 0, & (\text{VPD}) > (\text{VPD})_{\max} \\ \frac{(\text{VPD})_{\max} - (\text{VPD})}{(\text{VPD})_{\max} - (\text{VPD})_{\min}}, & (\text{VPD})_{\min} < (\text{VPD}) < (\text{VPD})_{\max} \\ 1, & (\text{VPD}) < (\text{VPD})_{\min} \end{cases} \quad (4)$$

where, VPD(Pa) is the average vapor pressure deficit. VPD_{\min} is the lower limit value of daytime mean vapor pressure deficit; and VPD_{\max} is the upper limit value of daytime mean vapor pressure deficit.

2.4 FPAR reconstruction

The temporal profile of FPAR should be smooth, as the result of the FPAR of the canopy changes slowly throughout the year. However, the FPAR from remote sensing data sometimes changes abruptly due to the noise of bad weather conditions such as clouds, persistent rainy days, and fog. To reduce the noise of the contaminated FPAR, a time-series reconstructing algorithm called the Savitzky_Golay filter (Chen et al., 2004) was employed in this study as follows:

$$Y_i^* = \frac{1}{2m+1} \sum_{j=-m}^{j=m} C_j Y_{i+j}, \quad (5)$$

where Y is the original time-series data; Y_i^* is the reconstructed time-series data; C_j is the j^{th} weight of the filter window; and $2m+1$ is the size of filter window (Ma

and Veroustraete, 2006).

2.5 Calibrating the LUE

According to the LUE model, the parameter of ε was calibrated using the following formula:

$$\varepsilon = \text{GPP} \times \text{FPAR}^{-1} \times \text{PAR}^{-1}, \quad (6)$$

where GPP is from the eddy covariance measurements. FPAR is from the MOD15 product and PAR is estimated from incident shortwave radiation (SWR) multiplied by 0.45. The maximum ε value was defined corresponding to the maximum GPP in the growing seasons in Eq. (6). In addition, the maximum LUE in each site are shown in Table 2.

2.6 Statistical indicator for validation

Three statistical indicators were used to assess the performance of the model goodness, including determination coefficient (R^2), root mean square error (RMSE), and the relative error (RE). They were calculated as follows:

$$R^2 = \left(\frac{\sum_{i=1}^N (\text{GPP}_{\text{EC}}(i) - \overline{\text{GPP}_{\text{EC}}}) (\text{GPP}_{\text{sim}}(i) - \overline{\text{GPP}_{\text{sim}}})}{\sqrt{\sum_{i=1}^N (\text{GPP}_{\text{EC}}(i) - \overline{\text{GPP}_{\text{EC}}})^2} \sqrt{\sum_{i=1}^N (\text{GPP}_{\text{sim}}(i) - \overline{\text{GPP}_{\text{sim}}})^2}} \right)^2, \quad (7)$$

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{i=1}^N (\text{GPP}_{\text{sim}}(i) - \text{GPP}_{\text{EC}}(i))^2}, \quad (8)$$

$$\text{RE} = \frac{\text{GPP}_{\text{sim}} - \text{GPP}_{\text{EC}}}{\text{GPP}_{\text{EC}}} \times 100\%, \quad (9)$$

where GPP_{sim} is the GPP calculated using the GPP_{MOD}

algorithm; GPP_{EC} is the tower measured GPP; the overbars represent the mean value; and N is the sample number.

3 Results

3.1 Validation of MOD17 GPP product

In the American and European sites, the MODIS GPP

Table 2 The calibrated LUE of seven maize sites

Site	US_Ne1	US_Ne2	US_Ne3	DE_Kli	FR_Gri	CN_DM	CN_YC
Max/(g C · MJ ⁻¹)	3.31	2.42	3.19	2.17	2.29	2.25	2.25

could not capture the beginning of the growing season of the maize. Additionally, there was a large underestimation in the MODIS GPP during the growing season in seven maize sites, as well as substantial biases in the non-growing seasons in the American and European sites (Fig. 2). In terms of the overall amount of the GPP, the agreements between GPP_{EC} and GPP_{MOD} changed in different sites with an R^2 from 0.45 to 0.93 (Fig. 3).

3.2 Improving MOD17 GPP product

To understand the errors of the GPP_{MOD} algorithm, three simulations were conducted. In the Simulation meteor_cor, only meteorological data such as PAR, VPD, and T were replaced by the observation value from the flux tower and other parameters (FPAR and ε_{max}) were default in the MODIS algorithm. In the Simulation LUE_cor, based on the Simulation meteor_cor, LUE was calibrated by the eddy covariance flux tower observation. In the Simulation FPAR_cor, based on the Simulation LUE_cor, the parameter of FPAR from the MOD15 was reconstructed to reduce the noise. The parameter details of the three

simulations are shown in Table 3.

Compared with GPP_{MOD} , simulation meteor_cor was replaced by the local meteorological data in the MODIS GPP algorithm, which only slightly improved the MODIS GPP at seven sites. However, Simulation LUE_cor greatly improved the amount of the MODIS GPP by using the ε calibrated by the eddy covariance flux tower observation. Meanwhile, the Simulation FPAR_cor could reduce the GPP noise due to the contaminated FPAR in the site of DE-Kli (Fig. 4) with GPP improvement from 1501.9 ($g C \cdot m^{-2} \cdot yr^{-1}$) to 1798.5 ($g C \cdot m^{-2} \cdot yr^{-1}$) and R^2 from 0.64 to 0.78. After improving the MODIS GPP algorithm step by step, from Simulation meteor_cor and Simulation LUE_cor to Simulation FPAR_cor, the amount of GPP increased markedly (Fig. 2 and Table 4) while R^2 between the simulation GPP and the observed GPP remains unchanged (Fig. 3 and Table 5).

From a statistical point of view, the simulation GPPs are indeed overestimated. However, concerning the pattern of the whole year, the simulated GPPs perfectly fit EC GPPs in the growing season of the maize. In the American and European sites, the FPAR was high in the non-growing




Fig. 2 The figure of Simulation meteor_cor(GPP_{meteor_cor}), Simulation LUE_cor(GPP_{LUE_cor}), Simulation FPAR_cor(GPP_{FPAR_cor}), GPP_{EC} , GPP_{MOD} at the seven sites. GPP_{meteor_cor} was calculated using the MODIS_GPP algorithm which was driven by the observed meteorological data (PAR, VPD, and T_{min}), FPAR(MOD15A2), and other default parameters; GPP_{LUE_cor} was calculated by the calibrated ε_0 values on the base of GPP_{meteor_cor} ; GPP_{FPAR_cor} was calculated with the reconstructed FPAR based on the GPP_{LUE_cor} ; GPP_{EC} was the eddy covariance flux tower observed GPP; and GPP_{MOD} was the MODIS GPP.




Fig. 3 The scatter plots between GPP_{MOD} , GPP_{LUE_cor} , and GPP_{EC} at seven maize eddy flux tower sites.

Table 3 Parameters used for the improving of MODIS GPP algorithm

GPP	FPAR	Meteorology data	Max	T_{min_max}	T_{min_min}	VPD_{max}	VPD_{min}
GPP MOD	MOD15 FPAR	DAO	1.004	12.02	-8.00	43	6.5
GPP meteor_cor	MOD15 FPAR	Surface measure	1.004	12.02	-8.00	43	6.5
GPP LUE_cor	MOD15 FPAR	Surface measure	Calibrated	12.02	-8.00	43	6.5
GPPFPAR_cor	reconstruction	Surface measure	Calibrated	12.02	-8.00	43	6.5

seasons, which leads to the high amount of the simulated GPPs (Fig. 2 and Fig. 5). The deviation of the FPAR in American and European sites may cause the high R^2 between the simulated GPP and EC GPP. However, there is no deviation in the CN_YC and CN_DM sites, and the R^2 is lower than those in American and European sites.

4 Discussion

4.1 Statistical characteristics of carbon fluxes across different regions

This study analyzed the statistical characteristics of carbon fluxes from maize croplands in different regions around the world, which provides valuable information to evaluate the carbon cycle in maize farmland ecosystems. The largest productivity of maize crops appeared in one of the American sites with GPP of 1774.6 g C/m²/yr, which had the largest mean LUE with the value of 2.97 g C/MJ. The YC site (a special explanation) had a mean GPP of 1676.3 g C/m²/yr with spring wheat (717.3 g C/m²/yr) and summer maize (959 g C/m²/yr). Deducting the GPP of

spring wheat from the CN_YC site, the smallest productivity of these maize crop sites was the CN_YC site, which was in the continental monsoon climate zone. However, the GPP of CN_DM site was slightly higher than that of the DE-Kli site but lower than those of US_Ne1, US_Ne2, and US_Ne3.

4.2 The maximum LUE and its uncertainty in the GPP

The maximum LUE, indicating the potential conversion efficiency of absorbed PAR under ideal vegetation growing conditions, has significance in the LUE model in the GPP simulation (Xiao et al., 2011). The maximum LUE was considered as a universal constant across plant function types in previous models (Potter et al., 1993). In the GPP_{MOD} algorithm, the default maximum LUE of crops is 1.044 g C/MJ, which contains all types of crops without consideration of C3 and C4. Maize is a C4 plant, and its maximum LUE was 2.66 g C/MJ, in the middle stream of the Heihe River basin (Wang et al., 2013). The maximum LUE of maize was determined as 2.84±0.57 g C/MJ by the flux tower data (Chen et al., 2014). In this study, the mean maximum LUE of these seven sites was 2.55 g C/MJ,




Fig. 4 FPAR and reconstructed FPAR (FPAR_SG) at seven flux sites.

calculated by the GPP_{EC} (measured by eddy covariance method), FPAR (MOD15A2) and PAR (meteorological measurements).

In these seven maize sites, the uncertainties of the underestimates were present in the LUE because the tower observing meteorological data only gently improves the GPP_{MOD} algorithm. However, after using the LUE calibrated by the eddy covariance flux tower data, the magnitude of GPP_{MOD} can be dramatically improved. Many previous works on the validation of MODIS GPP proved that the LUE was the primary reason for GPP underestimation (Turner et al., 2003, 2006; Fu et al., 2012).

4.3 The FPAR and its uncertainty on the GPP

4.3.1 The uncertainty of contaminated FPAR

It is apparent that the accuracy of the MODIS GPP product is highly reliant on the MODIS FPAR product and that the retrieval of FPAR under poor conditions with persistent cloud cover, fog, rainy weather, and low solar angles was extremely difficult (Coops et al., 2007). As a result, extraction of a high FPAR data for each of the 8-day time intervals can be problematic, leading to the uncertainty of the MODIS GPP. To obtain FPAR with no noise from

seven maize sites under conditions with low solar angles and persistent cloud cover is exceptionally challenging.

In this study, the noise of FPAR primarily appears at the DE-Kli and FR-Gri sites, which were in the ocean climate with a lot of cloudy and rainy weather (Fig. 4). According to our calculation results, the GPP in the ocean climate at the DE-Kli site was most affected by the noise of FPAR at 14%, followed by the FR-Gri site at 7%. After using the reconstructed FPAR as the input in Simulation $FPAR_{cor}$, the R^2 between the $GPP_{FPAR_{cor}}$ and GPP_{EC} was improved from 0.64 to 0.78 and 0.48 to 0.53, respectively (Table 5). Meanwhile, the monsoon climate can experience significant rainfall in the summer growing season, such as the CN-YC site. At CN-YC site, the R^2 between the $GPP_{FPAR_{cor}}$ and GPP_{EC} was from 0.76 to 0.83 with the RMSE from 16 to 18.66. In the continental climate of the American sites and CN-DM site, the GPP was slightly influenced by the contaminated FPAR.

In the absence of field measurements of FPAR, this study inferred the parameter of FPAR assimilated from the MODIS product relying on peer-reviewed literature. The prevalence of persistent cloud cover coupled with fog at high relative daily humidity in the growing season resulted in large uncertainty in the MODIS FPAR (Gebremichael and Barros, 2006). This study considered that the current MODIS algorithm, relying on DAO data for meteorologi-




Fig. 5 The relationship between the GPP_{EC} , GPP_{MOD} , and FPAR at seven maize eddy flux tower sites.

Table 4 Different GPPs from seven maize eddy covariance flux towers.

GPPs/(g C·m ⁻² ·yr ⁻¹)	US_Ne1	US_Ne2	US_Ne3	DE_Kli	FR_Gri	CN_DM	CN_YC
GPP_{MOD}	790.9	753.4	782.6	1066.8	933.4	700.8	710.9
GPP_{meteor_cor}	880.4	1066.4	814.6	719.8	1170.6	628.9	754.0
GPP_{LUE_cor}	2793.4	2472.0	2486.8	1501.9	2577.6	1355.4	1689.2
GPP_{FPAR_cor}	2815.5	2496.8	2496.7	1798.5	2703	1373.0	1706.3
GPP_{EC}	1707.3	1774.6	1550.3	1133.2	1283.4	1296.9	1676.3

Table 5 Statistical indices of different GPPs at seven maize eddy flux tower sites

Sites	GPP_{MOD}			GPP_{meteor_cor}			GPP_{LUE_cor}			GPP_{FPAR_cor}		
	RE/%	RMSE	R^2	RE/%	RMSE	R^2	RE/%	RMSE	R^2	RE/%	RMSE	R^2
US_Ne1	-53.7	50.3	0.77	-48.4	48.3	0.79	38.9	37.9	0.79	39.4	37.3	0.81
US_Ne2	-57.5	54.2	0.74	-39.9	40.37	0.90	28.2	26.1	0.90	28.9	25.8	0.91
US_Ne3	-49.5	48.2	0.76	-47.5	43.7	0.76	37.7	34.9	0.76	37.9	34.8	0.77
DE_Kli	-5.9	24.5	0.43	-36.5	23.0	0.65	24.5	23.1	0.64	36.9	24.3	0.78
FR_Gri	-27.3	33.5	0.43	-8.8	30.3	0.49	50.2	45.94	0.48	52.5	46.8	0.53
CN_DM	-45.9	27.1	0.93	-51.5	27.1	0.97	4.3	6.84	0.97	6.2	6.9	0.97
CN_YC	-57.6	34.7	0.73	-55	32.2	0.76	0.8	18.66	0.76	1.7	16	0.83

cal input, worked well in sunny days but raised challenges in complex weather, such as rainy and foggy days. Because regional weather was influenced by climate change, the

spatial patterns of cloud cover filled with fog and rainfall cannot be captured by the MODIS FPAR, which causes uncertainty and oscillation of the growing season GPP.

4.3.2 The uncertainty of canopy heterogeneity

Spatial heterogeneity of natural vegetation and land-surface affect the surface exchange of energy, water, and carbon, and the lower atmospheric circulation over a wide range of scales (Falge et al., 2002). The orientation and size of footprints vary remarkably according to the wind speed and direction from season to season (Chen et al., 2009). There is also the problem of the mismatch of the representation between the flux tower and the satellite observation of the GPP_{MOD} product. This study took no account of the footprint of observation in the eddy covariance flux tower because the original footprint was not big enough to be a pixel in the MODIS GPP products. Similarly to most studies, there are some areas to be improved in future studies, such as the footprint of eddy covariance flux.

With the comparison of the GPP_{EC} and GPP_{MOD} , simulated GPP have a systematic bias in the non-growing season in the American and European sites. In other words, GPP_{MOD} and simulated GPP cannot capture the start of the growing season but go into the growing season ahead of time. The bias of GPP_{MOD} and simulated GPP at the American and European sites correspond well with the FPAR from the MOD15A2H, which is an important parameter reflecting the condition of vegetation growth in the MODIS GPP algorithm. There are obvious reasons that the FPAR of the American and European sites mix maize with other vegetation. At the DE-Kli site, the landscapes cultivate evergreen forests coupled with a diversity of both annual crops, causing the FPAR to mix with forest. Meanwhile, the cropping systems, such as an alternative cropping system, make a diversity of the crops scatter near European and American sites, and the FPAR of some crops that start growing earlier than the maize was caught by the MOD15A2H. Therefore, canopy heterogeneity and a diversity of crops (due to alternative cropping) caused a high value of FPAR in the non-growing season of maize, which brought biases to the FPAR and uncertainty to the MODIS GPP.

However, in the Asian sites, a single corn crop was planted in wide areas in CN-DM and CN-YC sites. Therefore, the canopy heterogeneity was relatively small to ensure the FPAR without the interference of other crops, which made the MODIS GPP and Simulation GPP correctly capture the seasonal dynamics of maize growing in Asian sites.

4.4 The uncertainty of eddy covariance

The uncertainty of eddy covariance also exists, notably in the estimation of ecosystem respiration (Goulden et al., 1996) and interpolation errors caused by missing data. GPP is calculated as the net ecosystem exchange (NEE) plus ecosystem respiration (ER). As a result, various flux NEE partition methods will lead to different GPP amounts,

even at the same site (Reichstein et al., 2005). In this study, daytime respiration employs the Van't Hoff function (nighttime-based method: GPP_{NB}). With this method, the result can be affected by the suppression of the turbulence and dominance of advective fluxes at night (Lasslop et al., 2010).

5 Conclusions

In this study, the MODIS GPP product of maize is validated by the eddy covariance tower flux data at seven sites in America, Europe, and Asia. The MODIS GPP was underestimated by approximately 6% to 58%. The reasons for underestimation are as follows:

The marked influence of the accuracy of MODIS GPP was the LUE in each site. In the MODIS algorithm, the problem of the underestimate in LUE is a common phenomenon on the global scale. In the MODIS GPP algorithm, the maximum LUE of crops is defaulted at 1.044 g C/MJ, which contains all the types of crop without differentiation of C3 and C4. In fact, the mean calibration maximum LUE of these seven sites was 2.55 g C/MJ. The disparity between the default LUE in the MODIS GPP algorithm and the calibrated LUE from eddy flux tower are the primary reason for the underestimation of MODIS GPP.

In addition, the contaminated FPAR is a big contributor to the underestimate of MODIS GPP.

Meanwhile, in America and Europe, canopy heterogeneity and the diversity of crops caused by alternative cropping led to the deviation of catching the growing season of the maize crop, which introduces errors to the FPAR and uncertainty to the MODIS GPP.

When the MODIS GPP is applied to a specific area, users should consider the regional weather, consider the canopy heterogeneity and calibrate LUE from the eddy flux tower to minimize the noise of the FPAR and LUE for better accuracy of the MODIS GPP product. After improving the parameter of LUE and FPAR, the MODIS GPP product is applicable for global GPP calculations in temporal and spatial scales.

Acknowledgements This work is jointly supported by the National Natural Science Foundation of China (Grant No. 41771453), National Key Technology R&D Program of China (2016YFC0500106), Special Project of Science and Technology Basic Work (2014FY210800-5), and the Fundamental Research Funds for the Central Universities in China (SWU116088). We are grateful to the availability of the tower flux data of maize sites from Fluxnet and ChinaFlux.

References

- Chen B Z, Black T A, Coops N C, Hilker T, Trofymow J A, Morgenstern K (2009). Assessing tower flux footprint climatology and scaling between remotely sensed and eddy covariance measurements.

- Boundary-Layer Meteorol, 130(2): 137–167
- Chen J, Jönsson P, Tamura M, Gu Z, Matsushita B, Eklundh L (2004). A simple method for reconstructing a high-quality NDVI time-series data set based on the Savitzky–Golay filter. *Remote Sens Environ*, 91(3–4): 332–344
- Chen T, van der Werf G R, Gobron N, Moors E J, Dolman A J (2014). Global cropland monthly gross primary production in the year 2000. *Biogeosciences*, 11(14): 2365–2366
- Cohen W B, Maieringer T K, Yang Z Q, Gower S T, Turner D P, Ritts W D, Berterretche M, Running S W (2003). Comparisons of land cover and LAI estimates derived from ETM plus and MODIS for four sites in North America: a quality assessment of 2000/2001 provisional MODIS products. *Remote Sens Environ*, 88(3): 233–255
- Coops N C, Black T A, Jassal R S, Trofymow J A, Morgenstern K (2007). Comparison of MODIS, eddy covariance determined and physiologically modelled gross primary production (GPP) in a Douglas-fir forest stand. *Remote Sens Environ*, 107(3): 385–401
- Falge E, Baldocchi D, Tenhunen J, Aubinet M, Bakwin P, Berbigier P, Bernhofer C, Burba G, Clement R, Davis K J, Elbers J A, Goldstein A H, Grelle A, Granier A, Guðmundsson J, Hollinger D, Kowalski A S, Katul G, Law B E, Malhi Y, Meyers T, Monson R K, Munger J W, Oechel W, Paw U K T, Pilegaard K, Rannik Ü, Rebmann C, Suyker A, Valentini R, Wilson K, Wofsy S (2002). Seasonality of ecosystem respiration and gross primary production as derived from FLUXNET measurements. *Agric Meteorol*, 113(1–4): 53–74
- Fensholt R, Sandholt I, Rasmussen M S (2004). Evaluation of MODIS LAI, fAPAR and the relation between fAPAR and NDVI in a semi-arid environment using in situ measurements. *Remote Sens Environ*, 91(3–4): 490–507
- Fu G, Shen Z, Zhang X, Shi P, He Y, Zhang Y, Sun W, Wu J, Zhou Y, Pan X (2012). Calibration of MODIS-based gross primary production over an alpine meadow on the Tibetan Plateau. *Can J Rem Sens*, 38(2): 157–168
- Gebremichael M, Barros A P (2006). Evaluation of MODIS gross primary productivity (GPP) in tropical monsoon regions. *Remote Sens Environ*, 100(2): 150–166
- Gilmanov T G, Aires L, Barcza Z, Baron V S, Belelli L, Beringer J, Billesbach D, Bonal D, Bradford J, Ceschia E, Cook D, Corradi C, Frank A, Gianelle D, Gimeno C, Gruenwald T, Guo H, Hanan N, Haszpra L, Heilman J, Jacobs A, Jones M B, Johnson D A, Kiely G, Li S, Magliulo V, Moors E, Nagy Z, Nasyrov M, Owensby C, Pinter K, Pio C, Reichstein M, Sanz M J, Scott R, Soussana J F, Stoy P C, Svejcar T, Tuba Z, Zhou G (2010). Productivity, respiration, and light-response parameters of world grassland and agroecosystems derived from flux-tower measurements. *Rangeland Ecol Manag*, 63(1): 16–39
- Gitelson A A, Vina A, Masek J G, Verma S B, Suyker A E (2008). Synoptic monitoring of gross primary productivity of maize using Landsat data. *IEEE Geosci Remote Sens Lett*, 5(2): 133–137
- Goulden M L, Munger J W, Fan S, Daube B C, Wofsy S C (1996). Measurements of carbon sequestration by long-term eddy covariance: methods and a critical evaluation of accuracy. *Glob Change Biol*, 2(3): 169–182
- Hansen M C, Defries R S, Townshend J R G, Sohlberg R (2000). Global land cover classification at 1 km spatial resolution using a classification tree approach. *Int J Remote Sens*, 21(6–7): 1331–1364
- Heinsch F A, Zhao M, Running S W, Kimball J S, Nemani R R, Davis K J, Bolstad P V, Cook B D, Desai A R, Ricciuto D M, Law B E, Oechel W C, Kwon H, Luo H, Wofsy S C, Dunn A L, Munge J W, Baldocchi D D, Xu L, Hollinger D Y, Richardson A D, Stoy P C, Siqueira M B S, Monson R K, Burns S P, Flanagan L B (2006). Evaluation of remote sensing based terrestrial productivity from MODIS using regional tower eddy flux network observations. *IEEE Trans Geosci Remote Sens*, 44(7): 1908–1925
- Jung M, Reichstein M, Margolis H A, Cescatti A, Richardson A D, Arain M A, Arneth A, Bernhofer C, Bonal D, Chen J, Gianelle D, Gobron N, Kiely G, Kutsch W, Lasslop G, Law B E, Lindroth A, Merbold L, Montagnani L, Moors E J, Papale D, Sottocornola M, Vaccari F, Williams C (2015). Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations. *J Geophys Res*, 116(G3): 245–255
- Lai L, Huang X, Yang H, Chuai X, Zhang M, Zhong T, Chen Z, Chen Y, Wang X, Thompson J R (2016). Carbon emissions from land-use change and management in China between 1990 and 2010. *Sci Adv*, 2(11): e1611063
- Lasslop G, Reichstein M, Papale D, Richardson A D, Arneth A, Barr A, Stoy P, Wohlfahrt G (2010). Separation of net ecosystem exchange into assimilation and respiration using a light response curve approach: critical issues and global evaluation. *Glob Change Biol*, 16(1): 187–208
- Lehuger S, Gabrielle B, Cellier P, Loubet B, Roche R, Béziat P, Ceschia E, Wattenbach M (2010). Predicting the net carbon exchanges of crop rotations in Europe with an agro-ecosystem model. *Agric Ecosyst Environ*, 139(3): 384–395
- Ma M, Veroustraete F (2006). Reconstructing Pathfinder AVHRR land NDVI time-series data for the Northwest of China. *Advances in Space Research*, 37(4): 835–840
- Monteith J L (1972). Solar-radiation and productivity in tropical ecosystems. *J Appl Ecol*, 9(3): 747–766
- Peng Y, Gitelson A A (2011). Application of chlorophyll-related vegetation indices for remote estimation of maize productivity. *Agric Meteorol*, 151(9): 1267–1276
- Potter C S, Randerson J T, Field C B, Matson P A, Vitousek P M, Mooney H A, Klooster S A (1993). Terrestrial ecosystem production—A process model-based on global satellite and surface data. *Global Biogeochem Cycles*, 7(4): 811–841
- Reichstein M, Falge E, Baldocchi D, Papale D, Aubinet M, Berbigier P, Bernhofer C, Buchmann N, Gilmanov T, Granier A, Grünwald T, Havránková K, Ilvesniemi H, Janous D, Knohl A, Laurila T, Lohila A, Loustau D, Matteucci G, Meyers T, Miglietta F, Ourcival J M, Pumpanen J, Rambal S, Rotenberg E, Sanz M, Tenhunen J, Seufert G, Vaccari F, Vesala T, Yakir D, Valentini R (2005). On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm. *Glob Change Biol*, 11(9): 1424–1439
- Running S W, Baldocchi D D, Turner D P, Gower S T, Bakwin P S, Hibbard K A (1999). A global terrestrial monitoring network integrating tower fluxes, flask sampling, ecosystem modeling and EOS satellite data. *Remote Sens Environ*, 70(1): 108–127
- Running S W, Nemani R R, Heinsch F A, Zhao M S, Reeves M, Hashimoto H (2004). A continuous satellite-derived measure of

- global terrestrial primary production. *Bioscience*, 54(6): 547–560
- Sun X, Zhu Z, Wen X, Yuan G, Yu G (2006). The impact of averaging period on eddy fluxes observed at ChinaFLUX sites. *Agric Meteorol*, 137(3–4): 188–193
- Tang X, Li H, Huang N, Li X, Xu X, Ding Z, Xie J (2015). A comprehensive assessment of MODIS-derived GPP for forest ecosystems using the site-level FLUXNET database. *Environ Earth Sci*, 74(7): 5907–5918
- Tang X, Ma M, Ding Z, Xu X, Yao L, Huang X, Gu Q, Song L (2017). Remotely monitoring ecosystem water use efficiency of grassland and cropland in China's arid and semi-arid regions with MODIS data. *Remote Sens*, 9(6): 616
- Turner D P, Ritts W D, Cohen W B, Gower S T, Running S W, Zhao M, Costa M H, Kirschbaum A A, Ham J M, Saleska S R, Ahl D E (2006). Evaluation of MODIS NPP and GPP products across multiple biomes. *Remote Sens Environ*, 102(3–4): 282–292
- Turner D P, Ritts W D, Cohen W B, Gower S T, Zhao M, Running S W, Wofsy S C, Urbanski S, Dunn A L, Munger J W (2003). Scaling Gross Primary Production (GPP) over boreal and deciduous forest landscapes in support of MODIS GPP product validation. *Remote Sens Environ*, 88(3): 256–270
- Verma S B, Dobermann A, Cassman K G, Walters D T, Knops J M, Arkebauer T J, Suyker A E, Burba G G, Amos B, Yang H, Ginting D, Hubbard K G, Gitelson A A, Walter-Shea E A (2005). Annual carbon dioxide exchange in irrigated and rainfed maize-based agroecosystems. *Agric Meteorol*, 131(1–2): 77–96
- Wang X, Ma M, Li X, Song Y, Tan J, Huang G, Zhang Z, Zhao T, Feng J, Ma Z, Wei W, Bai Y (2013). Validation of MODIS-GPP product at 10 flux sites in northern China. *Int J Remote Sens*, 34(2): 587–599
- Xiao J, Davis K J, Urban N M, Keller K, Saliendra N Z (2011). Upscaling carbon fluxes from towers to the regional scale: influence of parameter variability and land cover representation on regional flux estimates. *J Geophys Res Biogeosci*, 116(G3): 115–132
- Zhang M, Huang X, Chuai X, Yang H, Lai L, Tan J (2015). Impact of land use type conversion on carbon storage in terrestrial ecosystems of China: a spatial-temporal perspective. *Sci Rep*, 5(1): 10233
- Zhang Y, Yu Q, Jiang J, Tang Y (2008). Calibration of Terra/MODIS gross primary production over an irrigated cropland on the North China Plain and an alpine meadow on the Tibetan Plateau. *Glob Change Biol*, 14(4): 757–767