

NFκB signaling regulates embryonic and adult neurogenesis

Yonggang ZHANG, Wenhui HU (✉)

Department of Neuroscience, Temple University School of Medicine, Philadelphia, PA 19140, USA

© Higher Education Press and Springer-Verlag Berlin Heidelberg 2012

Abstract Both embryonic and adult neurogenesis involves the self-renewal/proliferation, survival, migration and lineage differentiation of neural stem/progenitor cells. Such dynamic process is tightly regulated by intrinsic and extrinsic factors and complex signaling pathways. Misregulated neurogenesis contributes much to a large range of neurodevelopmental defects and neurodegenerative diseases. The signaling of NFκB regulates many genes important in inflammation, immunity, cell survival and neural plasticity. During neurogenesis, NFκB signaling mediates the effect of numerous niche factors such as cytokines, chemokines, growth factors, extracellular matrix molecules, but also cross-talks with other signaling pathways such as Notch, Shh, Wnt/β-catenin. This review summarizes current progress on the NFκB signaling in all aspects of neurogenesis, focusing on the novel role of NFκB signaling in initiating early neural differentiation of neural stem cells and embryonic stem cells.

Keywords neural stem cells, transcriptional factors, NFκB, neurogenesis, embryonic stem cells, signal transduction

Introduction

Neurogenesis is a process of neural cell formation and includes three terms based on the lineage differentiation into neurons, astrocytes and oligodendrocytes: neurogenesis, astrocytogenesis and oligodendrocytogenesis. Neurogenesis occurs not only during embryonic and perinatal development but also in adult nervous system including central (Conti and Cattaneo, 2010; Ming and Song, 2011), peripheral (Le Douarin et al., 2008; Pardal et al., 2010) and enteric nervous system (Heanue and Pachnis, 2007; Metzger, 2010). Defects or impairments in embryonic and neonatal neurogenesis contribute significantly to a large range of neuropsychiatric diseases including intellectual disability (mental retardation), autism spectrum disorders, childhood-onset schizophrenia, epilepsy, pediatric bipolar disorder, attention-deficit hyperactivity disorder, as well as genetic or neurodevelopmental disorders such as Fragile X syndrome, Rett syndrome, Down syndrome, Hirschsprung's disease (Heanue and Pachnis, 2007; Vaillend et al., 2008; Ma et al., 2009; Schäfer et al., 2009; Kishi and Macklis, 2010; Callan and Zarnescu, 2011). Adult neurogenesis is a relatively new concept and abnormal

adult neurogenesis plays an important role in the pathogenesis of many cognitive/mood dysfunction in adult such as depression (Hanson et al., 2011), schizophrenia (Inta et al., 2011), epilepsy (Andres-Mach et al., 2011), stroke (Zhang et al., 2011), HIV-associated neurocognitive disorders (Okamoto et al., 2007; Kaul, 2008) and neurodegenerative diseases including Alzheimer's disease (Mu and Gage, 2011), Parkinson's disease (van den Berge et al., 2011), multiple sclerosis (Huehnchen et al., 2011; Tepavčević et al., 2011), and others (Winner et al., 2011; Curtis et al., 2012).

Both embryonic and adult neurogenesis includes the self-renewal/proliferation, survival, migration and lineage differentiation of neural stem/progenitor cells (NSCs/NPCs). For better understanding, neurogenesis can be divided into early, middle and late phases though they are indistinguishable in most cases. Early neurogenesis includes self-renewal/proliferation of NSCs/NPCs and cell fate commitment. Middle neurogenesis covers the migration of lineage-restricted cells (neuroblast or glioblast) and terminal differentiation. Late neurogenesis includes the maturation and functional integrity of nascent cells. The duration of each phase depends upon the neurogenic region, developmental stage, and species. The progression from the early to late neurogenesis is tightly regulated by intrinsic and extrinsic factors and complex signaling pathways (Ma et al., 2009; Pathania et al., 2010). At the early stage, NSCs represent a very small population of quiescent, slow dividing cells while NPCs represent a large

Received April 10, 2012; accepted May 20, 2012

Correspondence: Wenhui HU

E-mail: whu@temple.edu

population of amplifying, rapid dividing cells (Encinas and Sierra, 2012). NSCs are the most primitive and normally quiescent neural precursor cells and are exceptionally sensitive to cosmic radiation (Encinas et al., 2008). Distinguishing NSCs from NPCs both *in vitro* and *in vivo* remains a big challenge (Bonaguidi et al., 2011; Siebzehnrubl et al., 2011).

Neurogenesis is regulated on various conceptual levels, from behavior, systemic response, specific nervous system activity, niche factors to intracellular/molecular regulation. Although many extrinsic factors and intrinsic epigenic/transcriptional proteins to regulate both embryonic and adult neurogenesis have been identified (Zhao et al., 2008; Conti and Cattaneo, 2010), the signaling pathways and molecular mechanisms remain poorly understood. In particular, what factors control the initiation of NSC differentiation into NPCs? After initial differentiation, what factors guide lineage-specific terminal differentiation? Several signaling pathways such as Wnt/ β -catenin, Notch, sonic hedgehog (Shh) as well as transcriptional factors such as Sox2, Pax6, Mash1, TLX, Hes1/5, NeuroD, Tbr2, etc. in neurogenesis have been identified (Ma et al., 2009; Hodge and Hevner, 2011), but little is known about the signaling of nuclear factor κ B (NF κ B) during neurogenesis (Schölzke and Schwaninger, 2007; Widera et al., 2008; Kaltschmidt and Kaltschmidt, 2009; Gutierrez and Davies, 2011; Zhang et al., 2011). Recently, inflammatory mediators including cytokines, chemokines, growth factors, adhesion molecules, are receiving increased attentions in the field of embryonic and adult neurogenesis because inflammatory and immune responses are known to play critical roles in various diseases and injuries of the nervous system (Das and Basu, 2008; Taupin, 2008; Widera et al., 2008; Whitney et al., 2009; Teng and Tang, 2010). Most inflammatory mediators act as both sources and targets of NF κ B signaling. The role of NF κ B signaling in regulating the proliferation/apoptosis of NSCs/NPCs, migration of neuroblast, maturation and plasticity of nascent neurons has been reviewed in several previous publications (Widera et al., 2006b; Schölzke and Schwaninger, 2007; Widera et al., 2008; Gutierrez and Davies, 2011; Zhang et al., 2011). In this review, we will focus on the role of NF κ B signaling in regulating selfrenewal and early differentiation of NSCs.

Signaling pathways of NF κ B activation

The transcriptional factor NF κ B plays a pivotal role in inflammation, immunity, cancer and neural plasticity (Häcker and Karin, 2006; Perkins, 2007). Constitutive and inducible activation of NF κ B has been reported in many types of human tumors and chronic diseases (Wong and Tergaonkar, 2009; Boyce et al., 2010; Lin et al., 2010; Mancino and Lawrence, 2010). Like other signaling transduction pathways, NF κ B is activated through a series of signaling cascades. The NF κ B

family contains 5 members including RelA(p65), RelB, c-Rel, p50/p105 (NF κ B1) and p52/p100 (NF κ B2), which form various combinations of homodimers or heterodimers (Chen and Greene, 2004; Perkins, 2007). All contain REL homology domain (RHD, 300 aa) responsible for DNA binding, dimerization, I κ B binding and nuclear translocation. The class I (p65, RelB, c-Rel) contains transactivation domain (TAD) and thus has intrinsic ability to activate transcription. The class II (p50 and p52) lacks TAD and thus their homodimers generally repress transcription of target genes but their heterodimers with class I have transcriptional activity. In non-stimulated cells, the NF κ B dimer is retained in the cytoplasm by the inhibitor of NF κ B (I κ B). Upon stimulation, I κ B is degraded via a phosphorylation-dependent proteasome-mediated mechanism and the released NF κ B is translocated to the nucleus where it binds to the κ B-sites and regulates the transcription of target genes. The degradation of I κ B is regulated by the I κ B kinase (IKK) that is activated by its upstream IKK kinases (Fig. 1). The I κ B family contains 8 members including the classical inhibitors I κ B α , I κ B β , I κ B γ (p105), I κ B δ (p100) and I κ B ϵ , as well as the atypical regulators Bcl-3, I κ B ξ and I κ Bns. All of them are characterized by the hallmark 5–7 ankyrin-repeats that mediate the interaction with NF κ B. The classical IKK complex contains 2 catalytic subunits IKK1/2 or IKK α / β and 1 regulatory subunit IKK γ (Chen and Greene, 2004; Perkins, 2007).

Three distinct signaling pathways for NF κ B activation have been identified: classical (canonical), non-classical (non-canonical, alternative) and atypical pathways (Viatour et al., 2005), all of them rely on sequentially activated kinases (Fig. 2). The classical pathway is a well-established original pathway and is crucial for the activation of innate immunity and inflammation. It involves the activation of the classical IKK complex (Häcker and Karin, 2006). This pathway generally regulates the activation of NF κ B complexes (e.g. p65/p50), in response to a wide range of stimuli such as pro-inflammatory cytokines tumor necrosis factor α (TNF α) and interleukin (IL) 1 β , Toll-like receptor agonists (LPS), antigens, or viruses (HTLV1, EBV). The activated IKK complex phosphorylates I κ B members (I κ B α , I κ B β , I κ B ϵ and p105) on a consensus motif DSGFxS (e.g. Ser-32/Ser-36 for I κ B α and Ser-19/Ser-23 for I κ B β) and the phosphorylated serines act as binding site for β -TrCP, the substrate recognition subunit of a Skp1-Cullin-F-box (SCF)-type E3 ubiquitin-protein ligase complex, named SCF $^{\beta$ -TrCP. This process, then, leads to the ubiquitination on specific lysine and the ubiquitinated I κ Bs are directed to 26S proteasome for full degradation, leaving free NF κ B complexes to enter into the nucleus. The kinetics of phosphorylation and degradation of I κ B β or I κ B ϵ are much slower than that of I κ B α and may reflect different substrate specificities of the IKK complex. The non-classical pathway is critical for the secondary lymphoid organ development, maturation of B cells and adaptive humoral immunity. It involves TRAF3-mediated




Figure 1 Signaling cascade of NFκB activation and identified members of each family. The red highlighted text denotes the best-studied original member(s) for each family.

activation of the NFκB-inducing kinase (NIK) and IKKα (Razani et al., 2011; Sun, 2012). Activated IKKα phosphorylates p100 on specific serine residues. After phosphorylation, p100 is ubiquitinated by SCF^{β-TrCP} E3 ligase and cleaved by 19S proteasome, instead of completely degraded by 26S proteasome, to generate the NFκB subunit p52. This process is generally slower than the activation of the classical pathway and leads to a delayed activation of nuclear p52-containing complexes, such as RelB/p52. The mechanisms of p52 generation are either constitutive (by cotranslational processing) or inducible (by post-translational cleavage). The non-classical pathway is triggered by some particular members of TNF family (Lymphotoxin β, B cell activation factor, CD40 ligand). Early studies demonstrated that almost all inducers of NFκB lead to activation of the classical IKKα/β/γ complex. Additional IKK complex, NIK/IKKα (Woronicz et al., 1997; Senftleben et al., 2001) and IKKβ/IKKγ (Quirling et al., 2004), has been identified. NIK/IKKα mediates the non-classical NFκB pathway while IKKβ/IKKγ subcomplex remains functional for the classical NFκB pathway. Our studies identified a novel NFκB signaling regulator, NIBP (NIK- and IKKβ binding protein) that forms a novel subcomplex comprised of NIK-NIBP-IKKβ without IKKα and IKKγ (Hu et al., 2005). Such subcomplex may represent a novel signaling pathway of NFκB activation and regulate protein transport and subcellular trafficking (Cox et al., 2007; Kümmel et al., 2008; Zong et al., 2011; Westlake et al., 2011; Zong et al., 2012).

Most stimuli activate NFκB through IKK-mediated IκB phosphorylation on N-terminal serine residues. A third signaling pathway for NFκB activation is classified as

atypical because it is independent of IKK (Fig. 2). Although the phosphorylation and proteasome-mediated degradation of IκB are involved, the phosphorylation sites are different. One atypical pathway is triggered by DNA damage such as UV or doxorubicin, leading to IκBα degradation via the proteasome, but the targeted serine residues are located within a C-terminal cluster, which is recognized by the p38 MAP kinase-activated casein kinase 2 (CK2) (Kato et al., 2003; Huillard et al., 2010). Another atypical pathway is stimulated by oxidative stress, which involves the activation of NADPH oxidase (NOX), generation of superoxide and hydrogen peroxide (H₂O₂) that leads to NFκB activation via phosphorylation of IκBα on the N-terminal Tyr-42 residue (Gloire et al., 2006). The upstream kinases for tyrosine phosphorylation of IκBα remain unclear, and the spleen tyrosine kinase (Syk) may be involved (Gallagher et al., 2007).

NFκB signaling in embryonic neurogenesis (neurodevelopment)

During early embryonic development, a subset of cells in the ectoderm overlying the notochord of the mesoderm proliferate/differentiate and become neural plate in response to the diffusible inhibitory signals (neural inducer) produced from the notochord. Neural plate folds to form the neural groove, which fuses to form the neural tube. Within the neural tube, NSCs (radial glia cells) derive from neuroepithelial cells (NECs) and differentiate sequentially into NPCs and various lineage-restricted precursor cells (RPCs), which include neuroblast and glioblast. These RPCs migrate to the marginal




Figure 2 Signaling pathways of NF κ B activation. In the classical pathway, the classical stimulators like TNF α bind to receptor, leading to recruitment of the adaptor molecule TRADD and the signaling proteins TRAF2 and RIP1. TRAF2 recruits the IKK complex that phosphorylates I κ B α at serine residue 32 and 36, leading to its ubiquitination and degradation by the proteasome, allowing the release of the classical heterodimer of p65/p50, which then translocate to the nucleus to regulate the transcription of target genes. In the non-classical pathway, NIK-induced phosphorylation of IKK α leads to the phosphorylation of p100 and proteasome-mediated processing of p100 into p52, allowing the nuclear translocation of RelB/p52 to regulate target gene expression. In the atypical pathway, the p38-activated casein kinase 2 (CK2) phosphorylates I κ B α at a cluster of serine residues within C-terminal domain, while oxidative stress induces spleen tyrosine kinase (Syk)-mediated phosphorylation of I κ B α at tyrosine residue (Y-42) within N-terminal domain.

zone and beyond to assume their terminal fate. The generation of different lineage occurs in a temporally distinct yet overlapping pattern. In rodents, neurogenesis peaks at embryonic day (E) 14, astrocytogenesis at postnatal day (P) 2, and oligodendrocytogenesis at P14 (Wang and Bordey, 2008; Sauvageot and Stiles, 2002).

The self-renewal and cell fate decision of NSCs during embryonic neurogenesis are regulated by various transcription factors and their signaling pathways including the nuclear hormone receptor TLX (tailless), the high-mobility-group transcription factor Sox2, the basic helix-loop-helix transcriptional factor Hes (hairy and enhancer of split), the tumor suppressor phosphatase Pten (phosphatase and tensin homolog deleted on chromosome 10), and the *Drosophila* membrane-associated protein Numb homologs, Numb and Numbl like (Shi et al., 2008). The Hes family plays key but opposing role in regulating neurodevelopment. Hes1 and Hes5 are activated by Notch signaling and repress the

expression of proneuronal factors such as Mash1, Neurogenin, Math and NeuroD (Cau et al., 2000; Kageyama et al., 2008). In contrast, Hes6 promotes neuronal differentiation but inhibits astrocyte differentiation (Fior and Henrique, 2005; Vilas-Boas and Henrique, 2010).

NF κ B is essential for embryonic development because p65 knockout mice died on E15 and p65/p50 or p65/c-rel double knockout mice died on E13 due to liver degeneration (Beg et al., 1995; Grossmann et al., 1999). Such embryonic lethality precluded further investigation on the role of NF κ B in late embryonic brain development. Additional knockout of TNF receptor 1 (TNFR1) in these p65 null mice rescued embryonic lethality (Alcamo et al., 2001), providing an opportunity to investigate the role of NF κ B signaling in regulating embryonic neurogenesis (Young et al., 2006). However, the distribution pattern of NSCs/NPCs and cell lineage analysis in neurogenic zones of these mutants have not yet been examined. IKK α /IKK β double knockout mice died on E12

due to apoptosis of NECs leading to impairments in neurogenesis (Li et al., 2000). In *Drosophila* (Chen et al., 2000; Ayyar et al., 2007; DeLotto et al., 2007; Reeves and Stathopoulos, 2009) and *Xenopus laevis* (Lake et al., 2001), the graded activation of NF κ B/c-rel protein in dorsal region determine the dorsal-ventral patterning during the very early embryonic development. During mouse embryogenesis, virtually all members of the NF κ B pathway are expressed in embryonic, trophoblastic, and uterine cells (Torchinsky and Toder, 2004). NF κ B may protect the embryos exposed to embryopathic stresses, possibly through its anti-apoptotic effect (Torchinsky and Toder, 2004).

Although little is known regarding the role of NF κ B in the early step of embryonic neurogenesis, several lines of evidence demonstrate that NF κ B is implicated in early differentiation of other stem cells. During murine spermatogenesis, NF κ B is activated in a stage-specific manner (Lilienbaum et al., 2000). During oocyte maturation and early embryonic development, NF κ B is activated (Nishikimi et al., 1999; Paciolla et al., 2011). In *Drosophila melanogaster*, the mRNA of the p65 homolog, named Dorsal, is maternally expressed and is concentrated in the egg cortex (Chen et al., 2000). In *Xenopus*, NF κ B activation is observed during oocyte maturation (Dominguez et al., 1993) and in late blastulae and gastrulae (Richardson et al., 1994). In mouse embryos, NF κ B activation is crucial to engage development beyond the 2-cell stage (Nishikimi et al., 1999). NF κ B mediates the neurogenic effect of erythropoietin in neurosphere cultures from E14 mouse ganglionic eminence (Shingo et al., 2001). Neural induction from cultured embryonic stem (ES) cells or induced pluripotent stem (iPS) cells has been established (Osakada and Takahashi, 2011; Denham et al., 2012), but the signaling mechanisms remain largely unknown. Such induction is an excellent *in vitro* model to recapture the embryonic neurogenesis. The signaling pathways identified during endogenous embryonic morphogenesis and neurogenesis can be applied to the neural induction and patterning, such as bone morphogenetic protein (BMP), fibroblast growth factor (FGF), Wnt, Shh and Notch signaling. We speculate that NF κ B signaling play an important role in the neuronal induction from ES or iPS cells. Recently, it has been shown that murine and human ES cells possess a low level of NF κ B activity that increases significantly during the differentiation process (Kang et al., 2007; Kim et al., 2008; Torres and Watt, 2008). In human ES cells, the classical NF κ B pathway regulates differentiation while the non-classical pathway maintains pluripotency (Yang et al., 2010). The transcription factor Nanog is essential in maintaining pluripotency of ES cells (Mitsui et al., 2003). During ES cell differentiation, endogenous NF κ B activity and target-gene expression increased. NF κ B inhibition increased expression of pluripotency markers. Nanog binds to NF κ B proteins, inhibits NF κ B activity and cooperates with Stat3 to maintain pluripotency (Torres and Watt, 2008). ES cell-specific miR-290 maintains the

pluripotency and self-renewal of ES cells through repressing NF κ B classical signaling (Lüningschrör et al., 2012). Forced expression of p65 causes loss of pluripotency, promotes differentiation of ES cells, and leads to an epithelial to mesenchymal transition. These data define p65 as a novel target gene of miR-290 cluster and provide new insight into the function of ES cell-specific miRNAs (Lüningschrör et al., 2012). Taken altogether, NF κ B signaling is activated and required during the early differentiation of various stem cells including ES/iPS, hematopoietic (Xiao et al., 2009; Montano-Almendras et al., 2012), mesenchymal (Hess et al., 2009; Cho et al., 2010) and cancer stem cells (Reikvam et al., 2009; Nogueira et al., 2011).

NF κ B signaling in adult neurogenesis

NF κ B signaling is well known to regulate cell survival or apoptosis in many cell types, depending upon the generation of its target genes. In adult nervous system, inflammation-related NF κ B signaling plays a sword-edge role after injuries or diseases. Neuronal NF κ B signaling is neuroprotective via its crucial role in maintaining neuronal survival, synaptogenesis, neural plasticity, learning and memory (Fridmacher et al., 2003; Mattson and Meffert, 2006; Camandola and Mattson, 2007; Boersma et al., 2011). A striking enrichment of phosphorylated I κ B α and IKK is observed in the axon initial segment (Schultz et al., 2006; Sanchez-Ponce et al., 2008) and the nodes of Ranvier (Politi et al., 2008), suggesting a novel role of NF κ B signaling in regulating axonal polarity and initial axonal formation. In the zones of active neurogenesis in both postnatal and adult mouse brain, various members of the NF κ B family are highly expressed (Denis-Donini et al., 2005), indicating for the first time that NF κ B is actively involved in the proliferation, migration and differentiation of adult NSCs/NPCs (Rolls et al., 2007; Widera et al., 2008). The presence of NF κ B in adult neurogenic zone is further validated using immunofluorescent microscopy (Meneghini et al., 2010; Zhang et al., 2012b). Early studies using κ B-dependent lacZ transgenic mice show that NF κ B activation in the central nervous system starts on day E12.5 and major changes are not observed until after birth, when the structures start their final maturation processes, but details about neurogenic zones are not reported (Schmidt-Ullrich et al., 1996). Recent studies in the hippocampal subgranular zone of adult NF κ B lacZ reporter mice have identified the activation of NF κ B signaling in NSCs instead of NPCs (Koo et al., 2010), suggesting a potential role of NF κ B signaling in early neurogenesis. Direct evidence for the *in vivo* effect of NF κ B signaling on the proliferation of NSCs/NPCs derives from p65 and p50 double knockout mice (Young et al., 2006) as well as overexpression of super inhibitor I κ B α mutant in NSCs/NPCs (Wu et al., 2000; Widera et al., 2006a; Rubio-Araiz et al., 2008; Zhu et al., 2008).

Little is known about the role of NF κ B signaling in regulating neural differentiation of NSCs/NPCs. In PC12 cells, both NIK and IKK β and their binding protein NIBP are required for nerve growth factor-induced neuronal differentiation via NF κ B signaling (Foehr et al., 2000; Azoitei et al., 2005; Hu et al., 2005), though several other signaling pathways are also involved (Wooten et al., 2000; Azoitei et al., 2005). In neuroblastoma cells, NF κ B promotes neuronal differentiation (Feng and Porter, 1999). Toll-like receptor 2 (TLR2) induces neuronal differentiation via protein kinase C (PKC)-dependent activation of NF κ B whereas TLR4 inhibits proliferation and neuronal differentiation of NPCs (Rolls et al., 2007). In p50-deficient mice, the neuronal differentiation of adult hippocampal NSCs is reduced by 50% while the proliferation does not change (Denis-Donini et al., 2008). In our recent study, we demonstrate that NF κ B signaling regulates the early differentiation of NSCs (Zhang et al., 2012b). During early differentiation, NF κ B signaling becomes active. Addition of TNF α to activate NF κ B signaling under proliferation conditions induces neural differentiation of NSCs/NPCs (Lou et al., 2003; Bernardino et al., 2008; Zhang et al., 2012b). TNF-like weak inducer of apoptosis (TWEAK) induces neuronal differentiation of NSCs/NPCs, under proliferation condition, through NF κ B-dependent downregulation of Hes1 that prevents neuronal differentiation (Schölzke et al., 2011). Selective inhibition of canonical NF κ B signaling by various pharmacologic inhibitors, shRNA and NSC-specific transgene dnIkB α retain the tripotential ability of differentiation and restore or enhance self-renewal capability of NSCs, suggesting that NF κ B signaling is essential for early neural differentiation (Zhang et al., 2012b). The critical role of NF κ B in the initial differentiation step of NSCs highlights a novel molecular mechanism for neurogenesis. Under physiologic conditions, moderate activation of NF κ B signaling promotes NSC differentiation into NPCs and maintains a continuous source for adult neurogenesis. In a mouse inducible IkB α transgenic model, NF κ B in NSCs/NPCs is necessary for axogenesis and maturation (Imielski et al., 2012). However, persistent and repeated overactivation of NF κ B signaling in NSCs may exhaust the NSC pool and thus lead to reduced neurogenesis as seen in aging patients (Villeda et al., 2011; Artegiani and Calegari, 2012; Encinas and Sierra, 2012) and chronic stress (Koo et al., 2010).

Cytokine-induced regulation of NF κ B signaling during neurogenesis

Several cytokines have been shown to regulate neurogenesis, such as TNF α , IL-1 β , IL-6, TWEAK, and IFN γ (Das and Basu, 2008; Whitney et al., 2009; Schölzke et al., 2011; Yirmiya and Goshen, 2011), and most of them stimulate NF κ B signaling. TNF α is extensively studied in affecting neurogenesis but the reports remain controversial. The

enhancing effect of TNF α on neuronal differentiation is first reported in rat mesencephalic NSCs (Lou et al., 2003). TNF α stimulation promotes the reaggregation of 3D neurospheres *in vitro* (Widera et al., 2006c). TNFR1-derived peptide promotes neuronal differentiation of the hippocampal NSCs (Kajiwara et al., 2005). However, TNF α has no effect on neuronal differentiation and neurite outgrowth of murine adult NSCs/NPCs (Wong et al., 2004; Keohane et al., 2010) and mouse ES-derived NPCs (Ideguchi et al., 2008). Other data suggest that TNF α promote gliogenesis but not neuronogenesis in the NSCs/NPCs (Ricci-Vitiani et al., 2006; Ideguchi et al., 2008; Johansson et al., 2008; Peng et al., 2008). Recent study demonstrates that TNF α -induced neuronal differentiation in NSCs/NPCs is dose-dependent (Bernardino et al., 2008). These opposite results regarding the effects of TNF α on neural differentiation may have resulted from the use of different protocols (culture media, coating matrix, and exposure time), passage numbers, brain regions, species, or doses (Keohane et al., 2010). In addition, these reports have targeted the late stages (from NPCs to RPCs and terminal cells) of neural differentiation. Using stemness assay and pharmacological approach, we demonstrate that TNF α treatment under proliferation conditions induced neural differentiation through NF κ B activation at a very early stage of neurogenesis. We also confirmed previous reports (Bernardino et al., 2008; Keohane et al., 2010) showing that TNF α controls the survival and neuronal differentiation of neural cells at some late stages during neural differentiation. The neurogenic effects of inflammatory cytokines highlight the important role of inflammatory and immune responses in early neurogenesis after injury or diseases of the nervous system (Rolls et al., 2007).

IL-1 β , another major proinflammatory cytokine, also plays an important role in neurogenesis. Several lines of evidence implicate that IL-1 β mediates stress-induced depression via inhibiting adult neurogenesis (Ben Menachem-Zidon et al., 2008; Goshen et al., 2008; Koo and Duman, 2008; Goshen and Yirmiya, 2009). Blockade of IL-1 β signaling could be a novel therapeutic approach for the treatment of depression (Koo and Duman, 2009). Further studies demonstrate a critical role of NF κ B signaling in mediating IL-1 β -induced antineurogenic effect by exhausting NSC pool (Koo et al., 2010). IL-1 β treatment decreases the proliferation and self-renewal of proliferating NSCs/NPCs cultured from embryonic rat hippocampus, but inhibits neuronal differentiation and promotes astroglial differentiation (Green et al., 2012), consistent with the effect of TNF α (Ricci-Vitiani et al., 2006; Ideguchi et al., 2008; Johansson et al., 2008; Peng et al., 2008).

Increasing evidence shows that another cytokine, IL-6, inhibits neuronogenesis but promotes gliogenesis both *in vitro* and *in vivo* (Okano, 2006; Nakanishi et al., 2007; Ideguchi et al., 2008; Johansson et al., 2008; Mukaino et al., 2008; Islam et al., 2009). However, interferon- γ is shown to inhibit the proliferation but promote neuronal differentiation

of NSCs (Ben-Hur et al., 2003; Wong et al., 2004; Ideguchi et al., 2008; Johansson et al., 2008; Lum et al., 2008).

Taken together, transient NF κ B activation by these cytokines under normal conditions or early stage of injuries or diseases may be beneficial by promoting NSC differentiation for neural repair or maintaining daily neurogenesis. However, chronic NF κ B activation may delete the NSC pool, leading to aberrant and insufficient neurogenesis (Koo et al., 2010; Artegiani and Calegari, 2012; Schwarz et al., 2012).

Non-classical and atypical NF κ B signaling pathways during neurogenesis

Most studies in terms of neurogenesis target the classical IKK β /I κ B α /p65 pathway of NF κ B activation (Widera et al., 2008; Kaltschmidt and Kaltschmidt, 2009). Little is known about the role of non-classical and atypical pathways in neurogenesis. RelB is expressed in migrating neuroblast, c-Rel is present in a few cells located at the edges of the rostral migratory stream, but only the classical p65/p50 is detectable in NSCs/NPCs (Denis-Donini et al., 2005). The p65 knockout mice dies on E15 (Beg et al., 1995), but RelB (Weih et al., 1997), c-Rel (Köntgen et al., 1995) or p50 (Sha et al., 1995) knockout mice are not embryonically lethal. There are no reports on the neurogenic pattern and lineage differentiation in the neurogenic zones of these knockout mice. In NSCs/NPCs, the pigment epithelium-derived factor (PEDF) induces a non-canonical activation of the NF κ B pathway, leading to the dismissal of the transcriptional co-repressor N-CoR from specific Notch-responsive promoters (Andreu-Agulló et al., 2009). However, several studies have shown that non-classical pathway plays important role in the differentiation of ES cells and other tissue stem cells. For example, the canonical NF κ B pathway regulates differentiation while the noncanonical pathway maintains the pluripotency of human ES cells (Yang et al., 2010). The p100/RelB signaling positively and intrinsically regulates self-renewal of hematopoietic stem/progenitor cell (Zhao et al., 2012). NIK/RelB signaling is required for osteoclast differentiation (Vaira et al., 2008). In mesenchymal stem cells, the non-classical NF κ B pathway promotes but the classical pathway inhibits the late and sustained CD69 expression (Saldanha-Araujo et al., 2011). LIGHT, a stimulator for non-classical NF κ B pathway, negatively regulates neurite growth from developing sensory neurons via inhibition of classical NF κ B pathway (Gavaldà et al., 2009). The mechanisms underlying the different and frequently opposing effects of the classical and non-classical NF κ B pathways in stem cell self-renewal and differentiation remain to be determined. NIBP may act as a switch between two pathways by binding to IKK β (classical) and NIK (non-classical) (Hu et al., 2005).

Although there is no direct evidence, we speculate that the atypical NF κ B pathway is involved in neurogenesis, because the key components of these atypical pathways have been

shown to regulate neurogenesis. NOX-mediated ROS is well known to activate NF κ B in various cell types (Gloire and Piette, 2009; Morgan and Liu, 2011). NSCs/NPCs maintain a higher level of ROS compared to differentiated neurons (Le Belle et al., 2011). Endogenous ROS and nitric oxide are essential for the proliferation and differentiation of NSCs/NPCs (Yoneyama et al., 2010; Le Belle et al., 2011). Controlled generation of ROS at low-to-moderate levels stimulates neural differentiation of NSCs/NPCs (Vieira et al., 2011; Kennedy et al., 2012). Numb-interacting protein 1 (Nip1), a component of the ROS-generating Duox system, directs neuronal differentiation of NSCs through ROS generation (Kennedy et al., 2010). The mechanisms underlying ROS-mediated neurogenesis may involve ROS-stimulated signaling pathways such as PI3K/Akt (Le Belle et al., 2011), p38 MAPK (Kim and Wong, 2009), ERK (Kim et al., 2011a), JNK (Yeo and Kang, 2007) and NF κ B signaling (Piao et al., 2005; Li et al., 2009; Qin and Crews, 2012). Syk mediates ciliary neurotrophic factor (CNTF)-induced tyrosine phosphorylation of I κ B α and promotes neurite growth of developing neurons (Gallagher et al., 2007; Gutierrez and Davies, 2011). Syk is involved in ephrinA-induced promotion of neuritogenesis (Richards et al., 2006; Angibaud et al., 2011). CK2 β functions as a positive regulator for NSC proliferation and multipotency (Ziercher et al., 2011) and controls oligodendrocytogenesis (Huillard et al., 2010).

Crosstalk between NF κ B and other signaling pathways

Many signaling pathways have been identified to regulate both embryonic and adult neurogenesis, including Notch, Shh, Wnt/ β -catenin, etc. (Mu et al., 2010). The functional interaction of these pathways with NF κ B signaling has been reported in neuron, microglia, astrocytes and other cells (Cao et al., 2011; Maniati et al., 2011; Zhang et al., 2012a). Such crosstalk may occur also to NSCs/NPCs (Bonini et al., 2011), but experimental evidence is needed. For example, Notch signaling is well known to maintain NSC self-renewal and multipotency, and thus it is speculated that NF κ B signaling may regulate NSC stemness through Notch signaling (Ang and Tergaonkar, 2007; Fujita et al., 2011). PEDF modulates Notch-dependent stemness of mouse NSCs through activation of non-canonical NF κ B pathway (Andreu-Agulló et al., 2009). In mouse mesencephalic neural crest cells, NF κ B signaling promotes gliogenesis through the interaction with Notch signaling (Fujita et al., 2011). In *Xenopus laevis*, NF κ B activates Shh signaling for anterior neural patterning (Lake et al., 2001). In mouse, Shh signaling regulates neural induction from ES cells (Cai et al., 2008) or NSCs (Dave et al., 2011). TLR3 regulates NSC proliferation by modulating Shh signaling (Yaddanapudi et al., 2011), perhaps through activation of NF κ B (Kasperczyk et al., 2009). Many studies have demonstrated the importance of Wnt/ β -catenin signaling

in neurogenesis (Kondo et al., 2011; Pei et al., 2012) and its interaction with NFκB signaling in cancer stem cells (Gavert et al., 2011; Pan et al., 2012) and mesenchymal stem cells (Hyun Hwa et al., 2008; Kim et al., 2011b). However, the crosstalk between NFκB and Wnt/β-catenin signaling during neural induction and neurogenesis need further investigation.

Correlation of NFκB dysfunction to neurodevelopmental and neurodegenerative diseases

The link of NFκB signaling defects to various neurodevelopmental disorders remains to be determined, though many immune and inflammatory responses have been implicated in these diseases and NFκB signaling mediates most actions of these immune/inflammatory factors. Among 6 genes associated with nonsyndromic autosomal-recessive mental retardation (Mir et al., 2009; Mochida et al., 2009; Philippe et al., 2009), two, NIBP (Mir et al., 2009; Mochida et al., 2009; Philippe et al., 2009) and CC2D1A (Noor et al., 2008; Zhao et al., 2010), have been shown to regulate NFκB signaling through the classical IKKβ pathway, implying the important role of NFκB signaling in mental retardation. In the cerebellum and front cortex from autism patients and animal model, IKKα expression is significantly increased, but the expression and activity of the classical NFκB signaling components such as IκBα and p65 do not display any significant change (Song et al., 2009), although the role of non-classical and atypical NFκB pathways need further investigation. In schizophrenia patients, NFκB activity is increased in peripheral blood mononuclear cells (Song et al., 2009) and the expression of NFκB-responsive genes is also increased (Hashimoto et al., 2011). Three single nucleotide polymorphism variants of the p65 gene are associated with schizophrenia in a Japanese population (Hashimoto et al., 2011).

NFκB signaling has been widely shown to regulate the pathogenesis of many neurodegenerative diseases such as Alzheimer's (Granic et al., 2009; Mu and Gage, 2011), Parkinson's (Ghosh et al., 2007; Sha et al., 2010), Huntington's (Marcora and Kennedy, 2010) diseases and multiple sclerosis (Huehnchen et al., 2011; Tepavčević et al., 2011), as well as traumatic or ischemic injuries of nervous system (Dong et al., 2011; Zhang et al., 2011). NFκB signaling has been targeted as a therapeutic strategy for neurodegenerative diseases (Camandola and Mattson, 2007; Mattson et al., 2000). On the other hand, adult neurogenesis has been shown to contribute to these neurodegenerative diseases (Mu and Gage, 2011; Winner et al., 2011). NFκB signaling acts as a double-edge sword during neurodegenerative disease, and the outcome depends upon the cell type, environmental factors and disease stages (Massa et al., 2006; Yang et al., 2007; Teng and Tang, 2010). As discussed above, NFκB signaling regulates all aspects of adult neurogenesis,

which may contribute to the development and progression of nervous system injury and diseases.

Concluding remark and future direction

NFκB signaling is a key mediator for numerous niche factors that regulate various stages or phases of embryonic and adult neurogenesis. The classical pathway of NFκB activation plays important role in regulating selfrenewal/multipotency and early differentiation of NSCs, as well as the proliferation/apoptosis of NPCs, migration of neuroblast, maturation and plasticity of nascent neurons. NFκB signaling is also important in regulating the early differentiation of other stem cells such as embryonic stem cells, hematopoietic stem cell, and mesenchymal stem cells. During neural induction both *in vitro* and *in vivo*, NFκB signaling is required. Further investigation of the upstream regulation and downstream mechanism underlying the essential role of NFκB signaling in initiating early differentiation of both neural induction and neurogenesis will open a potential avenue for the development of therapeutics for the treatment of neurodevelopmental disorders and neurodegenerative diseases. Emerging evidence suggests that non-classical and atypical NFκB pathways are implicated in ES cell differentiation. It will be important to evaluate the different role of three NFκB pathways during neurogenesis. NFκB signaling selectively regulates cell fate decision and lineage differentiation. The major barrier for NSC-based transplantation is the predominant gliogenesis *in vivo* when NSCs/NPCs encounter the complicate microenvironmental niches. Thus, selective blockade of gliogenesis and promotion of neuronal differentiation will be an important strategy for stem cell transplantation.

Acknowledgements

This work was supported by the National Institutes of Diabetes and Digestive and Kidney Diseases (WH, DK075964) and the National Center for Research Resources (WH, RR032123).

References

- Alcama E, Mizgerd J P, Horwitz B H, Bronson R, Beg A A, Scott M, Doerschuk C M, Hynes R O, Baltimore D (2001). Targeted mutation of TNF receptor I rescues the RelA-deficient mouse and reveals a critical role for NF-κB in leukocyte recruitment. *J Immunol*, 167(3): 1592–1600
- Andres-Mach M, Fike J R, Łuszczki J J (2011). Neurogenesis in the epileptic brain: a brief overview from temporal lobe epilepsy. *Pharmacol Rep*, 63(6): 1316–1323
- Andreu-Agulló C, Morante-Redolat J M, Delgado A C, Fariñas I (2009). Vascular niche factor PEDF modulates Notch-dependent stemness in the adult subependymal zone. *Nat Neurosci*, 12(12): 1514–1523
- Ang H L, Tergaonkar V (2007). Notch and NFκB signaling pathways: Do they collaborate in normal vertebrate brain development and

- function? *Bioessays*, 29(10): 1039–1047
- Angibaud J, Louveau A, Baudouin S J, Nerrière-Daguin V, Evain S, Bonnamain V, Hulin P, Csaba Z, Dournaud P, Thinarth R, Naveilhan P, Noraz N, Pellier-Monnin V, Boudin H (2011). The immune molecule CD3zeta and its downstream effectors ZAP-70/Syk mediate ephrin signaling in neurons to regulate early neurogenesis. *J Neurochem*, 119(4): 708–722
- Artegiani B, Calegari F (2012). Age-related cognitive decline: Can neural stem cells help us? *Aging (Albany NY)*, 4(3): 176–186
- Ayyar S, Pistillo D, Calleja M, Brookfield A, Gittins K, Goldstone C, Simpson P (2007). NF- κ B/Rel-mediated regulation of the neural fate in *Drosophila*. *PLoS ONE*, 2(11): e1178
- Azoitei N, Wirth T, Baumann B (2005). Activation of the I κ B kinase complex is sufficient for neuronal differentiation of PC12 cells. *J Neurochem*, 93(6): 1487–1501
- Beg A A, Sha W C, Bronson R T, Ghosh S, Baltimore D (1995). Embryonic lethality and liver degeneration in mice lacking the RelA component of NF- κ B. *Nature*, 376(6536): 167–170
- Ben-Hur T, Ben-Menachem O, Furer V, Einstein O, Mizrachi-Kol R, Grigoriadis N (2003). Effects of proinflammatory cytokines on the growth, fate, and motility of multipotential neural precursor cells. *Mol Cell Neurosci*, 24(3): 623–631
- Ben Menachem-Zidon O, Goshen I, Kreisel T, Ben Menahem Y, Reinhartz E, Ben Hur T, Yirmiya R (2008). Intrahippocampal transplantation of transgenic neural precursor cells overexpressing interleukin-1 receptor antagonist blocks chronic isolation-induced impairment in memory and neurogenesis. *Neuropsychopharmacology*, 33(9): 2251–2262
- Bernardino L, Agasse F, Silva B, Ferreira R, Grade S, Malva J O (2008). Tumor necrosis factor- α modulates survival, proliferation, and neuronal differentiation in neonatal subventricular zone cell cultures. *Stem Cells*, 26(9): 2361–2371
- Boersma M C, Dresselhaus E C, De Biase L M, Mihalas A B, Bergles D E, Meffert M K (2011). A requirement for nuclear factor- κ B in developmental and plasticity-associated synaptogenesis. *J Neurosci*, 31(14): 5414–5425
- Bonaguidi M A, Wheeler M A, Shapiro J S, Stadel R P, Sun G J, Ming G L, Song H (2011). *In vivo* clonal analysis reveals self-renewing and multipotent adult neural stem cell characteristics. *Cell*, 145(7): 1142–1155
- Bonini S A, Ferrari-Toninelli G, Uberti D, Montinaro M, Buizza L, Lanni C, Grilli M, Memo M (2011). Nuclear factor κ B-dependent neurite remodeling is mediated by Notch pathway. *J Neurosci*, 31(32): 11697–11705
- Boyce B F, Yao Z, Xing L (2010). Functions of nuclear factor κ B in bone. *Ann N Y Acad Sci*, 1192(1): 367–375
- Cai C, Thorne J, Grabel L (2008). Hedgehog serves as a mitogen and survival factor during embryonic stem cell neurogenesis. *Stem Cells*, 26(5): 1097–1108
- Callan M A, Zarnescu D C (2011). Heads-up: new roles for the fragile X mental retardation protein in neural stem and progenitor cells. *Genesis*, 49(6): 424–440
- Camandola S, Mattson M P (2007). NF- κ B as a therapeutic target in neurodegenerative diseases. *Expert Opin Ther Targets*, 11(2): 123–132
- Cao Q, Kaur C, Wu C Y, Lu J, Ling E A (2011). Nuclear factor- κ B regulates notch signaling in production of proinflammatory cytokines and nitric oxide in murine BV-2 microglial cells. *Neuroscience*, 192: 140–154
- Cau E, Gradwohl G, Casarosa S, Kageyama R, Guillemot F (2000). Hes genes regulate sequential stages of neurogenesis in the olfactory epithelium. *Development*, 127(11): 2323–2332
- Chen G, Handel K, Roth S (2000). The maternal NF- κ B/dorsal gradient of *Tribolium castaneum*: dynamics of early dorsoventral patterning in a short-germ beetle. *Development*, 127(23): 5145–5156
- Chen L F, Greene W C (2004). Shaping the nuclear action of NF- κ B. *Nat Rev Mol Cell Biol*, 5(5): 392–401
- Cho H H, Shin K K, Kim Y J, Song J S, Kim J M, Bae Y C, Kim C D, Jung J S (2010). NF- κ B activation stimulates osteogenic differentiation of mesenchymal stem cells derived from human adipose tissue by increasing TAZ expression. *J Cell Physiol*, 223(1): 168–177
- Conti L, Cattaneo E (2010). Neural stem cell systems: physiological players or *in vitro* entities? *Nat Rev Neurosci*, 11(3): 176–187
- Cox R, Chen S H, Yoo E, Segev N (2007). Conservation of the TRAPP-II-specific subunits of a Ypt/Rab exchanger complex. *BMC Evol Biol*, 7(1): 12
- Curtis MA, Low VF, Faull RL (2012). Neurogenesis and progenitor cells in the adult human brain: a comparison between hippocampal and subventricular progenitor proliferation. *Dev Neurobiol*, Online Available April 27, 2012
- Das S, Basu A (2008). Inflammation: a new candidate in modulating adult neurogenesis. *J Neurosci Res*, 86(6): 1199–1208
- Dave R K, Ellis T, Toumpas M C, Robson J P, Julian E, Adolphe C, Bartlett P F, Cooper H M, Reynolds B A, Wainwright B J (2011). Sonic hedgehog and notch signaling can cooperate to regulate neurogenic divisions of neocortical progenitors. *PLoS ONE*, 6(2): e14680
- DeLotto R, DeLotto Y, Steward R, Lippincott-Schwartz J (2007). Nucleocytoplasmic shuttling mediates the dynamic maintenance of nuclear Dorsal levels during *Drosophila* embryogenesis. *Development*, 134(23): 4233–4241
- Denham M, Parish C L, Leaw B, Wright J, Reid C A, Petrou S, Dottori M, Thompson L H (2012). Neurons derived from human embryonic stem cells extend long-distance axonal projections through growth along host white matter tracts after intra-cerebral transplantation. *Front Cell Neurosci*, 6: 11
- Denis-Donini S, Caprini A, Frassoni C, Grilli M (2005). Members of the NF- κ B family expressed in zones of active neurogenesis in the postnatal and adult mouse brain. *Brain Res Dev Brain Res*, 154(1): 81–89
- Denis-Donini S, Dellarole A, Crociara P, Francese M T, Bortolotto V, Quadrato G, Canonico P L, Orsetti M, Ghi P, Memo M, Bonini S A, Ferrari-Toninelli G, Grilli M (2008). Impaired adult neurogenesis associated with short-term memory defects in NF- κ B p50-deficient mice. *J Neurosci*, 28(15): 3911–3919
- Dominguez I, Sanz L, Arenzana-Seisdedos F, Diaz-Meco M T, Virelizier J L, Moscat J (1993). Inhibition of protein kinase C zeta subspecies blocks the activation of an NF- κ B-like activity in *Xenopus laevis* oocytes. *Mol Cell Biol*, 13(2): 1290–1295
- Dong J, Liu B, Song L, Lu L, Xu H, Gu Y (2011). Neural stem cells in the ischemic and injured brain: endogenous and transplanted. *Cell Tissue Bank Online Available Dec 21, 2011*
- Encinas J M, Sierra A (2012). Neural stem cell deforestation as the main force driving the age-related decline in adult hippocampal

- neurogenesis. *Behav Brain Res*, 227(2): 433–439
- Encinas J M, Vazquez M E, Switzer R C, Chamberland D W, Nick H, Levine H G, Scarpa P J, Enikolopov G, Steindler D A (2008). Quiescent adult neural stem cells are exceptionally sensitive to cosmic radiation. *Exp Neurol*, 210(1): 274–279
- Feng Z, Porter A G (1999). NF- κ B/Rel proteins are required for neuronal differentiation of SH-SY5Y neuroblastoma cells. *J Biol Chem*, 274(43): 30341–30344
- Fior R, Henrique D (2005). A novel *hes5/hes6* circuitry of negative regulation controls Notch activity during neurogenesis. *Dev Biol*, 281(2): 318–333
- Foehr E D, Bohuslav J, Chen L F, DeNoronha C, Gelezianas R, Lin X, O'Mahony A, Greene W C (2000). The NF- κ B-inducing kinase induces PC12 cell differentiation and prevents apoptosis. *J Biol Chem*, 275(44): 34021–34024
- Fridmacher V, Kaltschmidt B, Goudeau B, Ndiaye D, Rossi F M, Pfeiffer J, Kaltschmidt C, Israël A, Mémet S (2003). Forebrain-specific neuronal inhibition of nuclear factor- κ B activity leads to loss of neuroprotection. *J Neurosci*, 23(28): 9403–9408
- Fujita K, Yasui S, Shinohara T, Ito K (2011). Interaction between NF- κ B signaling and Notch signaling in gliogenesis of mouse mesencephalic neural crest cells. *Mech Dev*, 128(7–10): 496–509
- Gallagher D, Gutierrez H, Gavalda N, O'Keefe G, Hay R, Davies A M (2007). Nuclear factor- κ B activation via tyrosine phosphorylation of inhibitor κ B- α is crucial for ciliary neurotrophic factor-promoted neurite growth from developing neurons. *J Neurosci*, 27(36): 9664–9669
- Gavalda N, Gutierrez H, Davies A M (2009). Developmental regulation of sensory neurite growth by the tumor necrosis factor superfamily member LIGHT. *J Neurosci*, 29(6): 1599–1607
- Gavert N, Vivanti A, Hazin J, Brabletz T, Ben-Ze'ev A (2011). L1-mediated colon cancer cell metastasis does not require changes in EMT and cancer stem cell markers. *Mol Cancer Res*, 9(1): 14–24
- Ghosh A, Roy A, Liu X, Kordower J H, Mufson E J, Hartley D M, Ghosh S, Mosley R L, Gendelman H E, Pahan K (2007). Selective inhibition of NF- κ B activation prevents dopaminergic neuronal loss in a mouse model of Parkinson's disease. *Proc Natl Acad Sci USA*, 104(47): 18754–18759
- Gloire G, Legrand-Poels S, Piette J (2006). NF- κ B activation by reactive oxygen species: fifteen years later. *Biochem Pharmacol*, 72(11): 1493–1505
- Gloire G, Piette J (2009). Redox regulation of nuclear post-translational modifications during NF- κ B activation. *Antioxid Redox Signal*, 11(9): 2209–2222
- Goshen I, Kreisel T, Ben-Menachem-Zidon O, Licht T, Weidenfeld J, Ben-Hur T, Yirmiya R (2008). Brain interleukin-1 mediates chronic stress-induced depression in mice via adrenocortical activation and hippocampal neurogenesis suppression. *Mol Psychiatry*, 13(7): 717–728
- Goshen I, Yirmiya R (2009). Interleukin-1 (IL-1): a central regulator of stress responses. *Front Neuroendocrinol*, 30(1): 30–45
- Granic I, Dolga A M, Nijholt I M, van Dijk G, Eisel U L (2009). Inflammation and NF- κ B in Alzheimer's disease and diabetes. *J Alzheimers Dis*, 16(4): 809–821
- Green H F, Treacy E, Keohane A K, Sullivan A M, O'Keefe G W, Nolan Y M (2012). A role for interleukin-1 β in determining the lineage fate of embryonic rat hippocampal neural precursor cells. *Mol Cell Neurosci*, 49(3): 311–321
- Grossmann M, Metcalf D, Merryfull J, Beg A, Baltimore D, Gerondakis S (1999). The combined absence of the transcription factors Rel and RelA leads to multiple hemopoietic cell defects. *Proc Natl Acad Sci USA*, 96(21): 11848–11853
- Gutierrez H, Davies A M (2011). Regulation of neural process growth, elaboration and structural plasticity by NF- κ B. *Trends Neurosci*, 34(6): 316–325
- Häcker H, Karin M (2006). Regulation and function of IKK and IKK-related kinases. *Sci STKE*, 2006(357): re13
- Hanson N D, Owens M J, Nemeroff C B (2011). Depression, antidepressants, and neurogenesis: a critical reappraisal. *Neuropsychopharmacology*, 36(13): 2589–2602
- Hashimoto R, Ohi K, Yasuda Y, Fukumoto M, Yamamori H, Takahashi H, Iwase M, Okochi T, Kazui H, Saitoh O, Tatsumi M, Iwata N, Ozaki N, Kamijima K, Kunugi H, Takeda M (2011). Variants of the RELA gene are associated with schizophrenia and their startle responses. *Neuropsychopharmacology*, 36(9): 1921–1931
- Heanue T A, Pachnis V (2007). Enteric nervous system development and Hirschsprung's disease: advances in genetic and stem cell studies. *Nat Rev Neurosci*, 8(6): 466–479
- Hess K, Ushmorov A, Fiedler J, Brenner R E, Wirth T (2009). TNF α promotes osteogenic differentiation of human mesenchymal stem cells by triggering the NF- κ B signaling pathway. *Bone*, 45(2): 367–376
- Hodge R D, Hevner R F (2011). Expression and actions of transcription factors in adult hippocampal neurogenesis. *Dev Neurobiol*, 71(8): 680–689
- Hu W H, Pendergast J S, Mo X M, Brambilla R, Bracchi-Ricard V, Li F, Walters W M, Blits B, He L, Schaal S M, Bethea J R (2005). NIBP, a novel NIK and IKK(β)-binding protein that enhances NF- κ B activation. *J Biol Chem*, 280(32): 29233–29241
- Huehnchen P, Prozorovski T, Klaisle P, Lesemann A, Ingwersen J, Wolf S A, Kupsch A, Aktas O, Steiner B (2011). Modulation of adult hippocampal neurogenesis during myelin-directed autoimmune neuroinflammation. *Glia*, 59(1): 132–142
- Huillard E, Ziercher L, Blond O, Wong M, Deloulme J C, Souchelnytskyi S, Baudier J, Cochet C, Buchou T (2010). Disruption of CK2 β in embryonic neural stem cells compromises proliferation and oligodendrogenesis in the mouse telencephalon. *Mol Cell Biol*, 30(11): 2737–2749
- Hyun Hwa C, Hye Joon J, Ji Sun S, Yong Chan B, Jin Sup J (2008). Crossregulation of beta-catenin/Tcf pathway by NF- κ B is mediated by *lzts2* in human adipose tissue-derived mesenchymal stem cells. *Biochim Biophys Acta*, (3): 419–428
- Ideguchi M, Shinoyama M, Gomi M, Hayashi H, Hashimoto N, Takahashi J (2008). Immune or inflammatory response by the host brain suppresses neuronal differentiation of transplanted ES cell-derived neural precursor cells. *J Neurosci Res*, 86(9): 1936–1943
- Imielski Y, Schwamborn J C, Lüningschrör P, Heimann P, Holzberg M, Werner H, Leske O, Püschel A W, Memet S, Heumann R, Israel A, Kaltschmidt C, Kaltschmidt B (2012). Regrowing the adult brain: NF- κ B controls functional circuit formation and tissue homeostasis in the dentate gyrus. *PLoS ONE*, 7(2): e30838
- Inta D, Meyer-Lindenberg A, Gass P (2011). Alterations in postnatal neurogenesis and dopamine dysregulation in schizophrenia: a hypothesis. *Schizophr Bull*, 37(4): 674–680

- Islam O, Gong X, Rose-John S, Heese K (2009). Interleukin-6 and neural stem cells: more than gliogenesis. *Mol Biol Cell*, 20(1): 188–199
- Johansson S, Price J, Modo M (2008). Effect of inflammatory cytokines on major histocompatibility complex expression and differentiation of human neural stem/progenitor cells. *Stem Cells*, 26(9): 2444–2454
- Kageyama R, Ohtsuka T, Kobayashi T (2008). Roles of Hes genes in neural development. *Dev Growth Differ*, 50(Suppl 1): S97–S103
- Kajiwara K, Ogata S, Tanihara M (2005). Promotion of neurite outgrowth from fetal hippocampal cells by TNF- α receptor 1-derived peptide. *Cell Transplant*, 14(9): 665–672
- Kaltschmidt B, Kaltschmidt C (2009). NF- κ B in the nervous system. *Cold Spring Harb Perspect Biol*, 1(3): a001271
- Kang H B, Kim Y E, Kwon H J, Sok D E, Lee Y (2007). Enhancement of NF- κ B expression and activity upon differentiation of human embryonic stem cell line SNUhES3. *Stem Cells Dev*, 16(4): 615–624
- Kasperczyk H, Baumann B, Debatin K M, Fulda S (2009). Characterization of sonic hedgehog as a novel NF- κ B target gene that promotes NF- κ B-mediated apoptosis resistance and tumor growth *in vivo*. *FASEB J*, 23(1): 21–33
- Kato T Jr, Delhase M, Hoffmann A, Karin M (2003). CK2 Is a C-Terminal I κ B Kinase Responsible for NF- κ B Activation during the UV Response. *Mol Cell*, 12(4): 829–839
- Kaul M (2008). HIV's double strike at the brain: neuronal toxicity and compromised neurogenesis. *Front Biosci*, 13(13): 2484–2494
- Kennedy K A, Ostrakhovitch E A, Sandiford S D, Dayarathna T, Xie X, Waese E Y, Chang W Y, Feng Q, Skerjanc I S, Stanford W L, Li S S (2010). Mammalian numb-interacting protein 1/dual oxidase maturation factor 1 directs neuronal fate in stem cells. *J Biol Chem*, 285(23): 17974–17985
- Kennedy K A, Sandiford S D, Skerjanc I S, Li S S (2012). Reactive oxygen species and the neuronal fate. *Cell Mol Life Sci*, 69(2): 215–221
- Keohane A, Ryan S, Maloney E, Sullivan A M, Nolan Y M (2010). Tumour necrosis factor- α impairs neuronal differentiation but not proliferation of hippocampal neural precursor cells: Role of Hes1. *Mol Cell Neurosci*, 43(1): 127–135
- Kim J, Wong P K (2009). Loss of ATM impairs proliferation of neural stem cells through oxidative stress-mediated p38 MAPK signaling. *Stem Cells*, 27(8): 1987–1998
- Kim J H, Park S H, Park S G, Choi J S, Xia Y, Sung J H (2011a). The pivotal role of reactive oxygen species generation in the hypoxia-induced stimulation of adipose-derived stem cells. *Stem Cells Dev*, 20(10): 1753–1761
- Kim J M, Song J S, Cho H H, Shin K K, Bae Y C, Lee B J, Jung J S (2011b). Effect of the modulation of leucine zipper tumor suppressor 2 expression on proliferation of various cancer cells functions as a tumor suppressor. *Mol Cell Biochem*, 346(1–2): 125–136
- Kim Y E, Kang H B, Park J A, Nam K H, Kwon H J, Lee Y (2008). Upregulation of NF- κ B upon differentiation of mouse embryonic stem cells. *BMB Rep*, 41(10): 705–709
- Kishi N, Macklis J D (2010). MeCP2 functions largely cell-autonomously, but also non-cell-autonomously, in neuronal maturation and dendritic arborization of cortical pyramidal neurons. *Exp Neurol*, 222(1): 51–58
- Kondo T, Matsuo A J, Shimomura A, Koehler K R, Chan R J, Miller J M, Srour E F, Hashino E (2011). Wnt signaling promotes neuronal differentiation from mesenchymal stem cells through activation of Tlx3. *Stem Cells*, 29(5): 836–846
- Köntgen F, Grumont R J, Strasser A, Metcalf D, Li R, Tarlinton D, Gerondakis S (1995). Mice lacking the c-rel proto-oncogene exhibit defects in lymphocyte proliferation, humoral immunity, and interleukin-2 expression. *Genes Dev*, 9(16): 1965–1977
- Koo J W, Duman R S (2008). IL-1 β is an essential mediator of the antineurogenic and anhedonic effects of stress. *Proc Natl Acad Sci USA*, 105(2): 751–756
- Koo J W, Duman R S (2009). Evidence for IL-1 receptor blockade as a therapeutic strategy for the treatment of depression. *Curr Opin Investig Drugs*, 10(7): 664–671
- Koo J W, Russo S J, Ferguson D, Nestler E J, Duman R S (2010). Nuclear factor- κ B is a critical mediator of stress-impaired neurogenesis and depressive behavior. *Proc Natl Acad Sci USA*, 107(6): 2669–2674
- Kümmel D, Oeckinghaus A, Wang C, Krappmann D, Heinemann U (2008). Distinct isocomplexes of the TRAPP trafficking factor coexist inside human cells. *FEBS Lett*, 582(27): 3729–3733
- Lake B B, Ford R, Kao K R (2001). Xrel3 is required for head development in *Xenopus laevis*. *Development*, 128(2): 263–273
- Le Belle J E, Orozco N M, Paucar A A, Saxe J P, Mottahedeh J, Pyle A D, Wu H, Kornblum H I (2011). Proliferative neural stem cells have high endogenous ROS levels that regulate self-renewal and neurogenesis in a PI3K/Akt-dependant manner. *Cell Stem Cell*, 8(1): 59–71
- Le Douarin N M, Calloni G W, Dupin E (2008). The stem cells of the neural crest. *Cell Cycle*, 7(8): 1013–1019
- Li Q, Estepa G, Memet S, Israel A, Verma I M (2000). Complete lack of NF- κ B activity in IKK1 and IKK2 double-deficient mice: additional defect in neurulation. *Genes Dev*, 14(14): 1729–1733
- Li Q, Spencer N Y, Oakley F D, Buettner G R, Engelhardt J F (2009). Endosomal Nox2 facilitates redox-dependent induction of NF- κ B by TNF- α . *Antioxid Redox Signal*, 11(6): 1249–1263
- Lilienbaum A, Sage J, Mémet S, Rassoulzadegan M, Cuzin F, Israël A (2000). NF- κ B is developmentally regulated during spermatogenesis in mice. *Dev Dyn*, 219(3): 333–340
- Lin Y, Bai L, Chen W, Xu S (2010). The NF- κ B activation pathways, emerging molecular targets for cancer prevention and therapy. *Expert Opin Ther Targets*, 14(1): 45–55
- Lou S J, Gu P, Xu H, Xu X H, Wang M W, He C, Lu C L (2003). Effect of tumor necrosis factor- α on differentiation of mesencephalic neural stem cells and proliferation of oligodendrocytes in the rat. *Sheng Li Xue Bao*, 55(2): 183–186
- Lum M, Croze E, Wagner C, McLenachan S, Mitrovic B, Turnley A M (2008). Inhibition of neurosphere proliferation by IFN γ but not IFN β is coupled to neuronal differentiation. *J Neuroimmunol*, 206(1–2): 32–38
- Lüningschrör P, Stöcker B, Kaltschmidt B, Kaltschmidt C (2012). miR-290 cluster modulates pluripotency by repressing canonical NF- κ B signaling. *Stem Cells*, 30(4): 655–664
- Ma D K, Bonaguidi M A, Ming G L, Song H (2009). Adult neural stem cells in the mammalian central nervous system. *Cell Res*, 19(6): 672–682
- Mancino A, Lawrence T (2010). Nuclear factor- κ B and tumor-associated macrophages. *Clin Cancer Res*, 16(3): 784–789
- Maniati E, Bossard M, Cook N, Candido J B, Emami-Shahri N,

- Nedospasov S A, Balkwill F R, Tuveson D A, Hagemann T (2011). Crosstalk between the canonical NF- κ B and Notch signaling pathways inhibits Ppary expression and promotes pancreatic cancer progression in mice. *J Clin Invest*, 121(12): 4685–4699
- Marcora E, Kennedy M B (2010). The Huntington's disease mutation impairs Huntingtin's role in the transport of NF- κ B from the synapse to the nucleus. *Hum Mol Genet*, 19(22): 4373–4384
- Massa P T, Aleyasin H, Park D S, Mao X, Barger S W (2006). NF κ B in neurons? The uncertainty principle in neurobiology. *J Neurochem*, 97(3): 607–618
- Mattson M P, Culmsee C, Yu Z, Camandola S (2000). Roles of nuclear factor κ B in neuronal survival and plasticity. *J Neurochem*, 74(2): 443–456
- Mattson M P, Meffert M K (2006). Roles for NF- κ B in nerve cell survival, plasticity, and disease. *Cell Death Differ*, 13(5): 852–860
- Meneghini V, Francese M T, Carraro L, Grilli M (2010). A novel role for the receptor for advanced glycation end-products in neural progenitor cells derived from adult SubVentricular Zone. *Mol Cell Neurosci*, 45(2): 139–150
- Metzger M (2010). Neurogenesis in the enteric nervous system. *Arch Ital Biol*, 148(2): 73–83
- Ming G L, Song H (2011). Adult neurogenesis in the mammalian brain: significant answers and significant questions. *Neuron*, 70(4): 687–702
- Mir A, Kaufman L, Noor A, Motazacker M M, Jamil T, Azam M, Kahrizi K, Rafiq M A, Weksberg R, Nasr T, Naeem F, Tzschach A, Kuss A W, Ishak G E, Doherty D, Ropers H H, Barkovich A J, Najmabadi H, Ayub M, Vincent J B (2009). Identification of mutations in TRAPPC9, which encodes the NIK- and IKK-beta-binding protein, in nonsyndromic autosomal-recessive mental retardation. *Am J Hum Genet*, 85(6): 909–915
- Mitsui K, Tokuzawa Y, Itoh H, Segawa K, Murakami M, Takahashi K, Maruyama M, Maeda M, Yamanaka S (2003). The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell*, 113(5): 631–642
- Mochida G H, Mahajnah M, Hill A D, Basel-Vanagaite L, Gleason D, Hill R S, Bodell A, Crosier M, Straussberg R, Walsh C A (2009). A truncating mutation of TRAPPC9 is associated with autosomal-recessive intellectual disability and postnatal microcephaly. *Am J Hum Genet*, 85(6): 897–902
- Montano-Almendras C P, Essaghir A, Schoemans H, Varis I, Noel LA, Velghe AI, Latinne D, Knoops L, Demoulin J B (2012). ETV6-PDGFRB and FIP1L1-PDGFR A stimulate human hematopoietic progenitor proliferation and differentiation into eosinophils: role of NF- κ B. *Haematologica*, Online Available Jan 22, 2012
- Morgan M J, Liu Z G (2011). Crosstalk of reactive oxygen species and NF- κ B signaling. *Cell Res*, 21(1): 103–115
- Mu Y, Gage F H (2011). Adult hippocampal neurogenesis and its role in Alzheimer's disease. *Mol Neurodegener*, 6(1): 85
- Mu Y, Lee S W, Gage F H (2010). Signaling in adult neurogenesis. *Curr Opin Neurobiol*, 20(4): 416–423
- Mukaino M, Nakamura M, Okada S, Toyama Y, Liu M, Okano H (2008). [Role of IL-6 in regulation of inflammation and stem cell differentiation in CNS trauma]. *Nihon Rinsho Meneki Gakkai Kaishi*, 31(2): 93–98 (Role of IL-6 in regulation of inflammation and stem cell differentiation in CNS trauma)
- Nakanishi M, Niidome T, Matsuda S, Akaike A, Kihara T, Sugimoto H (2007). Microglia-derived interleukin-6 and leukaemia inhibitory factor promote astrocytic differentiation of neural stem/progenitor cells. *Eur J Neurosci*, 25(3): 649–658
- Nishikimi A, Mukai J, Yamada M (1999). Nuclear translocation of nuclear factor κ B in early 1-cell mouse embryos. *Biol Reprod*, 60(6): 1536–1541
- Nogueira L, Ruiz-Ontañón P, Vazquez-Barquero A, Lafarga M, Berciano M T, Aldaz B, Grande L, Casafont I, Segura V, Robles E F, Suarez D, Garcia L F, Martinez-Climent J A, Fernandez-Luna J L (2011). Blockade of the NF κ B pathway drives differentiating glioblastoma-initiating cells into senescence both *in vitro* and *in vivo*. *Oncogene*, 30(32): 3537–3548
- Noor A, Windpassinger C, Patel M, Stachowiak B, Mikhailov A, Azam M, Irfan M, Siddiqui Z K, Naeem F, Paterson A D, Lutfullah M, Vincent J B, Ayub M (2008). CC2D2A, encoding a coiled-coil and C2 domain protein, causes autosomal-recessive mental retardation with retinitis pigmentosa. *Am J Hum Genet*, 82(4): 1011–1018
- Okamoto S, Kang Y J, Brechtel C W, Siviglia E, Russo R, Clemente A, Harrop A, McKercher S, Kaul M, Lipton S A (2007). HIV/gp120 decreases adult neural progenitor cell proliferation via checkpoint kinase-mediated cell-cycle withdrawal and G1 arrest. *Cell Stem Cell*, 1(2): 230–236
- Okano H (2006). Adult neural stem cells and central nervous system repair. *Ernst Schering Res Found Workshop*, 60: 215–228
- Osakada F, Takahashi M (2011). Neural induction and patterning in Mammalian pluripotent stem cells. *CNS Neurol Disord Drug Targets*, 10(4): 419–432
- Paciolla M, Boni R, Fusco F, Pescatore A, Poeta L, Ursini M V, Lioi M B, Miano M G (2011). Nuclear factor- κ B-inhibitor alpha (NFKBIA) is a developmental marker of NF- κ B/p65 activation during *in vitro* oocyte maturation and early embryogenesis. *Hum Reprod*, 26(5): 1191–1201
- Pan J X, Ding K, Wang C Y (2012). Niclosamide, an old antihelminthic agent, demonstrates antitumor activity by blocking multiple signaling pathways of cancer stem cells. *Chin J Cancer*, Online Available Jan 9, 2012
- Pardal R, Ortega-Sáenz P, Durán R, Platero-Luengo A, López-Barneo J (2010). The carotid body, a neurogenic niche in the adult peripheral nervous system. *Arch Ital Biol*, 148(2): 95–105
- Pathania M, Yan L D, Bordey A (2010). A symphony of signals conducts early and late stages of adult neurogenesis. *Neuropharmacology*, 58(6): 865–876
- Pei Y, Brun S N, Markant S L, Lento W, Gibson P, Taketo M M, Giovannini M, Gilbertson R J, Wechsler-Reya R J (2012). WNT signaling increases proliferation and impairs differentiation of stem cells in the developing cerebellum. *Development*, 139(10): 1724–1733
- Peng H, Whitney N, Wu Y, Tian C, Dou H, Zhou Y, Zheng J (2008). HIV-1-infected and/or immune-activated macrophage-secreted TNF- α affects human fetal cortical neural progenitor cell proliferation and differentiation. *Glia*, 56(8): 903–916
- Perkins N D (2007). Integrating cell-signalling pathways with NF- κ B and IKK function. *Nat Rev Mol Cell Biol*, 8(1): 49–62
- Philippe O, Rio M, Carioux A, Plaza J M, Guigue P, Molinari F, Bodaert N, Bole-Feysot C, Nitschke P, Smahi A, Munnich A, Colleaux L (2009). Combination of linkage mapping and microarray-expression analysis identifies NF- κ B signaling defect as a cause of

- autosomal-recessive mental retardation. *Am J Hum Genet*, 85(6): 903–908
- Piao Y J, Seo Y H, Hong F, Kim J H, Kim Y J, Kang M H, Kim B S, Jo S A, Jo I, Jue D M, Kang I, Ha J, Kim S S (2005). Nox 2 stimulates muscle differentiation via NF- κ B/iNOS pathway. *Free Radic Biol Med*, 38(8): 989–1001
- Politi C, Del Turco D, Sie J M, Golinski P A, Tegeder I, Deller T, Schultz C (2008). Accumulation of phosphorylated I κ B alpha and activated IKK in nodes of Ranvier. *Neuropathol Appl Neurobiol*, 34(3): 357–365
- Qin L, Crews F T (2012). NADPH oxidase and reactive oxygen species contribute to alcohol-induced microglial activation and neurodegeneration. *J Neuroinflammation*, 9(1): 5
- Quirling M, Page S, Jilg N, Plenagl K, Peus D, Grubmüller C, Weingärtner M, Fischer C, Neumeier D, Brand K (2004). Detection of IKKbeta-IKKgamma subcomplexes in monocytic cells and characterization of associated signaling. *J Biol Chem*, 279(36): 37452–37460
- Razani B, Reichardt A D, Cheng G (2011). Non-canonical NF- κ B signaling activation and regulation: principles and perspectives. *Immunol Rev*, 244(1): 44–54
- Reeves G T, Stathopoulos A (2009). Graded dorsal and differential gene regulation in the *Drosophila* embryo. *Cold Spring Harb Perspect Biol*, 1(4): a000836
- Reikvam H, Olsnes A M, Gjertsen B T, Ersvar E, Bruserud O (2009). Nuclear factor- κ B signaling: a contributor in leukemogenesis and a target for pharmacological intervention in human acute myelogenous leukemia. *Crit Rev Oncog*, 15(1–2): 1–41
- Ricci-Vitiani L, Casalbore P, Petrucci G, Lauretti L, Montano N, Larocca L M, Falchetti M L, Lombardi D G, Gerevini V D, Cenciarelli C, D'Alessandris Q G, Fernandez E, De Maria R, Maira G, Peschle C, Parati E, Pallini R (2006). Influence of local environment on the differentiation of neural stem cells engrafted onto the injured spinal cord. *Neurol Res*, 28(5): 488–492
- Richards G R, Smith A J, Cudnon P, Ma Q P, Leveridge M, Kerby J, Roderick H L, Bootman M D, Simpson P B (2006). The JAK3 inhibitor WHI-P154 prevents PDGF-evoked process outgrowth in human neural precursor cells. *J Neurochem*, 97(1): 201–210
- Richardson J C, Garcia Estrabot A M, Woodland H R (1994). XrelA, a *Xenopus* maternal and zygotic homologue of the p65 subunit of NF- κ B. Characterisation of transcriptional properties in the developing embryo and identification of a negative interference mutant. *Mech Dev*, 45(2): 173–189
- Rolls A, Shechter R, London A, Ziv Y, Ronen A, Levy R, Schwartz M (2007). Toll-like receptors modulate adult hippocampal neurogenesis. *Nat Cell Biol*, 9(9): 1081–1088
- Rubio-Araiz A, Arévalo-Martín A, Gómez-Torres O, Navarro-Galve B, García-Ovejero D, Suetterlin P, Sánchez-Heras E, Molina-Holgado E, Molina-Holgado F (2008). The endocannabinoid system modulates a transient TNF pathway that induces neural stem cell proliferation. *Mol Cell Neurosci*, 38(3): 374–380
- Saldanha-Araujo F, Haddad R, Malmegrim de Farias K C, Alves Souza AD, Palma P V, Araujo A G, Orellana M D, Voltarelli J C, Covas D T, Zago M A, Panepucci R A (2011). Mesenchymal stem cells promote the sustained expression of CD69 on activated T-lymphocytes: roles of canonical and non-canonical NF- κ B signaling. *J Cell Mol Med*, Online Available July 21, 2011
- Sanchez-Ponce D, Tapia M, Muñoz A, Garrido J J (2008). New role of IKK alpha/beta phosphorylated I κ B alpha in axon outgrowth and axon initial segment development. *Mol Cell Neurosci*, 37(4): 832–844
- Sauvageot C M, Stiles C D (2002). Molecular mechanisms controlling cortical gliogenesis. *Curr Opin Neurobiol*, 12(3): 244–249
- Schäfer K H, Micci M A, Pasricha P J (2009). Neural stem cell transplantation in the enteric nervous system: roadmaps and roadblocks. *Neurogastroenterol Motil*, 21(2): 103–112
- Schmidt-Ullrich R, Mémet S, Lilienbaum A, Feuillard J, Raphaël M, Israel A (1996). NF- κ B activity in transgenic mice: developmental regulation and tissue specificity. *Development*, 122(7): 2117–2128
- Schölzke M N, Röttinger A, Murikinati S, Gehrig N, Leib C, Schwaninger M (2011). TWEAK regulates proliferation and differentiation of adult neural progenitor cells. *Mol Cell Neurosci*, 46(1): 325–332
- Schölzke M N, Schwaninger M (2007). Transcriptional regulation of neurogenesis: potential mechanisms in cerebral ischemia. *J Mol Med (Berl)*, 85(6): 577–588
- Schultz C, König H G, Del Turco D, Politi C, Eckert G P, Ghebremedhin E, Prehn J H, Kögel D, Deller T (2006). Coincident enrichment of phosphorylated I κ Balpha, activated IKK, and phosphorylated p65 in the axon initial segment of neurons. *Mol Cell Neurosci*, 33(1): 68–80
- Schwarz T J, Ebert B, Lie D C (2012). Stem cell maintenance in the adult mammalian hippocampus: A matter of signal integration? *Dev Neurobiol*, Online Available April 5, 2012
- Senftleben U, Cao Y, Xiao G, Greten F R, Krähn G, Bonizzi G, Chen Y, Hu Y, Fong A, Sun S C, Karin M (2001). Activation by IKKalpha of a second, evolutionary conserved, NF- κ B signaling pathway. *Science*, 293(5534): 1495–1499
- Sha D, Chin L S, Li L (2010). Phosphorylation of parkin by Parkinson disease-linked kinase PINK1 activates parkin E3 ligase function and NF- κ B signaling. *Hum Mol Genet*, 19(2): 352–363
- Sha W C, Liou H C, Tuomanen E I, Baltimore D (1995). Targeted disruption of the p50 subunit of NF- κ B leads to multifocal defects in immune responses. *Cell*, 80(2): 321–330
- Shi Y, Sun G, Zhao C, Stewart R (2008). Neural stem cell self-renewal. *Crit Rev Oncol Hematol*, 65(1): 43–53
- Shingo T, Sorokan S T, Shimazaki T, Weiss S (2001). Erythropoietin regulates the *in vitro* and *in vivo* production of neuronal progenitors by mammalian forebrain neural stem cells. *J Neurosci*, 21(24): 9733–9743
- Siebzehnrub F A, Vedam-Mai V, Azari H, Reynolds B A, Deleyrolle L P (2011). Isolation and characterization of adult neural stem cells. *Methods Mol Biol*, 750: 61–77
- Song X Q, Lv L X, Li W Q, Hao Y H, Zhao J P (2009). The interaction of nuclear factor- κ B and cytokines is associated with schizophrenia. *Biol Psychiatry*, 65(6): 481–488
- Sun S C (2012). The noncanonical NF- κ B pathway. *Immunol Rev*, 246(1): 125–140
- Taupin P (2008). Adult neurogenesis, neuroinflammation and therapeutic potential of adult neural stem cells. *Int J Med Sci*, 5(3): 127–132
- Teng F Y, Tang B L (2010). NF- κ B signaling in neurite growth and neuronal survival. *Rev Neurosci*, 21(4): 299–314
- Tepavčević V, Lazarini F, Alfaro-Cervello C, Kerninon C, Yoshikawa K, Garcia-Verdugo J M, Lledo P M, Nait-Oumesmar B, Baron-Van Evercooren A (2011). Inflammation-induced subventricular zone

- dysfunction leads to olfactory deficits in a targeted mouse model of multiple sclerosis. *J Clin Invest*, 121(12): 4722–4734
- Torchinsky A, Toder V (2004). To die or not to die: the function of the transcription factor NF- κ B in embryos exposed to stress. *Am J Reprod Immunol*, 51(2): 138–143
- Torres J, Watt F M (2008). Nanog maintains pluripotency of mouse embryonic stem cells by inhibiting NF- κ B and cooperating with Stat3. *Nat Cell Biol*, 10(2): 194–201
- Vaillend C, Poirier R, Laroche S (2008). Genes, plasticity and mental retardation. *Behav Brain Res*, 192(1): 88–105
- Vaira S, Johnson T, Hirbe A C, Alhawagri M, Anwisyte I, Sammut B, O'Neal J, Zou W, Weilbaecher K N, Faccio R, Novack D V (2008). RelB is the NF- κ B subunit downstream of NIK responsible for osteoclast differentiation. *Proc Natl Acad Sci USA*, 105(10): 3897–3902
- van den Berge S A, van Strien M E, Korecka J A, Dijkstra A A, Sluijs J A, Kooijman L, Eggers R, De Filippis L, Vescovi A L, Verhaagen J, van de Berg W D, Hol E M (2011). The proliferative capacity of the subventricular zone is maintained in the parkinsonian brain. *Brain*, 134(Pt 11): 3249–3263
- Viatour P, Merville M P, Bours V, Chariot A (2005). Phosphorylation of NF- κ B and I κ B proteins: implications in cancer and inflammation. *Trends Biochem Sci*, 30(1): 43–52
- Vieira H L, Alves P M, Vercelli A (2011). Modulation of neuronal stem cell differentiation by hypoxia and reactive oxygen species. *Prog Neurobiol*, 93(3): 444–455
- Vilas-Boas F, Henrique D (2010). HES6-1 and HES6-2 function through different mechanisms during neuronal differentiation. *PLoS ONE*, 5(12): e15459
- Villeda S A, Luo J, Mosher K I, Zou B, Britschgi M, Bieri G, Stan T M, Fainberg N, Ding Z, Egel A, Lucin K M, Czirr E, Park J S, Couillard-Després S, Aigner L, Li G, Peskind E R, Kaye J A, Quinn J F, Galasko D R, Xie X S, Rando T A, Wyss-Coray T (2011). The ageing systemic milieu negatively regulates neurogenesis and cognitive function. *Nature*, 477(7362): 90–94
- Wang D D, Bordey A (2008). The astrocyte odyssey. *Prog Neurobiol*, 86(4): 342–367
- Weih F, Durham S K, Barton D S, Sha W C, Baltimore D, Bravo R (1997). p50-NF- κ B complexes partially compensate for the absence of RelB: severely increased pathology in p50^{-/-}relB^{-/-} double-knockout mice. *J Exp Med*, 185(7): 1359–1370
- Westlake C J, Baye L M, Nachury M V, Wright K J, Ervin K E, Phu L, Chalouni C, Beck J S, Kirkpatrick D S, Slusarski D C, Sheffield V C, Scheller R H, Jackson P K (2011). Primary cilia membrane assembly is initiated by Rab11 and transport protein particle II (TRAPPII) complex-dependent trafficking of Rabin8 to the centrosome. *Proc Natl Acad Sci USA*, 108(7): 2759–2764
- Whitney N P, Eidem T M, Peng H, Huang Y, Zheng J C (2009). Inflammation mediates varying effects in neurogenesis: relevance to the pathogenesis of brain injury and neurodegenerative disorders. *J Neurochem*, 108(6): 1343–1359
- Widera D, Kaus A, Kaltschmidt C, Kaltschmidt B (2008). Neural stem cells, inflammation and NF- κ B: basic principle of maintenance and repair or origin of brain tumours? *J Cell Mol Med*, 12(2): 459–470
- Widera D, Mikenberg I, Elvers M, Kaltschmidt C, Kaltschmidt B (2006a). Tumor necrosis factor alpha triggers proliferation of adult neural stem cells via IKK/NF- κ B signaling. *BMC Neurosci*, 7(1): 64
- Widera D, Mikenberg I, Kaltschmidt B, Kaltschmidt C (2006b). Potential role of NF- κ B in adult neural stem cells: the underrated steersman? *Int J Dev Neurosci*, 24(2–3): 91–102
- Widera D, Mikenberg I, Kaus A, Kaltschmidt C, Kaltschmidt B (2006c). Nuclear factor- κ B controls the reaggregation of 3D neurosphere cultures *in vitro*. *Eur Cell Mater*, 11: 76–84, discussion 85
- Winner B, Kohl Z, Gage F H (2011). Neurodegenerative disease and adult neurogenesis. *Eur J Neurosci*, 33(6): 1139–1151
- Wong E T, Tergaonkar V (2009). Roles of NF- κ B in health and disease: mechanisms and therapeutic potential. *Clin Sci (Lond)*, 116(6): 451–465
- Wong G, Goldshmit Y, Turnley A M (2004). Interferon-gamma but not TNF alpha promotes neuronal differentiation and neurite outgrowth of murine adult neural stem cells. *Exp Neurol*, 187(1): 171–177
- Wooten M W, Seibenhener M L, Neidigh K B, Vandenplas M L (2000). Mapping of atypical protein kinase C within the nerve growth factor signaling cascade: relationship to differentiation and survival of PC12 cells. *Mol Cell Biol*, 20(13): 4494–4504
- Woronicz J D, Gao X, Cao Z, Rothe M, Goeddel D V (1997). I κ B kinase-beta: NF- κ B activation and complex formation with I κ B kinase-alpha and NIK. *Science*, 278(5339): 866–869
- Wu J P, Kuo J S, Liu Y L, Tzeng S F (2000). Tumor necrosis factor-alpha modulates the proliferation of neural progenitors in the subventricular/ventricular zone of adult rat brain. *Neurosci Lett*, 292(3): 203–206
- Xiao M, Inal C E, Parekh V I, Li X H, Whitnall M H (2009). Role of NF- κ B in hematopoietic niche function of osteoblasts after radiation injury. *Exp Hematol*, 37(1): 52–64
- Yaddanapudi K, De Miranda J, Hornig M, Lipkin W I (2011). Toll-like receptor 3 regulates neural stem cell proliferation by modulating the Sonic Hedgehog pathway. *PLoS ONE*, 6(10): e26766
- Yang C, Atkinson S P, Vilella F, Lloret M, Armstrong L, Mann D A, Lako M (2010). Opposing putative roles for canonical and noncanonical NF- κ B signaling on the survival, proliferation, and differentiation potential of human embryonic stem cells. *Stem Cells*, 28(11): 1970–1980
- Yang L, Tao L Y, Chen X P (2007). Roles of NF- κ B in central nervous system damage and repair. *Neurosci Bull*, 23(5): 307–313
- Yeo J E, Kang S K (2007). Selenium effectively inhibits ROS-mediated apoptotic neural precursor cell death *in vitro* and *in vivo* in traumatic brain injury. *Biochim Biophys Acta*, (11–12): 1199–1210
- Yirmiya R, Goshen I (2011). Immune modulation of learning, memory, neural plasticity and neurogenesis. *Brain Behav Immun*, 25(2): 181–213
- Yoneyama M, Kawada K, Gotoh Y, Shiba T, Ogita K (2010). Endogenous reactive oxygen species are essential for proliferation of neural stem/progenitor cells. *Neurochem Int*, 56(6–7): 740–746
- Young K M, Bartlett P F, Coulson E J (2006). Neural progenitor number is regulated by nuclear factor- κ B p65 and p50 subunit-dependent proliferation rather than cell survival. *J Neurosci Res*, 83(1): 39–49
- Zhang C, Wu H, Zhu X, Wang Y, Guo J (2011). Role of transcription factors in neurogenesis after cerebral ischemia. *Rev Neurosci*, 22(4): 457–465
- Zhang Q, Wang C, Liu Z, Liu X, Han C, Cao X, Li N (2012a). Notch signal suppresses Toll-like receptor-triggered inflammatory responses in macrophages by inhibiting extracellular signal-regulated kinase 1/2-mediated nuclear factor κ B activation. *J Biol Chem*, 287

- (9): 6208–6217
- Zhang Y, Liu J, Yao S, Li F, Xin L, Lai M, Bracchi-Ricard V, Xu H, Yen W, Meng W, Liu S, Yang L, Karmally S, Liu J, Zhu H, Gordon J, Khalili K, Srinivasan S, Bethea J R, Mo X, Hu W (2012b). Nuclear factor κ B signaling initiates early differentiation of neural stem cells. *Stem Cells*, 30(3): 510–524
- Zhao C, Deng W, Gage F H (2008). Mechanisms and functional implications of adult neurogenesis. *Cell*, 132(4): 645–660
- Zhao C, Xiu Y, Ashton J, Xing L, Morita Y, Jordan C T, Boyce B F (2012). Noncanonical NF- κ B signaling regulates hematopoietic stem cell self-renewal and microenvironment interactions. *Stem Cells*, 30(4): 709–718
- Zhao M, Li X D, Chen Z (2010). CC2D1A, a DM14 and C2 domain protein, activates NF- κ B through the canonical pathway. *J Biol Chem*, 285(32): 24372–24380
- Zhu C, Liu Z, Gui L, Yao W, Qian W, Zhang C (2008). Mutated I κ B α represses proliferation of immortalized neural progenitor cells and prevents their apoptosis after oxygen-glucose deprivation. *Brain Res*, 1244: 24–31
- Ziercher L, Filhol O, Laudet B, Prudent R, Cochet C, Buchou T (2011). Structure-function analysis of the beta regulatory subunit of protein kinase CK2 by targeting embryonic stem cell. *Mol Cell Biochem*, 356(1–2): 75–81
- Zong M, Satoh A, Yu M K, Siu K Y, Ng W Y, Chan H C, Tanner J A, Yu S (2012). TRAPPC9 mediates the interaction between p150 and COPII vesicles at the target membrane. *PLoS ONE*, 7(1): e29995
- Zong M, Wu X G, Chan C W, Choi M Y, Chan H C, Tanner J A, Yu S (2011). The adaptor function of TRAPPC2 in mammalian TRAPPs explains TRAPPC2-associated SEDT and TRAPPC9-associated congenital intellectual disability. *PLoS ONE*, 6(8): e23350