

Proteins moonlighting in tumor metabolism and epigenetics

Lei Lv (✉)¹, Qunying Lei (✉)^{2,3,4}

¹MOE Key Laboratory of Metabolism and Molecular Medicine, Department of Biochemistry and Molecular Biology, School of Basic Medical Sciences, Fudan University, Shanghai 200032, China; ²Fudan University Shanghai Cancer Center & Institutes of Biomedical Sciences; Cancer Institutes; Key Laboratory of Breast Cancer in Shanghai; The Shanghai Key Laboratory of Medical Epigenetics, Shanghai Medical College, Fudan University, Shanghai 200032, China; ³Department of Oncology, Shanghai Medical College, Fudan University, Shanghai 200032, China; ⁴State Key Laboratory of Medical Neurobiology, Fudan University, Shanghai 200032, China

© The Author(s) 2020. This article is published with open access at link.springer.com and journal.hep.com.cn

Abstract Cancer development is a complicated process controlled by the interplay of multiple signaling pathways and restrained by oxygen and nutrient accessibility in the tumor microenvironment. High plasticity in using diverse nutrients to adapt to metabolic stress is one of the hallmarks of cancer cells. To respond to nutrient stress and to meet the requirements for rapid cell proliferation, cancer cells reprogram metabolic pathways to take up more glucose and coordinate the production of energy and intermediates for biosynthesis. Such actions involve gene expression and activity regulation by the moonlighting function of oncoproteins and metabolic enzymes. The signal – moonlighting protein – metabolism axis facilitates the adaptation of tumor cells under varying environment conditions and can be therapeutically targeted for cancer treatment.

Keywords moonlighting function; tumor metabolism; epigenetics

Introduction

Proliferation of cancer cell is restrained by nutrient accessibility in the tumor microenvironment. Cancer cells have to reprogram metabolic pathways to adapt to nutrient stress. Metabolic reprogramming is a hallmark of cancer and plays a vital role in the tumorigenesis and maintenance of malignancy [1]. Tumor cells take up more glucose and convert the majority of them to lactate even in the presence of oxygen compared with normal cells [2]. This phenomenon was first observed during a study on the metabolic reprogramming in cancer and was uncovered by Otto Warburg, hence its designation as the Warburg effect. It is exploited to detect tumor clinically by the combined use of ¹⁸F-deoxyglucose and positron emission tomography (FDG-PET). Glutaminolysis, which involves glutamine utilization, is also upregulated in cancer cells. Aerobic glycolysis and glutaminolysis are two major metabolic alterations diverting carbon sources to biosynthesis to produce enough nucleotides, amino acids, and lipids for cell proliferation [3,4]. However, in non-glycolytic

cancers, such as prostate cancer and B cell lymphoma, exogenous free fatty acids (FFAs) are oxidized to produce energy [5,6] and to promote tumor growth and metastasis [7,8]. Intriguingly, emerging evidence demonstrated that adipocytes and lipid metabolism are correlated with the drug resistance of cancer [9]. Iwamoto and colleagues found that anti-angiogenic drugs trigger lipid-dependent metabolic reprogramming, leading to increased FFAs and drug resistance. This anti-angiogenic drug resistance can be overcome by the suppression of fatty acid oxidation [10]. Despite these fascinating observations, the underlying mechanism by which cancer cells reprogram metabolic pathways according to metabolic signals remains elusive. Results of many studies have demonstrated that glycolysis and other metabolic alterations observed in multiple cancer types are governed by the moonlighting function of oncoproteins and metabolic enzymes to promote anabolism and support cell growth and proliferation [11–13].

Recently, several metabolic enzymes that possess canonical and moonlighting functions have been discovered [14–16]. Moonlighting proteins perform several independent and often unrelated noncanonical functions. Many oncoproteins and metabolic enzymes are moonlighting in tumor metabolism. The canonical regulation of oncoproteins to tumor metabolism occurs as a secondary

Received May 12, 2020; accepted July 27, 2020

Correspondence: Lei Lv, lvlei@fudan.edu.cn;

Qunying Lei, qllei@fudan.edu.cn

response to nutrient stress and proliferation signals. The effector proteins need to be transcribed and translated before functioning. Moonlighting oncoproteins regulate metabolic flux as a primary response to metabolic stress and growth factor signaling associated with cell survival and proliferation, respectively, resulting in a more rapid and efficient response. Interestingly, most of the moonlighting proteins that regulate tumor metabolism are metabolic enzymes that directly sense metabolic signals and in turn activate gene expression through the regulation of transcription factor, DNA methylation, or histone acetylation. Here, we review proteins moonlighting in tumor metabolism and discuss their biological functions, translational implications, and research prospects.

Oncoproteins and tumor suppressor moonlighting in tumor metabolism

KRAS

The *RAS* oncogene family comprises more than 150 distinct members [17], and its signaling is frequently deregulated in tumorigenesis [18]. *RAS* proteins function as a molecular switch that cycles between active GTP-bound and inactive GDP-bound conformations [19]. *RAS* signaling controls many different cellular processes, such as cell proliferation, differentiation, and apoptosis [17]. Notably, multiple and recurrent gain-of-function mutations in *RAS* genes were identified in various types of human cancer, indicating that *RAS* signaling plays a fundamental role in tumor development [17,18]. Interestingly, not all

RAS mutations occur at a similar frequency. The most commonly mutated *RAS* genes are *KRAS* (85%), *NRAS* (12%), and *HRAS* (3%) [19]. *KRAS* gene encodes two different protein forms (*KRAS4A* and *KRAS4B*) through alternative splicing [20]. Mutation of *KRAS*, mostly at G12, abolishes its GTPase activity and constitutively activates downstream signaling, leading to unconstrained cell proliferation, tumorigenesis, and tumor drug resistance [21].

Recent studies have found that oncogenic *RAS* promotes metabolic reprogramming of tumor cells to provide biomass support for uncontrolled proliferation by transcriptional upregulation of the glucose transporters and glycolytic enzymes, which increases glucose uptake and glycolytic flux even in the presence of oxygen; this is known as the Warburg effect [18,22,23]. Besides its canonical role in transmitting signals from extracellular growth factors to the cell nucleus, *KRAS4A* but not *KRAS4B* directly binds to and regulates hexokinase 1 (*HK1*), which is the initiating enzyme of glycolysis that catalyzes the transfer of a high energy phosphate group from ATP to glucose and produces glucose-6-phosphate in a GTP- and prenylation-dependent manner. Binding with *HK1* on the outer mitochondrial membrane, *KRAS4A* represses the allosteric inhibition of *HK1*, thereby increasing its activity and enhancing glycolytic flux. Oncogenic *KRAS4A* has twice the effect of *KRAS4B* on the enhancement of the glucose consumption because of the unique function of *KRAS4A* on *HK1* regulation [24], thereby providing a therapeutic target for patients bearing the *KRAS* oncogenic mutation (Fig. 1).




Fig. 1 The canonical and moonlighting functions of KRAS. The canonical (left panel) and non-canonical (right panel) functions of KRAS are summarized. Abbreviations: EGFR, epidermal growth factor receptor; GTP, guanosine triphosphate; GDP, guanosine diphosphate; PI3K, phosphatidylinositol 3-kinase; HK1, hexokinase 1.

p53

Tumor suppressor p53 serves diverse functions in multiple physiological and pathological processes, including cell cycle arrest, DNA repair, apoptosis, and senescence [25]. Besides its role in the nucleus as a transcription factor, p53 retains several moonlighting functions in the cytoplasm, especially in metabolic reprogramming. IKK-NF-κB is a canonical signal pathway in the regulation of cancer metabolism. p53 restricts glycolysis and the expression of glucose transporter GLUT3 by suppressing the activation of IKK-NF-κB pathway to inhibit cell transformation [26]. As the energy factory of cell, mitochondria plays a crucial role in cell metabolism. Cytosolic p53 promotes the permeability of mitochondria and triggers apoptosis through the activation of the pro-apoptotic factor Bax independent of its transcription activity [27]. Mdm2-mediated monoubiquitylation of p53 induces its mitochondrial translocation [28]. Being the guardian of mitochondrial genome, p53 interacts with CHCHD4 and POLG to enhance the repair of oxidative mtDNA damage and to maintain mtDNA integrity [29]. Glucose-6-phosphate dehydrogenase (G6PD) is the first and rate-limiting enzyme of the pentose phosphate pathway (PPP). p53 binds to G6PD and suppresses the formation of active dimer, thereby repressing glucose consumption, NADPH production, and biosynthesis. Tumor-associated p53 mutation abolishes its activity to inhibit G6PD, thereby enhancing PPP flux, increasing glucose consumption, and directing more glucose to biosynthesis to support the rapid growth and proliferation of tumor cells [30] (Fig. 2).

Signal transducers and activators of transcription 3 (STAT3)

STAT3 plays a key role in multiple biological processes, including cell growth, differentiation inflammation, and

cancer development, depending on its function as a transcription factor [31–34]. Mechanistically, JAK kinases activate STAT3 via phosphorylation at Y705 residue upon cytokine or growth factor stimulation. Phosphorylated STAT3 forms homodimers, translocates into the nucleus, and activates target gene expression by binding to their promoter [35,36]. Besides serving as a transcription factor, a novel function of STAT3 in the mitochondria has also been discovered. Joanna Wegrzyn and colleagues found that STAT3 can get into the mitochondria and promote the activities of complexes I and II of the electron transport chain (ETC) and oxygen consumption, indicating the important role of STAT3 in regulating cell metabolism and maintaining cellular homeostasis independent of its transcriptional activity [37]. Subsequent studies further revealed the function of mitochondrial STAT3. STAT3 optimizes the function of ETC and regulates the opening of the mitochondrial permeability transition pore (mPTP), thereby manipulating the production of ATP and reactive oxygen species (ROS) and cell survival. Moreover, mitochondria STAT3 facilitates oncogenic transformation by increasing the ETC activity and sustaining altered glucose metabolism in cells [38] (Fig. 3). Therefore, targeting mitochondrial STAT3 function may be a promising treatment for cancer [38]. Recently, an inhibitor was developed to target mitochondrial STAT3 (mSTAT3). The binding of mSTAT3 with the inhibitor causes mitochondrial dysfunction and tumor death. Interestingly, the lethal consequence induced by mSTAT3 inhibition is enhanced by glucose starvation, which demonstrated the increase in sensitivity of cancer cells under metabolic stress to the mSTAT3 inhibitor; thus, mSTAT3 is a potential target for cancer therapy [39].

c-Myc

The proto-oncogene *c-Myc* is associated with many




Fig. 2 The canonical and moonlighting functions of p53. The canonical and non-canonical functions of p53 are summarized. Abbreviations: G6P, glucose-6-phosphate; G6PD, glucose-6-phosphate dehydrogenase; Bax, Bcl-2 associated X protein; Ub, ubiquitin.




Fig. 3 The canonical and moonlighting functions of STAT3. The canonical and non-canonical functions of STAT3 are summarized. Abbreviations: STAT3, signal transducers and activators of transcription 3; JAK, Janus kinase.

physiological processes, such as cell cycle progression, apoptosis, and cell transformation [40]. *c-Myc* dysregulation occurs in more than 70% of tumors; it affects gene and microRNA expressions, genomic amplification, and the overall organization of nucleus [41]. The *c-Myc* protein forms a complex with MAX in the nucleus and promotes the transcription of target genes [42]. Overexpression of *c-Myc* causes genomic instability by increasing the phosphorylation at S139 of histone H2AX and forming γ H2AX [43]. *c-Myc* can also reprogram metabolism in tumor cells, which induces metabolic stress and activates AMP-activated protein kinase (AMPK) [44–46]. *c-Myc* transcriptionally upregulates expression levels of glucose transporter 1 (GLUT1) and hexokinase 2 (HK2), as well as pyruvate kinase 2 (PKM2) through mRNA splicing, thereby increasing glycolytic flux and promoting tumor progression [47,48]. Besides its function in the nucleus, *c-Myc* cooperates with MCL1 to increase mitochondrial oxidative phosphorylation (mtOXPHOS), ROS production, and HIF-1 α expression to contribute to the chemotherapy resistance of cancer stem cells (CSCs) in triple negative breast cancer. The suppression of HIF-1 α restores chemotherapy sensitivity and inhibits CSC expansion [49].

Metabolic enzymes moonlighting in tumor metabolism

Aldolase (ALDO)

Aldolase (ALDO) is positioned midway in the glycolytic pathway and catalyzes the reversible cleavage of fructose-1,6-bisphosphate (FBP) to dihydroxyacetone-3-phosphate (DHAP) and glyceraldehyde-3-phosphate (G3P) [50]. Three aldolase family members are involved in metabo-

lism and glycolysis, namely, ALDOA, ALDOB, and ALDOC. Aldolase reportedly also plays a role in fructolysis. Fructose is catalyzed to fructose-1-phosphate by ketohexokinase and further converted to DHAP and glyceraldehyde by ALDOB and ALDOC [51]. ALDOA plays a major role in glycolysis and is highly expressed in several tumor types [52,53].

Besides its function as a glycolytic enzyme, recent studies demonstrated that aldolase can sense glucose level by binding to its substrate FBP. FBP-unoccupied aldolase binds to and suppresses ER-localized transient receptor potential V (TRPV) channels, thereby decreasing calcium level at the ER-lysosome contact, which in turn promotes the interaction of the channel proteins with lysosomal v-ATPase. Thus, the formation of a lysosomal complex containing v-ATPase, regulator, axin, liver kinase B1 (LKB1), and AMPK to induce AMPK activation is facilitated [14,54]. Once activated under low energy status, AMPK promotes ATP production by increasing the activity or expression level of proteins involved in catabolism and saves ATP by shutting down biosynthetic pathways, thereby contributing to the adaptation of cells to the nutrient limitation and the maintenance of energy homeostasis [55] (Fig. 4).

Glyceraldehyde-3-phosphate dehydrogenase (GAPDH)

GAPDH is a conventional enzyme in glycolysis; it converts glyceraldehyde-3-phosphate to 1,3-bisphosphoglycerate and generates NADH [56]. GAPDH is reportedly involved in synthetic lethality in KRAS or BRAF mutant colorectal cancer. High dose vitamin C causes GAPDH S-glutathionylation and inactivation, leading to cell death, thereby suggesting that GAPDH may be a therapeutic target for colorectal cancer with KRAS or BRAF mutations [57]. GAPDH can bind to the AU-rich elements (AREs) in




Fig. 4 The canonical and moonlighting functions of ALDO. The canonical and non-canonical functions of ALDO are summarized. Abbreviations: F6P, fructose-6-phosphate; FBP, fructose 1,6-bisphosphate; DHAP, dihydroxyacetone-3-phosphate; G3P, glyceraldehyde-3-phosphate; ER, endoplasmic reticulum; ALDO, aldolase; LKB1, liver kinase B1; AMPK, AMP-activated protein kinase.

the 3' UTR and destabilize the mRNA of vasoconstrictor endothelin-1 (ET-1). Interestingly, this process is also regulated by oxidative stress-induced S-glutathionylation of GAPDH. S-glutathionylated GAPDH cannot modify mRNA unwinding, thereby enhancing the mRNA stability of ET-1 [58].

In addition to its role in the cytoplasm, cytosolic GAPDH reportedly translocates into the nucleus in response to stresses and regulates physiological processes. Nitric oxide-induced S-nitrosylation at C150 induces GAPDH binding to Siah1 and initiates apoptosis [59]. Nuclear S-nitrosylated GAPDH transnitrosylates other nuclear proteins, such as SIRT1 and HDAC2, thereby suggesting that nitric oxide group transfer may represent a new signaling transmission mode [60]. The p300/CREB binding protein (CBP) acetylates nuclear GAPDH at K160, which in turn activates CBP and its downstream target *p53* to facilitate cell apoptosis [61]. During apoptosis, nuclear GAPDH cooperates with autophagy to protect caspase-independent cell death [62]. Under glucose starvation, GAPDH is phosphorylated at S122 by AMPK, leading to GAPDH nuclear translocation. Nuclear GAPDH stimulates SIRT1 activation and finally promotes autophagy to increase cell survival [63]. OCA-S is critical for S phase-dependent histone H2B transcription and may play a key role in cell metabolism. Nuclear GAPDH is a key component of OCA-S, whose function is stimulated by NAD^+ but repressed by NADH [64]. Apart from localization in cytoplasm and nucleus, free GAPDH protein transfers to the plasma membrane and forms a complex with transferrin, thereby becoming involved in iron metabolism [65] (Fig. 5).

Phosphoglycerate kinase 1 (PGK1)

PGK1 is a glycolytic enzyme that catalyzes the conversion of 1,3-bisphosphoglycerate (1,3-BPG) to 3-phosphoglycerate (3-PG) and produces ATP [66]. The activity and function of PGK1 are regulated by posttranslational modifications. Acetylation of PGK1 at K323 is reversibly regulated by P300/CBP associated factor (PCAF) and sirtuin 7 (SIRT7) and promotes the enzymatic activity and tumor metabolism of PGK1, thereby supporting tumor growth [67]. O-GlcNAcylation of PGK1 at T255 promotes PGK1 activity, lactate production, and mitochondria translocation, where PGK1 reduces oxidative phosphorylation by inhibiting pyruvate dehydrogenase (PDH), thereby promoting tumor growth [68]. Besides O-GlcNAcylation, phosphorylation of PGK1 also regulates its mitochondria localization. ERK phosphorylates PGK1 at S203 under oncogenic signal stimulation, which promotes its mitochondria translocation through PIN1-mediated cis-trans isomerization. Intriguingly, PGK1 acts as a protein kinase in the mitochondria and phosphorylates pyruvate dehydrogenase kinase 1 (PDK1) at T338, resulting in PDHK1 activation and phosphorylation and inactivation of PDH. These actions result in the blocking of pyruvate utilization in mitochondrial and ROS production and increase lactate production, thereby promoting Warburg effect to meet the needs of rapid proliferation of tumor cells by coordinating glycolysis and tricarboxylic acid (TCA) cycle [69]. Interestingly, PGK1 can also autophosphorylate and activate itself at Y324, thereby promoting glycolysis. Phosphatase PTEN dephosphorylates PGK1, thus suppressing glycolysis, ATP production, and tumor growth [70].




Fig. 5 The canonical and moonlighting functions of GAPDH. The canonical and non-canonical functions of GAPDH are summarized. Abbreviations: G6P, glucose-6-phosphate; 1,3-BPG, 1,3-bisphosphoglycerate; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; HDAC2, histone deacetylase 2; SIRT1, sirtuin 1; SNO, S-nitrosylation; SSG, S-glutathionylation.

S203 phosphorylation and T255 O-GlcNAcylation both mediate PGK1 mitochondrial localization, whereas all the modifications, including K323 acetylation, T255 O-GlcNAcylation, and Y324 phosphorylation, activate its enzymatic activity, thereby raising the question on whether these posttranslational modifications have crosstalk. S203 phosphorylation and T255 O-GlcNAcylation reportedly promote PGK1 mitochondrial translocation independently [68], but whether K323 acetylation, T255 O-GlcNAcylation, and Y324 phosphorylation have crosstalk is unclear. Possibly, these modifications enhance PGK1 activity collaboratively or they respond to different signals from the tumor microenvironment. These speculations require further investigation. In addition, under glutamine deprivation and hypoxia, PGK1 phosphorylates Beclin1 at S30, thereby enhancing VPS34-Beclin1-ATG14L complex activity, triggering autophagy, providing biomacromolecules for both energy production and biosynthesis, and maintaining the metabolism and homeostasis of tumor cells [71,72]. PGK1 reportedly translocates into the nucleus and functions as a transcription factor and driver of cell metastasis by repressing E-cadherin expression and metabolic reprogramming, thereby demonstrating that the subcellular localization of PGK1 creates a balance between proliferation and metastasis in SMAD4-negative pancreatic ductal adenocarcinoma [73] (Fig. 6).

Pyruvate kinase M2 (PKM2)

Pyruvate kinase regulates the final rate-limiting step of

glycolysis, thereby catalyzing the conversion of phosphoenolpyruvate (PEP) to pyruvate and producing ATP. Pyruvate kinase has four isoforms, namely, PKL, PKR, PKM1, and PKM2. PKL, PKR, and PKM1 are expressed in specific tissues; the expression of PKM2 declines in a number of adult tissues but is highly upregulated in developing embryos and tumors [74–76]. The cytoplasmic function and regulation of PKM2 is crucial for tumor growth [77,78]. Tumor cells take up more glucose than normal cells. Under the stimulation of glucose at a high level, PKM2 is acetylated by PCAF at K305 and recruited by chaperone HSC70 to lysosome for degradation, thereby accumulating glycolytic intermediates for biosynthesis to support rapid tumor cell proliferation [79,80]. Among these isoforms, only PKM2 is detected in the nucleus, as it contains a nuclear localization signal (NLS), which is buried when it forms a tetramer and functions as metabolic kinase in the cytoplasm. PKM2 is acetylated by p300 acetyltransferase at K433, which is unique for PKM2, upon mitogenic and oncogenic signal stimulation, thereby preventing the binding of its allosteric activator, fructose-1,6-bisphosphate (FBP), and promoting tetramer-dimer transition, exposure of NLS, and translocation into the nucleus [81]. Phosphorylation also regulates the cytoplasm–nucleus translocation of PKM2. ERK2 binds to and phosphorylates PKM2 at S37 under EGF stimulation, which causes a conformational change, promotes PKM2 binding to importin $\alpha 5$ and translocation into the nucleus, and acts as a protein kinase [82,83]. PKM2 directly binds to and phosphorylates histone H3 at T11




Fig. 6 The canonical and moonlighting functions of PGK1. The canonical and non-canonical functions of PGK1 are summarized. Abbreviations: 1,3-BPG, 1,3-bisphosphoglycerate; 3-PG, 3-phosphoglycerate (3-PG); ADP, adenosine diphosphate; ATP, adenosine triphosphate; PGK1, phosphoglycerate kinase 1; SIRT7, sirtuin 1; PDK1, pyruvate dehydrogenase kinase 1; PDH, pyruvate dehydrogenase; OXPHOS, oxidative phosphorylation; O-GlcNAc, O-GlcNAcylation.

using PEP as the phosphate donor, thereby enhancing the Warburg effect by upregulating the expressions of c-Myc target genes, such as glucose transporter type 1 (GLUT1) and lactate dehydrogenase A (LDHA) [82,84,85]. Moreover, nuclear PKM2 in the form of a dimer can also phosphorylate STAT3 at Y705 in the nucleus, promoting its transcriptional activity and activating MEK5 expression to promote cell proliferation and tumor growth [86,87]. Intriguingly, SAICAR, a metabolite abundant in proliferating cells, can bind to PKM2 and induce its protein kinase activity [88]. Besides its function as a protein kinase, nuclear PKM2 also serves as a co-transcription factor of HIF-1 α and activates downstream target gene expression to reprogram tumor metabolism and promote tumor angiogenesis [89–92] (Fig. 7). Vander Heiden and others have not found PKM2 kinase activity under basal conditions [93]; their study is similar to ours, because we can only detect its protein kinase activity by stimulating with EGF [81]. Thus, more metabolic enzymes could potentially function as protein kinases under varied appropriate stimuli.

The dual function of PKM2 as a metabolic kinase and as a protein kinase is vital for tumor development. Thus, PKM2 inhibitors and activators are developed to treat tumors by distinct mechanisms [94–99]. PKM2 inhibitors promote tumor cell death by decreasing glycolytic flux and energy supply, whereas PKM2 activators suppress tumor cell proliferation by diverting glycolytic intermediates away from biosynthesis and inhibiting the nuclear function.

Fructose-1,6-bisphosphatase 1 (FBP1)

FBP1 is one of the rate-limiting enzymes in gluconeogenesis and converts fructose-1,6-bisphosphate to fructose-6-phosphate. However, many studies have demonstrated that FBP1 moonlights in tumor metabolism and functions as a tumor suppressor by decreasing glucose uptake, reducing glycolysis, and limiting cancer cell proliferation [100–103]; all these actions depend on its function in the nucleus. Nuclear FBP1 binds to and represses the activity of both HIF-1 α and HIF-2 α independent of its enzymatic activity, thereby decreasing the expressions of HIF target genes, such as *VEGF*, *GLUT1*, and *LDHA*, and suppressing tumor metabolism. Notably, nucleus-excluded FBP1 failed to inhibit tumor cell growth, indicating the importance of FBP1 moonlighting function [104]. In addition, FBP1 can bind to Notch1 and facilitates its proteasomal degradation, thereby reducing the expressions of Notch1 target genes and inhibiting breast tumorigenesis [105] (Fig. 8). FBP1 inhibits tumor metabolism and is a tumor suppressor. However, this is not always the case. In the lung tumor microenvironment, the high expression level of FBP1 in NK cells decreases glycolysis and viability, leading to the dysfunction of NK cells, which can be reversed by FBP1 inhibition [106].

Fructose-1,6-bisphosphatase 2 (FBP2)

FBP2 is the isozyme of FBP1 that shares 76.6% sequence




Fig. 7 The canonical and moonlighting functions of PKM2. The canonical and non-canonical functions of PKM2 are summarized. Abbreviations: PEP, phosphoenolpyruvate; PKM2, pyruvate kinase M2; ERK2, extracellular signal-regulated kinase 2; HSC70, heat shock cognate 71 kDa protein; H3, histone H3; STAT3, signal transducers and activators of transcription 3; HIF-1 α , hypoxia-inducible factor-1 α .

identity with that in vertebrates. FBP1 is primarily expressed in the liver and kidney, whereas FBP2 is expressed more ubiquitously. Like FBP1, FBP2 suppresses a tumor and inhibits sarcoma progression independent of its function in gluconeogenesis, and these two actions involve two different mechanisms. FBP2 restrains

enhanced glycolysis in cytoplasm, thereby inhibiting the Warburg effect and cell proliferation. Moreover, FBP2 translocates into the nucleus and binds to and suppresses c-Myc mediated *TFAM* expression, which in turn represses mitochondrial biogenesis and respiration independent of its enzymatic activity [15] (Fig. 8).




Fig. 8 The canonical and moonlighting functions of FBP1/2. The canonical and non-canonical functions of FBP1/2 are summarized. Abbreviations: F-6-P, fructose-6-phosphate; F-1,6-P, fructose-1,6-bisphosphate; FBP1/2, fructose-1,6-bisphosphatase 1/2; HIF, hypoxia-inducible factor; VEGF, vascular endothelial growth factor; GLUT1, glucose transporter 1; LDHA, lactate dehydrogenase A; TFAM, transcription factor A, mitochondrial.

Phosphoenolpyruvate carboxykinase 1 (PCK1)

PCK1 is one of the rate-limiting enzymes of gluconeogenesis; it converts oxaloacetate and GTP to phosphoenolpyruvate and CO₂ [107]. PCK1 and its isoform, PCK2, shares 63.4% sequence identity and localizes in cytoplasm and mitochondria, respectively [108]. Rapid proliferative tumor cells require a large amount of amino acids, nucleosides, and fatty acids to synthesize proteins, DNA, and lipids. PCK1 reportedly shows protein kinase activity and promotes lipogenesis. Specifically, AKT phosphorylates PCK1 at S90 and promotes its translocation to the endoplasmic reticulum. In the endoplasmic reticulum, it phosphorylates INSIG1 at S207 and INSIG2 at S151, respectively, using GTP as the phosphate donor. This leads to the activation of SREBP proteins and the expression of genes required for lipogenesis. Thus, the proliferation of hepatocellular carcinoma (HCC) cells and tumorigenesis are promoted, demonstrating that the protein kinase activity of PCK1 is a promising target for the treatment of HCC [16].

Glutaminase (GLS) 1

Increased uptake and metabolism of glutamine is a hallmark of cancer. Glutamine is rapidly consumed to generate building blocks (amino acids, nucleotides, lipids, and carbohydrates) and energy (ATP) to support the growth and proliferation of cancer cells [1]. Glutamine also functions as a cell signaling molecule and promotes the synthesis of the antioxidant GSH [109]. GLS is the enzyme responsible for catalyzing the conversion of glutamine to glutamate, which is the first step of glutaminolysis. GLS has two isozymes, namely, GLS1 and GLS2. GLS1 is broadly expressed in various tissues and is highly expressed in many types of cancer. In contrast, the expression of GLS2 is restricted to the liver and brain [109].

Tumor development is challenged by nutrition limitation. Under glutamine deprivation, cancer cells undergo mitochondria fusion to promote the efficacy of respiratory chain and sustain the progression of tumor [110,111]. Cai *et al.* recently found that GLS1 can sense glutamine availability and initiates mitochondria fusion to help the cells overcome an energy crisis in an enzymatic activity-independent manner [112]. However, the detailed mechanism underlying this process has not yet been discovered.

Glutamate dehydrogenase (GDH) 1

GDH catalyzes glutamate to α -ketoglutarate (α -KG) and ammonia. α -KG is the key intermediate of the TCA cycle that interconnects amino acid and carbohydrate metabolisms [113]. There are two GDH isoforms in mammals,

namely, GDH1 and GDH2. The expression of GDH1 is upregulated in multiple human cancers, including glioma, breast cancer, lung cancer, and leukemia [114,115], suggesting its function in tumor development. GDH1 manipulates intracellular fumarate level by controlling α -KG production. Fumarate in turn binds to and activates glutathione peroxidase 1 (GPx1) to scavenge ROS, thereby maintaining redox homeostasis and promoting tumor growth. Blocking GDH1 by a small molecule R162 leads to the imbalance in redox homeostasis and to the inhibition of cancer cell proliferation and tumor growth [116]. As the product of GDH1, α -KG plays a variety of roles in different metabolic and cellular pathways [117,118]. The loss of LKB1 in lung cancer is correlated to the increase in metastasis and poor prognosis. In LKB1-deficient lung cancer, α -KG activates CamKK2-AMPK signaling by increasing their binding, resulting in energy production, anoikis resistance, and tumor metastasis; thus, inhibiting GDH1 with R162 impairs tumor metastasis [119]. Under low glucose stimulation, AMPK phosphorylates GDH1 at S384, promoting its interaction with RelA and IKK β . Thus, α -KG produced by GDH1 directly binds to and activates nuclear factor κ B (NF- κ B) signaling and upregulates GLUT1 expression, increasing glucose uptake and tumor cell survival [120,121] (Fig. 9). Collectively, GDH1 has unconventional functions in metabolic reprogramming through its product α -KG.

Ketohexokinase isoform A (KHK-A)

Ketohexokinase (KHK), also known as fructokinase, is the first rate-limiting enzyme in fructose metabolism. It catalyzes the conversion of fructose and ATP to fructose-1-phosphate (F-1-P) and ADP. F-1-P is then converted to dihydroxyacetone phosphate and glyceraldehyde-3-phosphate, which enter the later stage of glycolysis. KHK gene encodes two isoforms by alternative splicing, namely, KHK-A and KHK-C [122]. KHK-C is primarily expressed in liver, kidney, and intestines, whereas KHK-A is highly expressed in HCC cells and has a lower binding affinity to its substrate, fructose, than KHK-C; thus, KHK-A has a much lower activity than KHK-C [123,124].

c-Myc promotes the switch from the expression of KHK-C to KHK-A, resulting in much lower fructose catabolism rates, ATP consumption, and ROS production in HCC cells than in normal hepatocytes [125]. KHK-A interacts with and phosphorylates phosphoribosyl pyrophosphate synthetase 1 (PRPS1) at T225 in HCC cells, which activates PRPS1 by blocking the binding of its allosteric inhibitor ADP, leading to elevated *de novo* nucleic acid synthesis and HCC tumorigenesis through pentose phosphate pathway [125]. KHK-A is phosphorylated at S80, which is unique to KHK-A, by AMPK under oxidative stress. KHK-A in turn phosphorylates p62 at S28




Fig. 9 The canonical and moonlighting functions of GDH. The canonical and non-canonical functions of GDH are summarized. Abbreviations: ROS, reactive oxygen species; GPx1, glutathione peroxidase 1; Gln, glutamine; Glu, glutamate; α -KG, α -ketoglutarate; GLS, glutaminase; GDH, glutamate dehydrogenase; AMPK, AMP-activated protein kinase; CamKK2, calcium/calmodulin dependent protein kinase kinase 2; NF- κ B, nuclear factor κ B; GLUT1, glucose transporter 1.

and prevents its ubiquitylation, thereby promoting gene expression via Nrf2 activation to reduce ROS and to promote cell survival and HCC development [126].

Metabolic enzymes moonlighting in regulation of epigenetics

Lactate dehydrogenase A (LDHA)

The Warburg effect is characterized by increased glucose uptake and lactate production, which is produced by LDHA by catalyzing pyruvate, and plays a vital role in the regulation of metabolic pathways [127]. LDHA is acetylated at K5, thereby decreasing its enzymatic activity and promoting its degradation via chaperone mediated autophagy (CMA), which reduces tumor cell proliferation and migration [128]. Reduction of LDHA activity affects glycolytic flux and stimulates oxidative phosphorylation, decreases mitochondrial membrane potential and inhibits proliferation of tumor cells under hypoxia [129]. LDHA uses pyruvate as substrate but also catalyzes α -ketobutyrate (α -KB) [130]. However, the product and significance of this reaction is unknown. The collaboration study between the Canhua Huang and Qunying Lei groups found that LDHA translocates into the nucleus in response to HPV16E7-induced ROS accumulation, where it gains a moonlighting function to convert α -KB to α -hydroxybutyrate (α -HB), thereby promoting histone H3K79 trimethylation and activating antioxidant gene expression and Wnt signaling pathway to maintain cellular redox balance and cervical cancer cell proliferation under oxidative stress [131]. Besides α -HB, LDHA and MDH convert α -KG to L-2-hydroxyglutarate (L-2-HG) at acidic pH, thereby stabilizing HIF-1 α [132] (Figs. 10 and 11).

Isocitrate dehydrogenase 1/2 (IDH1/2)

Isocitrate dehydrogenase (IDH) catalyzes the oxidative decarboxylation of isocitrate to α -KG and produces NADPH [133]. IDH contains several isoforms, among which IDH1 and IDH2 are the most widely investigated and expressed in cytoplasm and mitochondria, respectively. Notably, IDH1 and IDH2 are frequently mutated in multiple tumors, including malignant gliomas [134,135], acute myeloid leukemia [136], cholangiocarcinoma [137], chondrosarcomas, and thyroid cancers [138]. Intriguingly, mutant IDH1/2 has gained the new function of converting α -KG to 2-HG [139–141], which accumulates in the tumor cells and inhibits multiple α -KG-dependent dioxygenases as a competitive inhibitor of α -KG, thereby leading to the dysregulation of histone and DNA demethylation to block cell differentiation and to promote tumorigenesis [142,143]. Elevated 2-HG level is observed in the serum of patients with IDH mutated glioma and acute myeloid leukemia, whereas the concentration of 2-HG is very low in normal tissues. 2-HG can be used as a biomarker for the clinical detection of cancer. mTOR is a key regulator of cancer metabolism. EGFR, NF1, and PTEN are upstream regulators of mTOR signaling pathway and are also frequently mutated in cancer. Interestingly, in the absence of EGFR/NF1/PTEN mutation, mTORC1/2 signaling is activated by IDH1/2 mutation and exogenous 2-HG, suggesting that 2-HG may regulate the activity of mTOR1/2 through an independent and non-canonical pathway to promote tumor cell survival and proliferation [144,145]. Besides histone and DNA demethylase, more than 60 other α -KG dependent dioxygenases reportedly exist, all of which could be inhibited by 2-HG; these include hypoxia inducible factors (HIF) and AlkB proteins that are responsible for the DNA repair of alkylation damage [146–148]. Thus, the oncometabolite 2-HG may




Fig. 10 The canonical and moonlighting functions of LDHA. The canonical and non-canonical functions of LDHA are summarized. Abbreviations: G6P, glucose-6-phosphate; LDHA, lactate dehydrogenase A; CMA, chaperone-mediated autophagy; ROS, reactive oxygen species; α-KG, α-ketoglutarate; α-KB, α-ketobutyrate; L-2-HG, L-2-hydroxyglutarate; HIF-1α, hypoxia-inducible factor-1α; O₂, oxygen; OH, hydroxylation; VHL, von Hippel-Lindau.

promote tumor development through multiple pathways, making mutant IDH a potential target for the reversal or relief of the biological consequences caused by 2-HG (Fig. 11).




Fig. 11 Metabolic enzymes moonlighting in regulation of epigenetics. The metabolic enzymes regulating epigenetics are summarized. Mutations in IDH, SDH, and FH accumulate 2-HG, succinate, and fumarate, respectively, thereby suppressing DNA and histone demethylation. The α-KB and Ac-CoA produced by LDHA and ACLY, respectively, promote histone H3K79 tri-methylation and acetylation. LDHA and MDH can also convert α-KG to L-2-HG at acidic pH. Abbreviations: G6P, glucose-6-phosphate; LDHA, lactate dehydrogenase A; α-KG, α-ketoglutarate; α-KB, α-ketobutyrate; L-2-HG, L-2-hydroxyglutarate; HIF-1α, hypoxia-inducible factor-1α; O₂, oxygen; OH, hydroxylation; VHL, von Hippel-Lindau; ACLY, ATP-citrate lyase; MDH, malate dehydrogenase; IDH, isocitrate dehydrogenase; SDH, succinate dehydrogenase; FH, fumarate hydratase; O-GlcNAc, O-GlcNAcylation; ATF2, activating transcription factor 2; ROS, reactive oxygen species; Me, methylation; H3K79me3, H3K79 tri-methylation; Ac, acetylation.

Table 1 Summary of metabolic enzymes moonlighting in tumor metabolism and epigenetics

Metabolic process	Enzyme	Canonical function		Moonlight function in tumor metabolism and epigenetics	
		Location	Function	Location	Function
Glycolysis	ALDO	Cytoplasm	Converts fructose-1,6-bisphosphate to dihydroxyacetone-3-phosphate and glyceraldehyde-3-phosphate	ER surface	Senses glucose availability through FBP binding and activates AMPK signaling under low glucose status [14,54]
	Cell membrane	Binds to OCA-S and increases histone H2B transcription, thereby affecting cell metabolism [64]			
		PGK1	Cytoplasm	Converts 1,3-bisphosphoglycerate to 3-phosphoglycerate and produces ATP	Mitochondria
	Cytoplasm				
		Nucleus	Phosphorylates Beclin1 at S30 and triggers autophagy under glutamine deprivation and hypoxia [71,72]		
			Functions as a transcription factor, drives cell metastasis via repression of E-cadherin expression and metabolic reprogramming [73]		
	PKM2	Cytoplasm	Converts phosphoenolpyruvate to pyruvate and produces ATP	Nucleus	Phosphorylates histone H3, upregulates c-Myc target genes, such as GLUT1 and LDHA, to promote the Warburg effect [82,84,85]
					Phosphorylates STAT3 at Y705, promotes its transcriptional activity, and activates MEK5 expression to promote cell proliferation and tumor growth [86,87]
	LDHA	Cytoplasm	Converts pyruvate to lactate	Nucleus	Functions as a co-transcription factor of HIF-1 α and activates downstream target gene expression to reprogram tumor metabolism [89–92]
Cytoplasm					Converts α -ketobutyrate to α -hydroxybutyrate, which promotes histone H3K79 tri-methylation and activates antioxidant gene expression and Wnt signaling pathway to maintain cellular redox balance and cervical cancer cell proliferation [130,131]
	Gluconeogenesis	FBP1/2	Cytoplasm	Converts fructose-1,6-bisphosphate to fructose-6-phosphate	Nucleus
PCK1					
	FBP1 binds to Notch1 and facilitates its degradation, thereby reducing the expressions of Notch1 target genes and inhibits breast tumorigenesis [105]				
PCK1	Cytoplasm	Converts oxaloacetate and GTP to phosphoenolpyruvate and CO ₂	ER	FBP2 binds to and suppresses c-Myc-mediated TFAM expression, which in turn represses mitochondrial biogenesis and respiration [15]	
				AKT-mediated PCK1 S190 phosphorylation functions as a protein kinase to phosphorylate INSIG1, leading to the activation of SREBP proteins and the expression of genes required for lipogenesis [16]	

(Continued)

Metabolic process	Enzyme	Canonical function		Moonlight function in tumor metabolism and epigenetics	
		Location	Function	Location	Function
TCA cycle	IDH1	Cytoplasm	Converts isocitrate to α -KG and produce NADPH (wild-type) Converts α -KG to 2-hydroxyglutarate (2-HG) (mutant)	Cytoplasm	2-HG accumulates in tumor cells, leading to the dysregulation of histone and DNA demethylation to block cell differentiation and promotes tumorigenesis, thereby activating mTORC1/2 signaling in the absence of EGFR/NF1/PTEN mutation [144,145] Accumulated succinate competitively inhibits the α -KG-dependent dioxygenases, including histone demethylases and TET family of 5mC hydroxylases [151] Accumulated fumarate competitively inhibits the α -KG dependent dioxygenases, including histone demethylases and TET family of 5mC hydroxylases [151] Inhibits KDM2B-mediated histone H3 K36 demethylation and promotes DNA repair and cell survival [13,152] O-GlcNAcylation of FH blocks AMPK-mediated phosphorylation; FH fails to form the FH-ATF2 complex and loses its transcriptional regulatory activity; tumor is maintained growth under glucose deficiency [153] Gains new activity to convert α -KG to L-2-hydroxyglutarate (L-2-HG) at acidic pH, thereby stabilizing HIF-1 α [132]
	IDH2	Mitochondria		Mitochondria	
	SDH	Mitochondria	Converts succinate to fumarate (wild type) Accumulates succinate (mutant)	Nucleus	
	FH	Mitochondria	Converts fumarate to L-malate (wild-type) Accumulates fumarate (mutant)	Nucleus	
	MDH1	Cytoplasm	Converts malate to oxaloacetate	Cytoplasm	
	MDH2	Mitochondria		Mitochondria	
Glutaminolysis	GLS1	Mitochondria	Converts glutamine to glutamate	Mitochondria	Senses glutamine availability and initiates mitochondria fusion to help the cells to overcome energy crisis under low glutamine status [112]
	GDH	Mitochondria	Converts glutamate to α -ketoglutarate and ammonia	Mitochondria	Manipulates the intracellular level of fumarate, which activates glutathione peroxidase 1 to scavenge reactive oxygen species, thereby maintaining redox homeostasis [116] In LK1-B deficient lung cancer, α -KG produced by GDH1 activates CamKK2-AMPK signaling, resulting in energy production [119] Under low glucose stimulation, α -KG produced by GDH1 directly activates NF- κ B signaling, upregulates GLUT1 expression, and increases glucose uptake [120,121]
Fructose metabolism	KHK-A	Cytoplasm	Converts fructose to fructose-1-phosphate	Cytoplasm	Phosphorylates and activates PRPS1, thereby promoting <i>de novo</i> nucleic acid synthesis and HCC tumorigenesis through pentose phosphate pathway [125] KHK-A S80 phosphorylation phosphorylates p62 and prevents its ubiquitylation, thereby promoting gene expression via Nrf2 activation to reduce reactive oxygen species [126]
Others	ACLY	Cytoplasm	Converts citrate to acetyl-CoA	Nucleus	Under growth factor stimulation and cell differentiation, ACLY-generated acetyl-CoA promotes histone acetylation and gene expression [155,156]

Fumarate hydratase (FH)

FH, also known as fumarase, is a key mitochondrial metabolic enzyme in the TCA cycle [149]. The typical function of FH is catalyzing the reversible conversion of fumarate to L-malate [150]. FH and another TCA cycle enzyme, succinate dehydrogenase (SDH), which converts succinate to fumarate, are both mutated in cancer cells, resulting in the accumulation of their substrates, fumarate and succinate, respectively. Fumarate and succinate are competitive inhibitors of multiple α -KG dependent dioxygenases, including histone demethylases and the ten-11 translocation (TET) family of 5mC hydroxylases [151]. Nuclear FH is phosphorylated by DNA-PK at T236 and recruited to double-strand break (DSB) regions in response to ionizing radiation (IR), where it generates fumarate to locally inhibit KDM2B-mediated histone H3 K36 demethylation, which in turn promotes DNA repair and cell survival [13,152]. While under glucose deprivation, AMPK phosphorylates FH at S75, thereby promoting FH-ATF2 binding. The complex translocates into the nucleus and binds to the promoters of ATF2 target genes. FH-catalyzed fumarate locally inhibits KDM2A activity and thus maintains the H3K36me2 level to facilitate target gene expression and to promote cell growth arrest. Notably, O-GlcNAcylation of FH blocks AMPK phosphorylation site. FH fails to form the FH-ATF2 complex and loses its transcriptional regulatory activity in the nucleus, thereby maintaining tumor growth under glucose deficiency [153] (Fig. 11).

ATP-citrate lyase (ACLY)

ACLY converts citrate into acetyl-CoA and is upregulated in several types of tumor. Under high glucose conditions, ACLY is acetylated by PCAF at lysine K540, K546, and K554; the blocking of its ubiquitylation and degradation stabilizes ACLY and promotes *de novo* lipid synthesis, cell proliferation, and tumor growth [154]. Surprisingly, Wellen *et al.* demonstrated that ACLY-generated acetyl-CoA is the main source for histone acetylation in mammalian cells. Growth factor stimulation and glucose availability promote ACLY-dependent histone acetylation, which is required for the expressions of hexokinase 2 (HK2), phosphofructokinase-1 (PFK-1), lactate dehydrogenase A (LDH-A), and glucose transporter 4 (GLUT4). Therefore, knockdown of *ACLY* results in a 32% decrease of glucose consumption, which links growth factor and glucose availability to the regulation of histone acetylation and metabolic alteration [155,156] (Fig. 11).

Discussion

Approximately 19 000 protein coding genes are present in

the human genome, but only hundreds of moonlighting proteins have been identified so far [157]. The number of moonlighting proteins will increase quickly, as the significance of protein moonlighting in biology and translational medicine is becoming increasingly apparent (Table 1). Study results have demonstrated that moonlighting proteins can execute novel biological functions, thereby broadening the pool of human functional proteome and simultaneously providing tumor cell growth advantage and therapeutic vulnerability.

Biological function: protein moonlighting provides rapid and efficient response to oncogenic mutations

Proteins moonlighting offers functional options for tumor cells to respond to metabolic stress without the transcription and translation of new proteins. Furthermore, some moonlighting proteins directly bind to metabolic enzymes and regulate their activity. For example, oncoprotein KRAS4A uniquely and directly binds to and activates HK1, thereby enhancing glycolytic flux [24]. Tumor suppressor p53 can also directly bind to G6PD and suppress its activity, thereby repressing glucose consumption, NADPH production, and biosynthesis. Intriguingly, tumor-derived p53 mutation abolishes its inhibition to G6PD, thereby enhancing PPP flux, promoting glucose consumption, and diverting more glucose to biosynthesis to support the rapid growth and proliferation of tumor cells [30]. These direct bindings provide rapid and efficient response to oncogenic mutations, which is important for the unconstrained proliferation of tumor cells.

Translational implication: protein moonlighting provides therapeutic vulnerability

Cancer cells reprogram metabolic pathways by protein moonlighting in response to external signals to support cell survival and proliferation, thereby providing a growth advantage and a therapeutic target for cancer treatment. Both the cytoplasmic and nuclear functions of PKM2 are important for cell growth and proliferation and are controlled by tetramer-dimer transition, thereby providing an excellent target for the treatment of cancer. PKM2 activator increases its metabolic activity in the cytoplasm, which diverts biomass away from biosynthesis, and inhibits protein kinase activity by promoting dimer-tetramer transition [98]. PKM2 is acetylated by p300 acetyltransferase at K433, and K433 acetylation acts as a switch for the shift of the metabolic and nuclear function of PKM2; thus, K433 acetylation is a promising target for cancer treatment [81]. Targeting p300 could increase its enzymatic activity and inhibit its nuclear function as protein kinase and co-transcription factor [158], thereby repressing tumor growth. The direct binding of KRAS4A to HK1 also can be targeted for patients bearing KRAS

oncogenic mutation to block the glycolytic flux. Exploring and targeting the moonlighting function is a promising strategy for the clinical treatment of cancer.

Research prospect: metabolites regulate signaling pathways

Cancer metabolism is regulated by multiple signaling pathways and vice versa. Metabolites provide feedback to coordinate signaling pathways to reprogram metabolism. α -KG generated by GDH1 plays an important role in the regulation of different cellular signaling pathways, such as activation of CamKK2-AMPK [119] and NF- κ B pathways [120,121], to reprogram metabolism. Fumarate binds to and activates GPx1 to scavenge ROS and to maintain redox homeostasis [116]. As analogs of α -KG, 2-HG, fumarate, and succinate can potentially inhibit more than 60 α -KG-dependent dioxygenases, including histone and DNA demethylase, HIF, and AlkB proteins. Identifying new signaling pathways regulated by metabolites and checking whether 2-HG, fumarate, succinate, and α -KG could regulate CamKK2-AMPK, NF- κ B, and GPx1 antioxidant pathways would be interesting.

More than 110 000 metabolites, nearly 3500 proteins related to metabolism dysregulation, and the tumor and metabolomics data of multiple tumors, including liver, breast, kidney, and colorectal cancers, are present in the Human Metabolome Database (HMDB). By means of bioinformatics, in-depth mining in the existing database would reveal the metabolic characteristics of different tumors, leading to the discovery of metabolic pathways and important metabolites closely related to tumor occurrence and development.

Acknowledgements

This work was supported by the Ministry of Science and Technology (No. 2019YFA0801703), the National Natural Science Foundation of China (Nos. 81790250, 81790253, 91959202, 81902823, and 81972620), and the Innovation Program of Shanghai Municipal Education Commission (No. N173606).

Compliance with ethics guidelines

Lei Lv and Qunying Lei declare that they have no conflict of interest. This manuscript is a review article and does not involve a research protocol requiring approval by the relevant institutional review board or ethics committee.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>.

References

- Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell* 2011; 144(5): 646–674
- Warburg O. On the origin of cancer cells. *Science* 1956; 123(3191): 309–314
- Allison KE, Coomber BL, Bridle BW. Metabolic reprogramming in the tumour microenvironment: a hallmark shared by cancer cells and T lymphocytes. *Immunology* 2017; 152(2): 175–184
- Rodrigues MF, Obre E, de Melo FH, Santos GC Jr, Galina A, Jasiulionis MG, Rossignol R, Rumjanek FD, Amoêdo ND. Enhanced OXPHOS, glutaminolysis and β -oxidation constitute the metastatic phenotype of melanoma cells. *Biochem J* 2016; 473(6): 703–715
- Caro P, Kishan AU, Norberg E, Stanley IA, Chapuy B, Ficarro SB, Polak K, Tondera D, Gounarides J, Yin H, Zhou F, Green MR, Chen L, Monti S, Marto JA, Shipp MA, Danial NN. Metabolic signatures uncover distinct targets in molecular subsets of diffuse large B cell lymphoma. *Cancer Cell* 2012; 22(4): 547–560
- Liu Y, Zuckier LS, Ghesani NV. Dominant uptake of fatty acid over glucose by prostate cells: a potential new diagnostic and therapeutic approach. *Anticancer Res* 2010; 30(2): 369–374
- Lazar I, Clement E, Dauvillier S, Milhas D, Ducoux-Petit M, LeGonidec S, Moro C, Soldan V, Dalle S, Balor S, Golzio M, Burlet-Schiltz O, Valet P, Muller C, Nieto L. Adipocyte exosomes promote melanoma aggressiveness through fatty acid oxidation: a novel mechanism linking obesity and cancer. *Cancer Res* 2016; 76(14): 4051–4057
- Nieman KM, Kenny HA, Penicka CV, Ladanyi A, Buell-Gutbrod R, Zillhardt MR, Romero IL, Carey MS, Mills GB, Hotamisligil GS, Yamada SD, Peter ME, Gwin K, Lengyel E. Adipocytes promote ovarian cancer metastasis and provide energy for rapid tumor growth. *Nat Med* 2011; 17(11): 1498–1503
- Cao Y. Adipocyte and lipid metabolism in cancer drug resistance. *J Clin Invest* 2019; 129(8): 3006–3017
- Iwamoto H, Abe M, Yang Y, Cui D, Seki T, Nakamura M, Hosaka K, Lim S, Wu J, He X, Sun X, Lu Y, Zhou Q, Shi W, Torimura T, Nie G, Li Q, Cao Y. Cancer lipid metabolism confers antiangiogenic drug resistance. *Cell Metab* 2018; 28(1): 104–117.e5
- Ward PS, Thompson CB. Metabolic reprogramming: a cancer hallmark even Warburg did not anticipate. *Cancer Cell* 2012; 21(3): 297–308
- Reina-Campos M, Moscat J, Diaz-Meco M. Metabolism shapes the tumor microenvironment. *Curr Opin Cell Biol* 2017; 48: 47–53
- Boukouris AE, Zervopoulos SD, Michelakis ED. Metabolic

- enzymes moonlighting in the nucleus: metabolic regulation of gene transcription. *Trends Biochem Sci* 2016; 41(8): 712–730
14. Li M, Zhang CS, Zong Y, Feng JW, Ma T, Hu M, Lin Z, Li X, Xie C, Wu Y, Jiang D, Li Y, Zhang C, Tian X, Wang W, Yang Y, Chen J, Cui J, Wu YQ, Chen X, Liu QF, Wu J, Lin SY, Ye Z, Liu Y, Piao HL, Yu L, Zhou Z, Xie XS, Hardie DG, Lin SC. Transient receptor potential V channels are essential for glucose sensing by aldolase and AMPK. *Cell Metab* 2019; 30(3): 508–524.e12
 15. Huangyang P, Li F, Lee P, Nissim I, Weljie AM, Mancuso A, Li B, Keith B, Yoon SS, Simon MC. Fructose-1,6-bisphosphatase 2 inhibits sarcoma progression by restraining mitochondrial biogenesis. *Cell Metab* 2020; 31(1): 174–188.e7
 16. Xu D, Wang Z, Xia Y, Shao F, Xia W, Wei Y, Li X, Qian X, Lee JH, Du L, Zheng Y, Lv G, Leu JS, Wang H, Xing D, Liang T, Hung MC, Lu Z. The gluconeogenic enzyme PCK1 phosphorylates INSIG1/2 for lipogenesis. *Nature* 2020; 580(7804): 530–535
 17. Fernández-Medarde A, Santos E. Ras in cancer and developmental diseases. *Genes Cancer* 2011; 2(3): 344–358
 18. Kerr EM, Gaude E, Turrell FK, Frezza C, Martins CP. Mutant Kras copy number defines metabolic reprogramming and therapeutic susceptibilities. *Nature* 2016; 531(7592): 110–113
 19. Cox AD, Der CJ. Ras history: the saga continues. *Small GTPases* 2010; 1(1): 2–27
 20. Simanshu DK, Nissley DV, McCormick F. RAS proteins and their regulators in human disease. *Cell* 2017; 170(1): 17–33
 21. Hallin J, Engstrom LD, Hargis L, Calinisan A, Aranda R, Briere DM, Sudhakar N, Bowcut V, Baer BR, Ballard JA, Burkard MR, Fell JB, Fischer JP, Vigers GP, Xue Y, Gatto S, Fernandez-Banet J, Pavlicek A, Velastagui K, Chao RC, Barton J, Pierobon M, Baldelli E, Patricoin EF 3rd, Cassidy DP, Marx MA, Rybkin II, Johnson ML, Ou SI, Lito P, Papadopoulos KP, Jänne PA, Olson P, Christensen JG. The KRAS^{G12C} inhibitor MRTX849 provides insight toward therapeutic susceptibility of KRAS-mutant cancers in mouse models and patients. *Cancer Discov* 2020; 10(1): 54–71
 22. Kimmelman AC. Metabolic dependencies in RAS-driven cancers. *Clin Cancer Res* 2015; 21(8): 1828–1834
 23. Ying H, Kimmelman AC, Lyssiotis CA, Hua S, Chu GC, Fletcher-Sanankone E, Locasale JW, Son J, Zhang H, Colloff JL, Yan H, Wang W, Chen S, Viale A, Zheng H, Paik JH, Lim C, Guimaraes AR, Martin ES, Chang J, Hezel AF, Perry SR, Hu J, Gan B, Xiao Y, Asara JM, Weissleder R, Wang YA, Chin L, Cantley LC, DePinho RA. Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* 2012; 149(3): 656–670
 24. Amendola CR, Mahaffey JP, Parker SJ, Ahearn IM, Chen WC, Zhou M, Court H, Shi J, Mendoza SL, Morten MJ, Rothenberg E, Gottlieb E, Wadghiri YZ, Possemato R, Hubbard SR, Balmain A, Kimmelman AC, Phillips MR. KRAS4A directly regulates hexokinase 1. *Nature* 2019; 576(7787): 482–486
 25. Vousden KH, Ryan KM. p53 and metabolism. *Nat Rev Cancer* 2009; 9(10): 691–700
 26. Kawauchi K, Araki K, Tobiume K, Tanaka N. p53 regulates glucose metabolism through an IKK-NF- κ B pathway and inhibits cell transformation. *Nat Cell Biol* 2008; 10(5): 611–618
 27. Chipuk JE, Kuwana T, Bouchier-Hayes L, Droin NM, Newmeyer DD, Schuler M, Green DR. Direct activation of Bax by p53 mediates mitochondrial membrane permeabilization and apoptosis. *Science* 2004; 303(5660): 1010–1014
 28. Marchenko ND, Wolff S, Erster S, Becker K, Moll UM. Monoubiquitylation promotes mitochondrial p53 translocation. *EMBO J* 2007; 26(4): 923–934
 29. Park JH, Zhuang J, Li J, Hwang PM. p53 as guardian of the mitochondrial genome. *FEBS Lett* 2016; 590(7): 924–934
 30. Jiang P, Du W, Wang X, Mancuso A, Gao X, Wu M, Yang X. p53 regulates biosynthesis through direct inactivation of glucose-6-phosphate dehydrogenase. *Nat Cell Biol* 2011; 13(3): 310–316
 31. Hillmer EJ, Zhang H, Li HS, Watowich SS. STAT3 signaling in immunity. *Cytokine Growth Factor Rev* 2016; 31: 1–15
 32. Bromberg JF, Wrzeszczynska MH, Devgan G, Zhao Y, Pestell RG, Albanese C, Darnell JE Jr. Stat3 as an oncogene. *Cell* 1999; 98(3): 295–303
 33. Levy DE, Lee CK. What does Stat3 do? *J Clin Invest* 2002; 109(9): 1143–1148
 34. Guanizo AC, Fernando CD, Garama DJ, Gough DJ. STAT3: a multifaceted oncoprotein. *Growth Factors* 2018; 36(1-2): 1–14
 35. Zhao S, Venkatasubbarao K, Lazor JW, Sperry J, Jin C, Cao L, Freeman JW. Inhibition of STAT3 Tyr705 phosphorylation by Smad4 suppresses transforming growth factor beta-mediated invasion and metastasis in pancreatic cancer cells. *Cancer Res* 2008; 68(11): 4221–4228
 36. Bollrath J, Pheesse TJ, von Burstin VA, Putoczki T, Bennecke M, Bateman T, Nebelsiek T, Lundgren-May T, Canli O, Schwitala S, Matthews V, Schmid RM, Kirchner T, Arkan MC, Ernst M, Greten FR. gp130-mediated Stat3 activation in enterocytes regulates cell survival and cell-cycle progression during colitis-associated tumorigenesis. *Cancer Cell* 2009; 15(2): 91–102
 37. Wegrzyn J, Potla R, Chwae YJ, Sepuri NB, Zhang Q, Koeck T, Derecka M, Szczepanek K, Szelag M, Gornicka A, Moh A, Moghaddas S, Chen Q, Bobbili S, Cichy J, Dulak J, Baker DP, Wolfman A, Stuehr D, Hassan MO, Fu XY, Avadhani N, Drake JL, Fawcett P, Lesnefsky EJ, Larner AC. Function of mitochondrial Stat3 in cellular respiration. *Science* 2009; 323(5915): 793–797
 38. Garama DJ, White CL, Balic JJ, Gough DJ. Mitochondrial STAT3: powering up a potent factor. *Cytokine* 2016; 87: 20–25
 39. Genini D, Brambilla L, Laurini E, Merulla J, Civenni G, Pandit S, D’Antuono R, Perez L, Levy DE, Priel S, Carbone GM, Catapano CV. Mitochondrial dysfunction induced by a SH2 domain-targeting STAT3 inhibitor leads to metabolic synthetic lethality in cancer cells. *Proc Natl Acad Sci USA* 2017; 114(25): E4924–E4933
 40. Pelengaris S, Khan M. The many faces of c-MYC. *Arch Biochem Biophys* 2003; 416(2): 129–136
 41. Kuzyk A, Mai S. c-MYC-induced genomic instability. *Cold Spring Harb Perspect Med* 2014; 4(4): a014373
 42. Dang CV. MYC on the path to cancer. *Cell* 2012; 149(1): 22–35
 43. Kumari A, Folk WP, Sakamuro D. The dual roles of MYC in genomic instability and cancer chemoresistance. *Genes (Basel)* 2017; 8(6): E158
 44. Dejure FR, Eilers M. MYC and tumor metabolism: chicken and egg. *EMBO J* 2017; 36(23): 3409–3420
 45. Pavlova NN, Thompson CB. The emerging hallmarks of cancer metabolism. *Cell Metab* 2016; 23(1): 27–47
 46. Shim H, Dolde C, Lewis BC, Wu CS, Dang G, Jungmann RA, Dalla-Favera R, Dang CV. c-Myc transactivation of LDH-A:

- implications for tumor metabolism and growth. *Proc Natl Acad Sci USA* 1997; 94(13): 6658–6663
47. Fang Y, Shen ZY, Zhan YZ, Feng XC, Chen KL, Li YS, Deng HJ, Pan SM, Wu DH, Ding Y. CD36 inhibits β -catenin/c-myc-mediated glycolysis through ubiquitination of GPC4 to repress colorectal tumorigenesis. *Nat Commun* 2019; 10(1): 3981
 48. David CJ, Chen M, Assanah M, Canoll P, Manley JL. HnRNP proteins controlled by c-Myc deregulate pyruvate kinase mRNA splicing in cancer. *Nature* 2010; 463(7279): 364–368
 49. Lee KM, Giltman JM, Balko JM, Schwarz LJ, Guerrero-Zotano AL, Hutchinson KE, Nixon MJ, Estrada MV, Sánchez V, Sanders ME, Lee T, Gómez H, Lluch A, Pérez-Fidalgo JA, Wolf MM, Andrejeva G, Rathmell JC, Fesik SW, Arteaga CL. MYC and MCL1 cooperatively promote chemotherapy-resistant breast cancer stem cells via regulation of mitochondrial oxidative phosphorylation. *Cell Metab* 2017; 26(4): 633–647.e7
 50. Pelicano H, Martin DS, Xu RH, Huang P. Glycolysis inhibition for anticancer treatment. *Oncogene* 2006; 25(34): 4633–4646
 51. Ali M, Rellos P, Cox TM. Hereditary fructose intolerance. *J Med Genet* 1998; 35(5): 353–365
 52. Chang YC, Yang YC, Tien CP, Yang CJ, Hsiao M. Roles of aldolase family genes in human cancers and diseases. *Trends Endocrinol Metab* 2018; 29(8): 549–559
 53. Rose IA, O'Connell EL. Studies on the interaction of aldolase with substrate analogues. *J Biol Chem* 1969; 244(1): 126–134
 54. Zhang CS, Hawley SA, Zong Y, Li M, Wang Z, Gray A, Ma T, Cui J, Feng JW, Zhu M, Wu YQ, Li TY, Ye Z, Lin SY, Yin H, Piao HL, Hardie DG, Lin SC. Fructose-1,6-bisphosphate and aldolase mediate glucose sensing by AMPK. *Nature* 2017; 548(7665): 112–116
 55. Herzig S, Shaw RJ. AMPK: guardian of metabolism and mitochondrial homeostasis. *Nat Rev Mol Cell Biol* 2018; 19(2): 121–135
 56. Tristan C, Shahani N, Sedlak TW, Sawa A. The diverse functions of GAPDH: views from different subcellular compartments. *Cell Signal* 2011; 23(2): 317–323
 57. Yun J, Mullarky E, Lu C, Bosch KN, Kavalier A, Rivera K, Roper J, Chio II, Giannopoulou EG, Rago C, Muley A, Asara JM, Paik J, Elemento O, Chen Z, Pappin DJ, Dow LE, Papadopoulos N, Gross SS, Cantley LC. Vitamin C selectively kills KRAS and BRAF mutant colorectal cancer cells by targeting GAPDH. *Science* 2015; 350(6266): 1391–1396
 58. Rodríguez-Pascual F, Redondo-Horcajo M, Magán-Marchal N, Lagares D, Martínez-Ruiz A, Kleinert H, Lamas S. Glyceraldehyde-3-phosphate dehydrogenase regulates endothelin-1 expression by a novel, redox-sensitive mechanism involving mRNA stability. *Mol Cell Biol* 2008; 28(23): 7139–7155
 59. Hara MR, Agrawal N, Kim SF, Cascio MB, Fujimuro M, Ozeki Y, Takahashi M, Cheah JH, Tankou SK, Hester LD, Ferris CD, Hayward SD, Snyder SH, Sawa A. S-nitrosylated GAPDH initiates apoptotic cell death by nuclear translocation following Siah1 binding. *Nat Cell Biol* 2005; 7(7): 665–674
 60. Kornberg MD, Sen N, Hara MR, Juluri KR, Nguyen JV, Snowman AM, Law L, Hester LD, Snyder SH. GAPDH mediates nitrosylation of nuclear proteins. *Nat Cell Biol* 2010; 12(11): 1094–1100
 61. Sen N, Hara MR, Kornberg MD, Cascio MB, Bae BI, Shahani N, Thomas B, Dawson TM, Dawson VL, Snyder SH, Sawa A. Nitric oxide-induced nuclear GAPDH activates p300/CBP and mediates apoptosis. *Nat Cell Biol* 2008; 10(7): 866–873
 62. Colell A, Ricci JE, Tait S, Milasta S, Maurer U, Bouchier-Hayes L, Fitzgerald P, Guio-Carrion A, Waterhouse NJ, Li CW, Mari B, Barbry P, Newmeyer DD, Beere HM, Green DR. GAPDH and autophagy preserve survival after apoptotic cytochrome c release in the absence of caspase activation. *Cell* 2007; 129(5): 983–997
 63. Chang C, Su H, Zhang D, Wang Y, Shen Q, Liu B, Huang R, Zhou T, Peng C, Wong CC, Shen HM, Lippincott-Schwartz J, Liu W. AMPK-dependent phosphorylation of GAPDH triggers Sirt1 activation and is necessary for autophagy upon glucose starvation. *Mol Cell* 2015; 60(6): 930–940
 64. Zheng L, Roeder RG, Luo Y. S phase activation of the histone H2B promoter by OCA-S, a coactivator complex that contains GAPDH as a key component. *Cell* 2003; 114(2): 255–266
 65. Sirover MA. Subcellular dynamics of multifunctional protein regulation: mechanisms of GAPDH intracellular translocation. *J Cell Biochem* 2012; 113(7): 2193–2200
 66. Zhang Y, Yu G, Chu H, Wang X, Xiong L, Cai G, Liu R, Gao H, Tao B, Li W, Li G, Liang J, Yang W. Macrophage-associated PGK1 phosphorylation promotes aerobic glycolysis and tumorigenesis. *Mol Cell* 2018; 71(2): 201–215.e7
 67. Hu H, Zhu W, Qin J, Chen M, Gong L, Li L, Liu X, Tao Y, Yin H, Zhou H, Zhou L, Ye D, Ye Q, Gao D. Acetylation of PGK1 promotes liver cancer cell proliferation and tumorigenesis. *Hepatology* 2017; 65(2): 515–528
 68. Nie H, Ju H, Fan J, Shi X, Cheng Y, Cang X, Zheng Z, Duan X, Yi W. O-GlcNAcylation of PGK1 coordinates glycolysis and TCA cycle to promote tumor growth. *Nat Commun* 2020; 11(1): 36
 69. Li X, Jiang Y, Meisenhelder J, Yang W, Hawke DH, Zheng Y, Xia Y, Aldape K, He J, Hunter T, Wang L, Lu Z. Mitochondria-translocated PGK1 functions as a protein kinase to coordinate glycolysis and the TCA cycle in tumorigenesis. *Mol Cell* 2016; 61(5): 705–719
 70. Qian X, Li X, Shi Z, Xia Y, Cai Q, Xu D, Tan L, Du L, Zheng Y, Zhao D, Zhang C, Lorenzi PL, You Y, Jiang BH, Jiang T, Li H, Lu Z. PTEN suppresses glycolysis by dephosphorylating and inhibiting autophosphorylated PGK1. *Mol Cell* 2019; 76(3): 516–527.e7
 71. Qian X, Li X, Cai Q, Zhang C, Yu Q, Jiang Y, Lee JH, Hawke D, Wang Y, Xia Y, Zheng Y, Jiang BH, Liu DX, Jiang T, Lu Z. Phosphoglycerate kinase 1 phosphorylates beclin1 to induce autophagy. *Mol Cell* 2017; 65(5): 917–931.e6
 72. Qian X, Li X, Lu Z. Protein kinase activity of the glycolytic enzyme PGK1 regulates autophagy to promote tumorigenesis. *Autophagy* 2017; 13(7): 1246–1247
 73. Liang C, Shi S, Qin Y, Meng Q, Hua J, Hu Q, Ji S, Zhang B, Xu J, Yu XJ. Localisation of PGK1 determines metabolic phenotype to balance metastasis and proliferation in patients with SMAD4-negative pancreatic cancer. *Gut* 2020; 69(5): 888–900
 74. Dayton TL, Jacks T, Vander Heiden MG. PKM2, cancer metabolism, and the road ahead. *EMBO Rep* 2016; 17(12): 1721–1730
 75. Mazurek S. Pyruvate kinase type M2: a key regulator of the metabolic budget system in tumor cells. *Int J Biochem Cell Biol* 2011; 43(7): 969–980
 76. Sun Q, Chen X, Ma J, Peng H, Wang F, Zha X, Wang Y, Jing Y, Yang H, Chen R, Chang L, Zhang Y, Goto J, Onda H, Chen T,

- Wang MR, Lu Y, You H, Kwiatkowski D, Zhang H. Mammalian target of rapamycin up-regulation of pyruvate kinase isoenzyme type M2 is critical for aerobic glycolysis and tumor growth. *Proc Natl Acad Sci USA* 2011; 108(10): 4129–4134
77. Christofk HR, Vander Heiden MG, Harris MH, Ramanathan A, Gerszten RE, Wei R, Fleming MD, Schreiber SL, Cantley LC. The M2 splice isoform of pyruvate kinase is important for cancer metabolism and tumour growth. *Nature* 2008; 452(7184): 230–233
 78. Christofk HR, Vander Heiden MG, Wu N, Asara JM, Cantley LC. Pyruvate kinase M2 is a phosphotyrosine-binding protein. *Nature* 2008; 452(7184): 181–186
 79. Lv L, Li D, Zhao D, Lin R, Chu Y, Zhang H, Zha Z, Liu Y, Li Z, Xu Y, Wang G, Huang Y, Xiong Y, Guan KL, Lei QY. Acetylation targets the M2 isoform of pyruvate kinase for degradation through chaperone-mediated autophagy and promotes tumor growth. *Mol Cell* 2011; 42(6): 719–730
 80. Macintyre AN, Rathmell JC. PKM2 and the tricky balance of growth and energy in cancer. *Mol Cell* 2011; 42(6): 713–714
 81. Lv L, Xu YP, Zhao D, Li FL, Wang W, Sasaki N, Jiang Y, Zhou X, Li TT, Guan KL, Lei QY, Xiong Y. Mitogenic and oncogenic stimulation of K433 acetylation promotes PKM2 protein kinase activity and nuclear localization. *Mol Cell* 2013; 52(3): 340–352
 82. Yang W, Zheng Y, Xia Y, Ji H, Chen X, Guo F, Lyssiotis CA, Aldape K, Cantley LC, Lu Z. ERK1/2-dependent phosphorylation and nuclear translocation of PKM2 promotes the Warburg effect. *Nat Cell Biol* 2012; 14(12): 1295–1304
 83. Yang W, Xia Y, Ji H, Zheng Y, Liang J, Huang W, Gao X, Aldape K, Lu Z. Nuclear PKM2 regulates β -catenin transactivation upon EGFR activation. *Nature* 2011; 480(7375): 118–122
 84. Li S, Swanson SK, Gogol M, Florens L, Washburn MP, Workman JL, Saganuma T. Serine and SAM responsive complex SESAME regulates histone modification crosstalk by sensing cellular metabolism. *Mol Cell* 2015; 60(3): 408–421
 85. Yang W, Xia Y, Hawke D, Li X, Liang J, Xing D, Aldape K, Hunter T, Alfred Yung WK, Lu Z. PKM2 phosphorylates histone H3 and promotes gene transcription and tumorigenesis. *Cell* 2012; 150(4): 685–696
 86. Hsu MC, Hung WC. Pyruvate kinase M2 fuels multiple aspects of cancer cells: from cellular metabolism, transcriptional regulation to extracellular signaling. *Mol Cancer* 2018; 17(1): 35
 87. Gao X, Wang H, Yang JJ, Liu X, Liu ZR. Pyruvate kinase M2 regulates gene transcription by acting as a protein kinase. *Mol Cell* 2012; 45(5): 598–609
 88. Keller KE, Doctor ZM, Dwyer ZW, Lee YS. SAICAR induces protein kinase activity of PKM2 that is necessary for sustained proliferative signaling of cancer cells. *Mol Cell* 2014; 53(5): 700–709
 89. Luo W, Hu H, Chang R, Zhong J, Knabel M, O’Meally R, Cole RN, Pandey A, Semenza GL. Pyruvate kinase M2 is a PHD3-stimulated coactivator for hypoxia-inducible factor 1. *Cell* 2011; 145(5): 732–744
 90. Wang HJ, Hsieh YJ, Cheng WC, Lin CP, Lin YS, Yang SF, Chen CC, Izumiya Y, Yu JS, Kung HJ, Wang WC. JMJD5 regulates PKM2 nuclear translocation and reprograms HIF-1 α -mediated glucose metabolism. *Proc Natl Acad Sci USA* 2014; 111(1): 279–284
 91. Demaria M, Poli V. PKM2, STAT3 and HIF-1 α : the Warburg’s vicious circle. *JAK-STAT* 2012; 1(3): 194–196
 92. Azoitei N, Becher A, Steinestel K, Rouhi A, Diepold K, Genze F, Simmet T, Seufferlein T. PKM2 promotes tumor angiogenesis by regulating HIF-1 α through NF- κ B activation. *Mol Cancer* 2016; 15(1): 3
 93. Hosios AM, Fiske BP, Gui DY, Vander Heiden MG. Lack of evidence for PKM2 protein kinase activity. *Mol Cell* 2015; 59(5): 850–857
 94. Vander Heiden MG, Christofk HR, Schuman E, Subtelny AO, Sharfi H, Harlow EE, Xian J, Cantley LC. Identification of small molecule inhibitors of pyruvate kinase M2. *Biochem Pharmacol* 2010; 79(8): 1118–1124
 95. Wang Y, Hao F, Nan Y, Qu L, Na W, Jia C, Chen X. PKM2 inhibitor shikonin overcomes the cisplatin resistance in bladder cancer by inducing necroptosis. *Int J Biol Sci* 2018; 14(13): 1883–1891
 96. Chen J, Xie J, Jiang Z, Wang B, Wang Y, Hu X. Shikonin and its analogs inhibit cancer cell glycolysis by targeting tumor pyruvate kinase-M2. *Oncogene* 2011; 30(42): 4297–4306
 97. Kung C, Hixon J, Choe S, Marks K, Gross S, Murphy E, DeLaBarre B, Cianchetta G, Sethumadhavan S, Wang X, Yan S, Gao Y, Fang C, Wei W, Jiang F, Wang S, Qian K, Saunders J, Driggers E, Woo HK, Kunii K, Murray S, Yang H, Yen K, Liu W, Cantley LC, Vander Heiden MG, Su SM, Jin S, Salituro FG, Dang L. Small molecule activation of PKM2 in cancer cells induces serine auxotrophy. *Chem Biol* 2012; 19(9): 1187–1198
 98. Anastasiou D, Yu Y, Israelsen WJ, Jiang JK, Boxer MB, Hong BS, Tempel W, Dimov S, Shen M, Jha A, Yang H, Mattaini KR, Metallo CM, Fiske BP, Courtney KD, Malstrom S, Khan TM, Kung C, Skoumbourdis AP, Veith H, Southall N, Walsh MJ, Brimacombe KR, Leister W, Lunt SY, Johnson ZR, Yen KE, Kunii K, Davidson SM, Christofk HR, Austin CP, Inglese J, Harris MH, Asara JM, Stephanopoulos G, Salituro FG, Jin S, Dang L, Auld DS, Park HW, Cantley LC, Thomas CJ, Vander Heiden MG. Pyruvate kinase M2 activators promote tetramer formation and suppress tumorigenesis. *Nat Chem Biol* 2012; 8(10): 839–847
 99. Parnell KM, Foulks JM, Nix RN, Clifford A, Bullough J, Luo B, Senina A, Vollmer D, Liu J, McCarthy V, Xu Y, Saunders M, Liu XH, Pearce S, Wright K, O’Reilly M, McCullar MV, Ho KK, Kanner SB. Pharmacologic activation of PKM2 slows lung tumor xenograft growth. *Mol Cancer Ther* 2013; 12(8): 1453–1460
 100. Dong C, Yuan T, Wu Y, Wang Y, Fan TW, Miriyala S, Lin Y, Yao J, Shi J, Kang T, Lorkiewicz P, St Clair D, Hung MC, Evers BM, Zhou BP. Loss of FBP1 by Snail-mediated repression provides metabolic advantages in basal-like breast cancer. *Cancer Cell* 2013; 23(3): 316–331
 101. Zhang J, Wang J, Xing H, Li Q, Zhao Q, Li J. Down-regulation of FBP1 by ZEB1-mediated repression confers to growth and invasion in lung cancer cells. *Mol Cell Biochem* 2016; 411(1–2): 331–340
 102. Hirata H, Sugimachi K, Komatsu H, Ueda M, Masuda T, Uchi R, Sakimura S, Nambara S, Saito T, Shinden Y, Iguchi T, Eguchi H, Ito S, Terashima K, Sakamoto K, Hirakawa M, Honda H, Mimori K. Decreased expression of fructose-1,6-bisphosphatase associates with glucose metabolism and tumor progression in hepatocellular carcinoma. *Cancer Res* 2016; 76(11): 3265–3276
 103. Son B, Lee S, Kim H, Kang H, Jeon J, Jo S, Seong KM, Lee SJ,

- Youn H, Youn B. Decreased FBP1 expression rewires metabolic processes affecting aggressiveness of glioblastoma. *Oncogene* 2020; 39(1): 36–49
104. Li B, Qiu B, Lee DS, Walton ZE, Ochocki JD, Mathew LK, Mancuso A, Gade TP, Keith B, Nissim I, Simon MC. Fructose-1,6-bisphosphatase opposes renal carcinoma progression. *Nature* 2014; 513(7517): 251–255
105. Lu C, Ren C, Yang T, Sun Y, Qiao P, Wang D, Lv S, Yu Z. A noncanonical role of fructose-1, 6-bisphosphatase 1 is essential for inhibition of Notch1 in breast cancer. *Mol Cancer Res* 2020; 18(5): 787–796
106. Cong J, Wang X, Zheng X, Wang D, Fu B, Sun R, Tian Z, Wei H. Dysfunction of natural killer cells by FBP1-induced inhibition of glycolysis during lung cancer progression. *Cell Metab* 2018; 28(2): 243–255.e5
107. Burgess SC, He T, Yan Z, Lindner J, Sherry AD, Malloy CR, Browning JD, Magnuson MA. Cytosolic phosphoenolpyruvate carboxykinase does not solely control the rate of hepatic gluconeogenesis in the intact mouse liver. *Cell Metab* 2007; 5(4): 313–320
108. Méndez-Lucas A, Hyroššová P, Novellasdemunt L, Viñals F, Perales JC. Mitochondrial phosphoenolpyruvate carboxykinase (PEPCK-M) is a pro-survival, endoplasmic reticulum (ER) stress response gene involved in tumor cell adaptation to nutrient availability. *J Biol Chem* 2014; 289(32): 22090–22102
109. Matés JM, Campos-Sandoval JA, Santos-Jiménez JL, Márquez J. Dysregulation of glutaminase and glutamine synthetase in cancer. *Cancer Lett* 2019; 467: 29–39
110. Mishra P, Chan DC. Metabolic regulation of mitochondrial dynamics. *J Cell Biol* 2016; 212(4): 379–387
111. Maycotte P, Marín-Hernández A, Goyri-Aguirre M, Anaya-Ruiz M, Reyes-Leyva J, Cortés-Hernández P. Mitochondrial dynamics and cancer. *Tumour Biol* 2017; 39(5): 1010428317698391
112. Cai WF, Zhang C, Wu YQ, Zhuang G, Ye Z, Zhang CS, Lin SC. Glutaminase GLS1 senses glutamine availability in a non-enzymatic manner triggering mitochondrial fusion. *Cell Res* 2018; 28(8): 865–867
113. Stillman TJ, Baker PJ, Britton KL, Rice DW. Conformational flexibility in glutamate dehydrogenase: role of water in substrate recognition and catalysis. *J Mol Biol* 1993; 234(4): 1131–1139
114. Zaganas I, Spanaki C, Plaitakis A. Expression of human GLUD2 glutamate dehydrogenase in human tissues: functional implications. *Neurochem Int* 2012; 61(4): 455–462
115. Michaelidis TM, Tzimagiorgis G, Moschonas NK, Papamatheakis J. The human glutamate dehydrogenase gene family: gene organization and structural characterization. *Genomics* 1993; 16(1): 150–160
116. Jin L, Li D, Alesi GN, Fan J, Kang HB, Lu Z, Boggon TJ, Jin P, Yi H, Wright ER, Duong D, Seyfried NT, Egnatchik R, DeBerardinis RJ, Magliocca KR, He C, Arellano ML, Khoury HJ, Shin DM, Khuri FR, Kang S. Glutamate dehydrogenase 1 signals through antioxidant glutathione peroxidase 1 to regulate redox homeostasis and tumor growth. *Cancer Cell* 2015; 27(2): 257–270
117. Fedeles BI, Singh V, Delaney JC, Li D, Essigmann JM. The AlkB family of Fe(II)/ α -ketoglutarate-dependent dioxygenases: repairing nucleic acid alkylation damage and beyond. *J Biol Chem* 2015; 290(34): 20734–20742
118. Yang C, Ko B, Hensley CT, Jiang L, Wasti AT, Kim J, Sudderth J, Calvaruso MA, Lumata L, Mitsche M, Rutter J, Merritt ME, DeBerardinis RJ. Glutamine oxidation maintains the TCA cycle and cell survival during impaired mitochondrial pyruvate transport. *Mol Cell* 2014; 56(3): 414–424
119. Jin L, Chun J, Pan C, Kumar A, Zhang G, Ha Y, Li D, Alesi GN, Kang Y, Zhou L, Yu WM, Magliocca KR, Khuri FR, Qu CK, Metallo C, Owonikoko TK, Kang S. The PLAG1–GDH1 axis promotes anoikis resistance and tumor metastasis through CamKK2–AMPK signaling in LKB1-deficient lung cancer. *Mol Cell* 2018; 69(1): 87–99.e7
120. Wang X, Liu R, Qu X, Yu H, Chu H, Zhang Y, Zhu W, Wu X, Gao H, Tao B, Li W, Liang J, Li G, Yang W. α -ketoglutarate-activated NF- κ B signaling promotes compensatory glucose uptake and brain tumor development. *Mol Cell* 2019; 76(1): 148–162.e7
121. Di Conza G, Tsai CH, Ho PC. Fifty shades of α -ketoglutarate on cellular programming. *Mol Cell* 2019; 76(1): 1–3
122. Diggle CP, Shires M, Leitch D, Brooke D, Carr IM, Markham AF, Hayward BE, Asipu A, Bonthron DT. Ketohexokinase: expression and localization of the principal fructose-metabolizing enzyme. *J Histochem Cytochem* 2009; 57(8): 763–774
123. Ishimoto T, Lanasa MA, Le MT, Garcia GE, Diggle CP, Maclean PS, Jackman MR, Asipu A, Roncal-Jimenez CA, Kosugi T, Rivard CJ, Maruyama S, Rodriguez-Iturbe B, Sánchez-Lozada LG, Bonthron DT, Sautin YY, Johnson RJ. Opposing effects of fructokinase C and A isoforms on fructose-induced metabolic syndrome in mice. *Proc Natl Acad Sci USA* 2012; 109(11): 4320–4325
124. Li X, Qian X, Lu Z. Fructokinase A acts as a protein kinase to promote nucleotide synthesis. *Cell Cycle* 2016; 15(20): 2689–2690
125. Li X, Qian X, Peng LX, Jiang Y, Hawke DH, Zheng Y, Xia Y, Lee JH, Cote G, Wang H, Wang L, Qian CN, Lu Z. A splicing switch from ketohexokinase-C to ketohexokinase-A drives hepatocellular carcinoma formation. *Nat Cell Biol* 2016; 18(5): 561–571
126. Xu D, Li X, Shao F, Lv G, Lv H, Lee JH, Qian X, Wang Z, Xia Y, Du L, Zheng Y, Wang H, Lyu J, Lu Z. The protein kinase activity of fructokinase A specifies the antioxidant responses of tumor cells by phosphorylating p62. *Sci Adv* 2019; 5(4): eaav4570
127. Ippolito L, Morandi A, Giannoni E, Chiarugi P. Lactate: a metabolic driver in the tumour landscape. *Trends Biochem Sci* 2019; 44(2): 153–166
128. Zhao D, Zou SW, Liu Y, Zhou X, Mo Y, Wang P, Xu YH, Dong B, Xiong Y, Lei QY, Guan KL. Lysine-5 acetylation negatively regulates lactate dehydrogenase A and is decreased in pancreatic cancer. *Cancer Cell* 2013; 23(4): 464–476
129. Fantin VR, St-Pierre J, Leder P. Attenuation of LDH-A expression uncovers a link between glycolysis, mitochondrial physiology, and tumor maintenance. *Cancer Cell* 2006; 9(6): 425–434
130. Rosalki SB, Wilkinson JH. Reduction of α -ketobutyrate by human serum. *Nature* 1960; 188(4756): 1110–1111
131. Liu Y, Guo JZ, Liu Y, Wang K, Ding W, Wang H, Liu X, Zhou S, Lu XC, Yang HB, Xu C, Gao W, Zhou L, Wang YP, Hu W, Wei Y, Huang C, Lei QY. Nuclear lactate dehydrogenase A senses ROS to produce α -hydroxybutyrate for HPV-induced cervical tumor growth. *Nat Commun* 2018; 9(1): 4429
132. Intlekofer AM, Wang B, Liu H, Shah H, Carmona-Fontaine C,

- Rustenburg AS, Salah S, Gunner MR, Chodera JD, Cross JR, Thompson CB. L-2-Hydroxyglutarate production arises from noncanonical enzyme function at acidic pH. *Nat Chem Biol* 2017; 13(5): 494–500
133. Yan H, Parsons DW, Jin G, McLendon R, Rasheed BA, Yuan W, Kos I, Batinic-Haberle I, Jones S, Riggins GJ, Friedman H, Friedman A, Reardon D, Herndon J, Kinzler KW, Velculescu VE, Vogelstein B, Bigner DD. IDH1 and IDH2 mutations in gliomas. *N Engl J Med* 2009; 360(8): 765–773
134. Kang MR, Kim MS, Oh JE, Kim YR, Song SY, Seo SI, Lee JY, Yoo NJ, Lee SH. Mutational analysis of IDH1 codon 132 in glioblastomas and other common cancers. *Int J Cancer* 2009; 125(2): 353–355
135. Nobusawa S, Yokoo H. IDH1/2 mutations in gliomas. *Brain Nerve* 2011; 63(12): 1378–1386
136. Mardis ER, Ding L, Dooling DJ, Larson DE, McLellan MD, Chen K, Koboldt DC, Fulton RS, Delehaunty KD, McGrath SD, Fulton LA, Locke DP, Magrini VJ, Abbott RM, Vickery TL, Reed JS, Robinson JS, Wylie T, Smith SM, Carmichael L, Eldred JM, Harris CC, Walker J, Peck JB, Du F, Dukes AF, Sanderson GE, Brummett AM, Clark E, McMichael JF, Meyer RJ, Schindler JK, Pohl CS, Wallis JW, Shi X, Lin L, Schmidt H, Tang Y, Haipek C, Wiechert ME, Ivy JV, Kalicki J, Elliott G, Ries RE, Payton JE, Westervelt P, Tomasson MH, Watson MA, Baty J, Heath S, Shannon WD, Nagarajan R, Link DC, Walter MJ, Graubert TA, DiPersio JF, Wilson RK, Ley TJ. Recurring mutations found by sequencing an acute myeloid leukemia genome. *N Engl J Med* 2009; 361(11): 1058–1066
137. Borger DR, Tanabe KK, Fan KC, Lopez HU, Fantin VR, Straley KS, Schenkein DP, Hezel AF, Ancukiewicz M, Liebman HM, Kwak EL, Clark JW, Ryan DP, Deshpande V, Dias-Santagata D, Ellisen LW, Zhu AX, Iafrate AJ. Frequent mutation of isocitrate dehydrogenase (IDH)1 and IDH2 in cholangiocarcinoma identified through broad-based tumor genotyping. *Oncologist* 2012; 17(1): 72–79
138. Yang H, Ye D, Guan KL, Xiong Y. IDH1 and IDH2 mutations in tumorigenesis: mechanistic insights and clinical perspectives. *Clin Cancer Res* 2012; 18(20): 5562–5571
139. Ward PS, Patel J, Wise DR, Abdel-Wahab O, Bennett BD, Collier HA, Cross JR, Fantin VR, Hedvat CV, Perl AE, Rabinowitz JD, Carroll M, Su SM, Sharp KA, Levine RL, Thompson CB. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting α -ketoglutarate to 2-hydroxyglutarate. *Cancer Cell* 2010; 17(3): 225–234
140. Reitman ZJ, Parsons DW, Yan H. IDH1 and IDH2: not your typical oncogenes. *Cancer Cell* 2010; 17(3): 215–216
141. Dang L, White DW, Gross S, Bennett BD, Bittinger MA, Driggers EM, Fantin VR, Jang HG, Jin S, Keenan MC, Marks KM, Prins RM, Ward PS, Yen KE, Liau LM, Rabinowitz JD, Cantley LC, Thompson CB, Vander Heiden MG, Su SM. Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature* 2009; 462(7274): 739–744
142. Xu W, Yang H, Liu Y, Yang Y, Wang P, Kim SH, Ito S, Yang C, Wang P, Xiao MT, Liu LX, Jiang WQ, Liu J, Zhang JY, Wang B, Frye S, Zhang Y, Xu YH, Lei QY, Guan KL, Zhao SM, Xiong Y. Oncometabolite 2-hydroxyglutarate is a competitive inhibitor of α -ketoglutarate-dependent dioxygenases. *Cancer Cell* 2011; 19(1): 17–30
143. Lu C, Ward PS, Kapoor GS, Rohle D, Turcan S, Abdel-Wahab O, Edwards CR, Khanin R, Figueroa ME, Melnick A, Wellen KE, O'Rourke DM, Berger SL, Chan TA, Levine RL, Mellinghoff IK, Thompson CB. IDH mutation impairs histone demethylation and results in a block to cell differentiation. *Nature* 2012; 483(7390): 474–478
144. Zhu H, Zhang Y, Chen J, Qiu J, Huang K, Wu M, Xia C. IDH1 R132H mutation enhances cell migration by activating AKT–mTOR signaling pathway, but sensitizes cells to 5-FU treatment as NADPH and GSH are reduced. *PLoS One* 2017; 12(1): e0169038
145. Parsons DW, Jones S, Zhang X, Lin JC, Leary RJ, Angenendt P, Mankoo P, Carter H, Siu IM, Gallia GL, Olivi A, McLendon R, Rasheed BA, Keir S, Nikolskaya T, Nikolsky Y, Busam DA, Tekleab H, Diaz LA Jr, Hartigan J, Smith DR, Strausberg RL, Marie SK, Shinjo SM, Yan H, Riggins GJ, Bigner DD, Karchin R, Papadopoulos N, Parmigiani G, Vogelstein B, Velculescu VE, Kinzler KW. An integrated genomic analysis of human glioblastoma multiforme. *Science* 2008; 321(5897): 1807–1812
146. Loenarz C, Schofield CJ. Expanding chemical biology of 2-oxoglutarate oxygenases. *Nat Chem Biol* 2008; 4(3): 152–156
147. Reiter-Brennan C, Semmler L, Klein A. The effects of 2-hydroxyglutarate on the tumorigenesis of gliomas. *Contemp Oncol (Pozn)* 2018; 22(4): 215–222
148. Wang P, Wu J, Ma S, Zhang L, Yao J, Hoadley KA, Wilkerson MD, Perou CM, Guan KL, Ye D, Xiong Y. Oncometabolite D-2-hydroxyglutarate inhibits ALKBH DNA repair enzymes and sensitizes IDH mutant cells to alkylating agents. *Cell Rep* 2015; 13(11): 2353–2361
149. Yogev O, Yogev O, Singer E, Shaulian E, Goldberg M, Fox TD, Pines O. Fumarase: a mitochondrial metabolic enzyme and a cytosolic/nuclear component of the DNA damage response. *PLoS Biol* 2010; 8(3): e1000328
150. Leshets M, Silas YBH, Lehming N, Pines O. Fumarase: from the TCA cycle to DNA damage response and tumor suppression. *Front Mol Biosci* 2018; 5: 68
151. Xiao M, Yang H, Xu W, Ma S, Lin H, Zhu H, Liu L, Liu Y, Yang C, Xu Y, Zhao S, Ye D, Xiong Y, Guan KL. Inhibition of α -KG-dependent histone and DNA demethylases by fumarate and succinate that are accumulated in mutations of FH and SDH tumor suppressors. *Genes Dev* 2012; 26(12): 1326–1338
152. Jiang Y, Qian X, Shen J, Wang Y, Li X, Liu R, Xia Y, Chen Q, Peng G, Lin SY, Lu Z. Local generation of fumarate promotes DNA repair through inhibition of histone H3 demethylation. *Nat Cell Biol* 2015; 17(9): 1158–1168
153. Wang T, Yu Q, Li J, Hu B, Zhao Q, Ma C, Huang W, Zhuo L, Fang H, Liao L, Eugene Chin Y, Jiang Y. O-GlcNAcylation of fumarase maintains tumour growth under glucose deficiency. *Nat Cell Biol* 2017; 19(7): 833–843
154. Lin R, Tao R, Gao X, Li T, Zhou X, Guan KL, Xiong Y, Lei QY. Acetylation stabilizes ATP-citrate lyase to promote lipid biosynthesis and tumor growth. *Mol Cell* 2013; 51(4): 506–518
155. Wellen KE, Hatzivassiliou G, Sachdeva UM, Bui TV, Cross JR, Thompson CB. ATP-citrate lyase links cellular metabolism to histone acetylation. *Science* 2009; 324(5930): 1076–1080
156. Lee JV, Carrer A, Shah S, Snyder NW, Wei S, Venneti S, Worth

- AJ, Yuan ZF, Lim HW, Liu S, Jackson E, Aiello NM, Haas NB, Rebbeck TR, Judkins A, Won KJ, Chodosh LA, Garcia BA, Stanger BZ, Feldman MD, Blair IA, Wellen KE. Akt-dependent metabolic reprogramming regulates tumor cell histone acetylation. *Cell Metab* 2014; 20(2): 306–319
157. Henderson B, Martin AC. Protein moonlighting: a new factor in biology and medicine. *Biochem Soc Trans* 2014; 42(6): 1671–1678
158. Palsson-McDermott EM, Dyck L, Zaslona Z, Menon D, McGettrick AF, Mills KHG, O'Neill LA. Pyruvate kinase M2 is required for the expression of the immune checkpoint PD-L1 in immune cells and tumors. *Front Immunol* 2017; 8: 1300