

Table S1. References containing published PKD/BFIE/ICCA pedigrees used for penetrance analysis

1. Wang JL, Cao L, Li XH, Hu ZM, Li JD, Zhang JG, Liang Y, San-A, Li N, Chen SQ, Guo JF, Jiang H, Shen L, Zheng L, Mao X, Yan WQ, Zhou Y, Shi YT, Ai SX, Dai MZ, Zhang P, Xia K, Chen SD, Tang BS. Identification of PRRT2 as the causative gene of paroxysmal kinesigenic dyskinesias. *Brain* 2011; 134: 3490–3498
2. Heron SE, Grinton BE, Kivity S, Afawi Z, Zuberi SM, Hughes JN, Pridmore C, Hodgson BL, Iona X, Sadleir LG, Pelekanos J, Herlenius E, Goldberg-Stern H, Bassan H, Haan E, Korczyn AD, Gardner AE, Corbett MA, Gécz J, Thomas PQ, Mulley JC, Berkovic SF, Scheffer IE, Dibbens LM. PRRT2 mutations cause benign familial infantile epilepsy and infantile convulsions with choreoathetosis syndrome. *Am J Hum Genet* 2012; 90(1): 152–160
3. Lee HY, Huang Y, Bruneau N, Roll P, Roberson ED, Hermann M, Quinn E, Maas J, Edwards R, Ashizawa T, Baykan B, Bhatia K, Bressman S, Bruno MK, Brunt ER, Caraballo R, Echenne B, Fejerman N, Frucht S, Gurnett CA, Hirsch E, Houlden H, Jankovic J, Lee WL, Lynch DR, Mohammed S, Müller U, Nespeca MP, Renner D, Rochette J, Rudolf G, Saiki S, Soong BW, Swoboda KJ, Tucker S, Wood N, Hanna M, Bowcock AM, Szepetowski P, Fu YH, Ptáček LJ. Mutations in the gene PRRT2 cause paroxysmal kinesigenic dyskinesia with infantile convulsions. *Cell Rep* 2012; 1(1): 2–12
4. Schubert J, Paravidino R, Becker F, Berger A, Bebek N, Bianchi A, Brockmann K, Capovilla G, Dalla Bernardina B, Fukuyama Y, Hoffmann GF, Jurkat-Rott K, Anttonen AK, Kurlemann G, Lehesjoki AE, Lehmann-Horn F, Mastrangelo M, Mause U, Müller S, Neubauer B, Püst B, Rating D, Robbiano A, Ruf S, Schroeder C, Seidel A, Specchio N, Stephani U, Striano P, Teichler J, Turkdogan D, Vigevano F, Viri M, Bauer P, Zara F, Lerche H, Weber YG. PRRT2 mutations are the major cause of benign familial infantile seizures. *Hum Mutat* 2012; 33(10): 1439–1443
5. Marini C, Conti V, Mei D, Battaglia D, Lettori D, Losito E, Bruccini G, Tortorella G, Guerrini R. PRRT2 mutations in familial infantile seizures, paroxysmal dyskinesia, and hemiplegic migraine. *Neurology* 2012; 79: 2109–2114
6. Cloarec R, Bruneau N, Rudolf G, Massacrier A, Salmi M, Bataillard M, Boulay C, Caraballo R, Fejerman N, Genton P, Hirsch E, Hunter A, Lesca G, Motte J, Roubertie A, Sanlaville D, Wong SW, Fu YH, Rochette J, Ptáček LJ, Szepetowski P. PRRT2 links infantile convulsions and paroxysmal dyskinesia with migraine. *Neurology* 2012; 79(21): 2097–2103
7. Gardiner AR, Bhatia KP, Stamelou M, Dale RC, Kurian MA, Schneider SA, Wali GM, Counihan T, Schapira AH, Spacey SD, Valente EM, Silveira-Moriyama L, Teive HA, Raskin S, Sander JW, Lees A, Warner T, Kullmann DM, Wood NW, Hanna M, Houlden H. PRRT2 gene mutations: From paroxysmal dyskinesia to episodic ataxia and hemiplegic migraine. *Neurology* 2012; 79(21): 2115–2121

8. Lee YC, Lee MJ, Yu HY, Chen C, Hsu CH, Lin KP, Liao KK, Chang MH, Liao YC, Soong BW. PRRT2 mutations in paroxysmal kinesigenic dyskinesia with infantile convulsions in a Taiwanese cohort. *PLoS ONE* 2012; 7(8): e38543. Doi: 10.1371/journal.pone.0038543.
9. Steinlein OK, Villain M, Korenke C. The PRRT2 mutation c.649dupC is the so far most frequent cause of benign familial infantile convulsions. *Seizure* 2012; 21(9): 740–742
10. Hedera P, Xiao J, Puschmann A, Momčilović D, Wu SW, LeDoux MS. Novel PRRT2 mutation in an African-American family with paroxysmal kinesigenic dyskinesia. *BMC Neurol* 2012; 12: 93
11. de Vries B, Callenbach PMC, Kamphorst JT, Weller CM, Koelewijn SC, ten Houten R, de Coo IFM, Brouwer OF, van den Maagdenberg AMJM. PRRT2 mutation causes benign familial infantile convulsions. *Neurology* 2012; 79(21): 2154–2155
12. Becker F, Schubert J, Striano P, Anttonen AK, Liukkonen E, Gaily E, Gerloff C, Müller S, Heußinger N, Kellinghaus C, Robbiano A, Polvi A, Zittel S, von Oertzen TJ, Rostasy K, Schöls L, Warner T, Münchau A, Lehesjoki AE, Zara F, Lerche H, Weber YG. PRRT2-related disorders: further PKD and ICCA cases and review of the literature. *J Neurol* 2013; 260(5): 1234–1244
13. Cai C, Shi O, Li WD. Missense mutations of the proline-rich transmembrane protein 2 gene cosegregate with mild paroxysmal kinesigenic dyskinesia and infantile convulsions in a Chinese pedigree. *Parkinsonism Relat Disord* 2013; 19(3): 402–403
14. Ishii A, Yasumoto S, Ihara Y, Inoue T, Fujita T, Nakamura N, Ohfu M, Yamashita Y, Takatsuka H, Taga T, Miyata R, Ito M, Tsuchiya H, Matsuoka T, Kitao T, Murakami K, Lee WT, Kaneko S, Hirose S. Genetic analysis of PRRT2 for benign infantile epilepsy, infantile convulsions with choreoathetosis syndrome, and benign convulsions with mild gastroenteritis. *Brain Dev* 2013; 35(6): 524–530
15. Okumura A, Shimojima K, Kubota T, Abe S, Yamashita S, Imai K, Okanishi T, Enoki H, Fukasawa T, Tanabe T, Dibbens LM, Shimizu T, Yamamoto T. PRRT2 mutation in Japanese children with benign infantile epilepsy. *Brain Dev* 2013; 35(7): 641–646
16. Labate A, Tarantino P, Palamara G, Gagliardi M, Cavalcanti F, Ferlazzo E, Sturniolo M, Incorpora G, Annesi G, Aguglia U, Gambardella A. Mutations in PRRT2 result in familial infantile seizures with heterogeneous phenotypes including febrile convulsions and probable SUDEP. *Epilepsy Res* 2013; 104(3): 280–284
17. Jing XY, Li XH, Yuan P, Deng J, Hu B, Wang Y. A novel mutation and functional implications of 5 variants in the PRRT2 gene in 20 paroxysmal kinesigenic dyskinesia pedigrees. *Parkinsonism Relat Disord* 2013; 19(6): 639–642
18. Wang JL, Mao X, Hu ZM, Li JD, Li N, Guo JF, Jiang H, Shen L, Li J, Shi YT, Xia K, Liu JY, Liao WP, Tang BS. Mutation analysis of PRRT2 in two Chinese BFIS families and nomenclature of PRRT2 related paroxysmal diseases. *Neurosci Lett* 2013; 552: 40–45
19. Yang X, Zhang Y, Xu X, Wang S, Yang Z, Wu Y, Liu X, Wu X. Phenotypes and PRRT2

- mutations in Chinese families with benign familial infantile epilepsy and infantile convulsions with paroxysmal choreoathetosis. *BMC Neurol* 2013; 13: 209
20. Djémié T, Weckhuysen S, Holmgren P, Hardies K, Van Dyck T, Hendrickx R, Schoonjans A-S, Van Paesschen W, Jansen AC, De Meirleir L, Selim LAM, Girgis MY, Buyse G, Lagae L, Smets K, Smouts I, Claeys KG, Van den Bergh V, Grisar T, Blatt I, Shorer Z, Roelens F, Afawi Z, Helbig I, Ceulemans B, De Jonghe P, Suls A. PRRT2 mutations: exploring the phenotypical boundaries. *J Neurol Neurosurg Psychiatry* 2014; 85(4): 462–465
 21. Brueckner F, Kohl B, Puest B, Gassner S, Osseforth J, Lindenau M, Stodieck S, Biskup S, Lohmann E. Unusual variability of PRRT2 linked phenotypes within a family. *Eur J Paediatr Neurol* 2014; 18(4): 540–542
 22. Ji Z, Su Q, Hu L, Yang Q, Liu C, Xiong J, Xiong F. Novel loss-of-function PRRT2 mutation causes paroxysmal kinesigenic dyskinesia in a Han Chinese family. *BMC Neurol* 2014; 14: 146
 23. Guerrero-López R, Ortega-Moreno L, Giráldez BG, Alarcón-Morcillo C, Sánchez-Martín G, Nieto-Barrera M, Gutiérrez-Delicado E, Gómez-Garre P, Martínez-Bermejo A, García-Peñas JJ, Serratos JM. Atypical course in individuals from Spanish families with benign familial infantile seizures and mutations in the PRRT2 gene. *Epilepsy Res* 2014; 108(8): 1274–1278
 24. Zhang LM, An Y, Pan G, Ding YF, Zhou YF, Yao YH, Wu BL, Zhou SZ. Reduced Penetrance of PRRT2 Mutation in a Chinese Family With Infantile Convulsion and Choreoathetosis Syndrome. *J Child Neurol* 2015; 30(10): 1263–1269
 25. Chen GH. Five cases of paroxysmal kinesigenic dyskinesia by genetic diagnosis. *Exp Ther Med* 2015; 9(3): 909–912
 26. Zhao G, Liu X, Zhang Q, Wang K. PRRT2 mutations in a cohort of Chinese families with paroxysmal kinesigenic dyskinesia and genotype-phenotype correlation reanalysis in literatures. *Int J Neurosci* 2018; 128(8): 751–760
 27. Ma H, Feng S, Deng X, Wang L, Zeng S, Wang C, Ma X, Sun H, Chen R, Du S, Mao J, Zhang X, Ma C, Jiang H, Zhang L, Tang B, Liu JY. A PRRT2 variant in a Chinese family with paroxysmal kinesigenic dyskinesia and benign familial infantile seizures results in loss of interaction with STX1B. *Epilepsia* 2018; 59(8): 1621–1630
 28. Yang L, You C, Qiu S, Yang X, Li Y, Liu F, Zhang D, Niu Y, Xu L, Xu N, Li X, Luo F, Yang J, Li B. Novel and de novo point and large microdeletion mutation in PRRT2-related epilepsy. *Brain Behav* 2020; 10(5): e01597. Doi: 10.1002/brb3.1597
 29. Baldi S, Zhu JL, Hu QY, Wang JL, Zhang JB, Zhang SH. A Novel PRRT2 Variant in Chinese Patients Suffering from Paroxysmal Kinesigenic Dyskinesia with Infantile Convulsion. *Behav Neurol* 2020; 2020: 2097059