

# Carbon dots based on targeting unit inheritance strategy for Golgi apparatus-targeting imaging

Yingying Wei (✉)<sup>1</sup>, Yudian Gao (✉)<sup>2,3</sup>, Lin Chen<sup>4</sup>, Qiang Li<sup>1</sup>, Jinglei Du<sup>1</sup>, Dongming Wang<sup>5</sup>, Fanggang Ren<sup>6</sup>, Xuguang Liu<sup>7</sup>, and Yongzhen Yang<sup>4</sup>

1 Interventional Treatment Department, Second Hospital of Shanxi Medical University (Second Clinical Medical College of Shanxi Medical University), Taiyuan 030001, China

2 Ophthalmology Department, Shanxi Bethune Hospital, Shanxi Academy of Medical Sciences, Tongji Shanxi Hospital, Third Hospital of Shanxi Medical University, Taiyuan 030032, China

3 Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China

4 Key Laboratory of Interface Science and Engineering in Advanced Materials (Ministry of Education), Taiyuan University of Technology, Taiyuan 030024, China

5 Key Laboratory of Molecular Diagnosis and Treatment of Blood Diseases of Shanxi Province, Second Hospital of Shanxi Medical University, Taiyuan 030001, China

6 Orthopedics Department, Second Hospital of Shanxi Medical University, Taiyuan 030001, China

7 Institute of Materials Science and Engineering, Taiyuan University of Technology, Taiyuan 030024, China

E-mails: weiyinying0726@126.com (Y.W.), gaoyudian@126.com (Y.G.)

## Supplementary material




Fig. S1 CD spectra of carbon dots synthesized at 140 °C for 5 h and 160 °C for 3 h.