

Recent results on nucleon spin structure study at RHIC

Qinghua Xu

*School of Physics, Key Laboratory of Particle Physics and Particle Irradiation (MoE), Shandong University,
Jinan 250100, China*

E-mail: xugh@sdu.edu.cn

Received August 3, 2015; accepted August 22, 2015

Both the PHENIX and STAR experiments at the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory are running polarized proton–proton collisions at $\sqrt{s} = 200$ and 500 GeV. The main goal of the RHIC spin physics program is to gain deeper insight into the spin structure of the nucleon. We will give an overview of recent spin results from RHIC, particularly the study of gluon polarization via jet/hadron production and sea quark polarization via W boson production in longitudinally polarized proton–proton collisions.

Keywords RHIC, nucleon spin structure, gluon polarization, sea quark polarization

PACS numbers 14.20.Dh, 13.88.+e, 13.87.Ce, 13.38.Be

Contents

1	Introduction	1
2	Gluon polarization (ΔG) measurements	2
3	Sea quark polarization measurements	4
4	Summary and outlook	6
	Acknowledgements	6
	References	6

experiments with polarized lepton and/or hadron beams have shown that the fraction of proton spin carried by the quark and antiquark spins is relatively small, with a typical value of about one-quarter (0.25) [1]. Inclusive DIS measurements, however, provided little constraint on the gluon polarization. The current best probes of Δg are offered by polarized proton–proton collisions, i.e., the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Laboratory, which is the first and currently the only high-energy polarized hadron–hadron collider in the world. On the other hand, the polarized proton–proton collider can provide a cleaner probe of the flavor separation of quark and antiquark polarization via parity-violating weak boson production than the semi-inclusive DIS (SIDIS) process, which suffers from uncertainty of the fragmentation functions.

Polarized proton–proton collisions at center-of-mass energies of 200 and 500 GeV at RHIC provide a unique way to gain new insights into the proton spin structure, especially the gluon polarization and sea quark polarization in the nucleon, as mentioned above [2]. There are two main experiments (detectors) for the spin program at RHIC: PHENIX and STAR. Both are carrying out a rich spin program using longitudinally polarized and transversely polarized proton–proton collisions.

For precise determination of the gluon polarization (Δg), the STAR detector is well suited for the reconstruction of various final states including jets, identified hadrons (π^0 , π^\pm), electrons/positrons, and photons, which allows for measurements of different gluon-

1 Introduction

The spin structure of the nucleon has been an hot topic in quantum chromodynamics (QCD) since the late 1980s, when polarized lepton–nucleon deep inelastic scattering (DIS) experiments revealed that only a small portion ($\sim 20\%$) of the nucleon spin is carried by the spins of quarks and antiquarks [1]. According to the spin sum rule in the light cone gauge, the proton spin of one-half can be decomposed into

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L_q + L_g, \quad (1)$$

where $\Delta\Sigma$ denotes the combined quark and antiquark spin contribution, and $L_{q(g)}$ is the quark (gluon) orbital angular momentum contribution. In the past three decades, significant efforts have been made both experimentally and theoretically to elucidate the contribution of the gluon spin and the flavor-separated contribution of quarks and antiquarks, as well as the contribution of the orbital angular momentum of quarks and gluons. DIS

*Special Topic: Spin Physics (Eds. Haiyan Gao & Bo-Qiang Ma).

involved processes. The recent STAR results on the inclusive jet double-spin asymmetry provide the first evidence of a nonzero gluon polarization in the Bjorken- x range $x > 0.05$, and the magnitude is even comparable to the total quark spin contribution, as indicated in the recent global analysis.

Another main objective of the STAR spin program is flavor separation of the quark and antiquark spin to the proton spin via W boson production in polarized proton-proton collisions at $\sqrt{s} = 500$ GeV. The parity-violating nature of the W boson provides a unique and clean way of measuring the anti- u and anti- d quark polarization without the involvement of hadronization as in the SIDIS process. The recent results of W boson single-spin asymmetry from the 2012 data at STAR, which provide new constraints on light sea quark polarization, will be discussed.

2 Gluon polarization (ΔG) measurements

In proton-proton collisions, the gluon polarization can be accessed with strongly interacting probes including jet or hadron production by measuring the double-spin asymmetries (A_{LL}),

$$A_{LL} = \frac{\sigma^{++} - \sigma^{+-}}{\sigma^{++} + \sigma^{+-}} = \frac{1}{P_1 P_2} \frac{N^{++} - RN^{+-}}{N^{++} + RN^{+-}}, \quad (2)$$

where $\sigma_{++/+-}$ denotes the cross section for jet/hadron production with proton beam helicity of “ $++/+-$ ”, P_1 and P_2 are the beam polarizations, $R = \mathcal{L}^{++}/\mathcal{L}^{+-}$ is the ratio of the luminosities, and N^{++} (N^{+-}) are the jet or hadron yields for equal (opposite) helicity beams. To see how A_{LL} provides sensitivity to the gluon polarization, at the leading order the asymmetry can be described as

$$A_{LL} = \frac{\sum_{f_a, f_b} \Delta f_a \times \Delta f_b \times d\hat{\sigma}^{f_a f_b \rightarrow fX} \cdot \hat{a}_{LL}^{f_a f_b \rightarrow fX}}{\sum_{f_a, f_b} f_a \times f_b \times d\hat{\sigma}^{f_a f_b \rightarrow fX}}, \quad (3)$$

where $\Delta f_{a(b)}$ denotes the helicity distribution function for parton $a(b)$, $f_{a(b)}$ is the unpolarized parton distribution function, and $\hat{a}_{LL}^{f_a f_b \rightarrow fX}$ is the double-spin asymmetry for the partonic process $f_a f_b \rightarrow fX$. The summation considers all the subprocesses, including qq , qg , and gg scattering. For most RHIC kinematics, qg and gg scatterings dominate the jet production [3, 4], making A_{LL} measurements for jets and hadrons sensitive to the gluon polarization, as shown in Fig. 1 for the fractional contributions of different subprocesses to the inclusive jet production in pp collisions at 200 and 510 GeV [5]. The measured A_{LL} versus jet transverse momenta cover




Fig. 1 Relative contributions from different partonic subprocesses to inclusive jet production as a function of $x_T (= 2p_T/\sqrt{s})$ at mid-rapidity at 200 and 510 GeV in pp collisions [5].

a wide range of the parton momentum fraction x of the proton, with a combination of different subprocesses. Ultimately, the data will be included in the global analysis, together with data on other processes such as DIS, to constrain the gluon polarization as different groups have done [6–10], similar to the unpolarized case.

At STAR, jets were reconstructed from charged tracks with $|\eta| < 1.3$ measured in the Time Projection Chamber (TPC) and energy deposits in the Barrel Electromagnetic Calorimeter (BEMC) with towers at $|\eta| < 1$ and the End-cap Electromagnetic Calorimeter (EEMC) at $1 < \eta < 2$. The jets were reconstructed using the midpoint cone algorithm [11] for early data and then the anti- k_t algorithm [12] (known as infrared-safe) was chosen as the jet algorithm at STAR since run 9 data. The jet cross section measurement and the first A_{LL} measurements of inclusive jets were done using 2003 and 2004 data at STAR [13], and inclusive jet production can be well described by pQCD. Updated measurements of A_{LL} with higher precision using 2005 and 2006 data have also been published [14, 15]. STAR also measured the double-spin asymmetries for π^0 at $|\eta| < 1$ [16] and the near-forward rapidity at $1 < \eta < 2$ [17]. The early A_{LL} results were included in the DSSV2008 QCD global analysis to provide information on the gluon polarization and ruled out large positive and negative scenarios with a small value of Δg in the range of $0.05 < x < 0.2$, resulting in a small gluon polarization within the uncertainties [7].

After the 2006 run, several improvements at STAR, including an increase in the jet trigger efficiency and a faster data acquisition system, enabled us to improve the A_{LL} precision significantly during the 2009 run. Figure 2 shows the A_{LL} results for inclusive jet production versus jet p_T in two pseudo-rapidity bins for the 2009 data [18] at STAR; theoretical predictions with inputs from different global analyses are also shown for comparison. These predictions were made by inserting the polarized




Fig. 2 Longitudinal double-spin asymmetry A_{LL} for inclusive jet production vs. jet p_T for $|\eta| < 1$ in pp collisions at 200 GeV at STAR [18].

parton distribution functions (PDFs) from BB [6], DSSV [7, 8], LSS [9], and NNPDF [10] into the NLO jet production code of Mukherjee and Vogelsang [4]. The BB10 and NNPDF polarized PDFs are based only on inclusive DIS data, whereas that of LSS includes both inclusive and SIDIS data sets. The LSS10 gluon density has a node at $x \simeq 0.2$, and the LSS10p gluon is positive definite at the input scale $Q_0^2 = 2.5 \text{ GeV}^2$. The DSSV [7] model, in addition to DIS and SIDIS data, also incorporates RHIC polarized proton–proton data before 2009. LSS10p provides the best description of these STAR jet data, whereas the STAR results are higher than the predictions of DSSV and NNPDF and below the predictions of BB10. However, the measurements fall within the combined data and model uncertainties for these three cases.

The impact of new STAR jet data on constraining the gluon polarization was studied using the reweighing method by the NNPDF group [19] and the truncated x -integral of $\Delta g(x, Q^2 = 10 \text{ GeV}^2) = 0.17 \pm 0.06$ in the range $0.05 < x < 0.2$, which was 0.05 ± 0.15 , as a comparison before having the 2009 STAR jet data. The STAR jet A_{LL} data are found to be consistent with the PHENIX π^0 A_{LL} values [20], as shown in Fig. 3, which are very small or consistent with zero within the uncertainty but cover a different scale range. The DSSV group also updated their global analysis by incorporating the

2009 A_{LL} results from the STAR inclusive jet channel and PHENIX π^0 channel, which suggested a consistent sizable value of the gluon polarization at the level of $\int_{0.05}^1 \Delta g dx (Q^2 = 10 \text{ GeV}^2) = 0.20_{-0.07}^{+0.06}$ [21], which is of similar magnitude as the total quark spin contribution itself.

In addition, STAR recently released new results on jet A_{LL} in pp collisions at 510 GeV, which cover a smaller x range and provide new constraints on the gluon polarization at low x . Figure 4 shows A_{LL} versus x_T for inclusive jet production at mid-rapidity at 510 GeV in pp collisions from 2012 STAR data [5], together with the results at 200 GeV, in comparison with theoretical predictions.

Di-jet production will allow a better constraint on the partonic kinematics to determine the shape of the gluon polarization. The partonic Bjorken- x of the incoming partons can be extracted at leading order from the




Fig. 3 Longitudinal double-spin asymmetry A_{LL} for π^0 production vs. p_T for $|\eta| < 1$ in pp collisions at 200 GeV at PHENIX [20].




Fig. 4 Inclusive jet A_{LL} vs. $x_T (= 2p_T/\sqrt{s})$ at mid-rapidity at 510 and 200 GeV in pp collisions at STAR, compared to recent NLO predictions. Red (blue) data points and curves represent 200 (510) GeV.

pseudo-rapidity plus the transverse momentum of the di-jets. The wide acceptance of the STAR experiment permits reconstruction of di-jet events with different topological configurations. Future inclusive and di-jet A_{LL} measurements at both $\sqrt{s} = 200$ and 500 GeV at STAR will extend the range toward smaller values of Bjorken- x .

3 Sea quark polarization measurements

While the valence quark helicity distributions are well determined at intermediate x from DIS experiments, the polarization of the sea quarks \bar{u} and \bar{d} is not well constrained by SIDIS data, especially regarding whether the light sea symmetry is broken as in the unpolarized case [22]. The production of $W^{+(-)}$ bosons in pp collisions with beams longitudinally polarized provides a perfect tool to access sea quark and valence quark helicity distributions without the need for hadron fragmentation functions. As W production is a parity-violating process, W bosons couple only to either left-handed particles or right-handed antiparticles, thus selecting only the corresponding helicity-positive or -negative quark and antiquark from the polarized proton. The longitudinal single-spin asymmetry A_L for W boson production with one proton beam polarized is defined as

$$A_L(W) \equiv \frac{\sigma(p_+p \rightarrow W) - \sigma(p_-p \rightarrow W)}{\sigma(p_+p \rightarrow W) + \sigma(p_-p \rightarrow W)}, \quad (4)$$

where σ_+/σ_- is the cross section with a helicity positive/negative proton beam.

As the production of W bosons violates parity maximally, the above A_L equation for W^\pm can be rewritten as follows at leading order:

$$A_L^{W^+} \equiv \frac{\Delta u(x_1)\bar{d}(x_2) - \Delta\bar{d}(x_1)u(x_2)}{u(x_1)\bar{d}(x_2) + \bar{d}(x_1)u(x_2)}, \quad (5)$$

$$A_L^{W^-} \equiv \frac{\Delta d(x_1)\bar{u}(x_2) - \Delta\bar{u}(x_1)d(x_2)}{d(x_1)\bar{u}(x_2) + \bar{u}(x_1)d(x_2)}. \quad (6)$$

We can see that $A_L^{W^+}$ approaches $\Delta u/u$ in the limit $y_W \gg 0$, as the rapidity along the polarized proton beam is defined as positive, and the valence quark usually carries a larger momentum fraction. Similarly, $A_L^{W^+}$ approaches $-\Delta\bar{d}/\bar{d}$ when $y_W \ll 0$. Thus, by identifying the rapidity of the W bosons, we can measure the polarizations of quark and antiquarks directly. For W^- , $\Delta\bar{u}/\bar{u}$ can be accessed in the backward region accordingly.

In experiments, W^\pm is usually detected via the leptonic channel with the kinetics of lepton decay, as the missing energy of the neutrino is hard to measure. The theoretical framework that describes inclusive lepton

production of W bosons has been developed, so the measurements of A_L versus the lepton rapidity can be compared with theoretical predictions [27, 28]. The first measurements of the spin asymmetry, A_L , and cross section for W^\pm production were reported by the STAR [23, 25] and PHENIX [24] experiments using data collected in 2009 at $\sqrt{s} = 500$ GeV. These measurements paved the way for further measurements with larger statistics and more pseudo-rapidity bins that will provide real constraints on \bar{u} and \bar{d} polarization in the nucleon.

At STAR, $W^{+(-)}$ are detected through $e^{+(-)}$ with large transverse momentum via the $W \rightarrow e\nu$ channel with the BEMC and EEMC. The $W \rightarrow e\nu$ events are characterized by an isolated e^\pm with a sizable transverse energy, E_T^e , that peaks near half the W mass (~ 40 GeV, referred to as the Jacobian peak). The neutrino from W decay, close to opposite in the azimuthal of decay e^\pm , with a large missing transverse energy, is undetected, which causes a large imbalance in the p_T sum of all the reconstructed final state particles. A p_T -balance variable is defined in terms of the p_T sum of the e^\pm candidate and the reconstructed jets outside an isolation cone around the electron track with a radius $\Delta R = 0.7$. This signed p_T -balance must be larger than a typical value of 14 GeV/c for the 2011 and 2012 data. Charge separation of W^+ and W^- is implemented on the basis of the e^\pm track curvature measured in the TPC. The W^\pm yield versus the e^\pm transverse energy E_T is shown in Fig. 5.

In 2012, the STAR experiment collected pp data with an integrated luminosity of 72 pb $^{-1}$ at $\sqrt{s} = 510$ GeV and an average beam polarization of 56%, which is six times larger than that in the 2009 dataset, thus allowing a pseudo-rapidity-dependent measurement of A_L for W bosons. In 2011, STAR also collected a relatively small data sample with an integrated luminosity of 11 pb $^{-1}$ with a similar beam polarization. The A_L results from the STAR 2011+2012 data [26] are shown in Fig. 6 for both W^+ and W^- versus the lepton pseudo-rapidity, with comparisons to theoretical predictions from NLO (CHE) [27] and fully resummed (RHICBOS) [28] calculations based on the DSSV08 helicity distributions [7]. The $A_L^{W^+}$ data are in good agreement with the predictions using the DSSV08 global analysis. However, the measured $A_L^{W^-}$ asymmetries are systematically larger than the theoretical predictions using the DSSV08 analysis, especially for $\eta_{e^-} < -0.5$, which indicates that these data provide new constraints for $\Delta\bar{u}$ compared to those from previous data. The region of $\eta_{e^-} < -0.5$ in particular is expected to be sensitive to the \bar{u} quark helicity distribution; hence, the enhancement of $A_L^{W^-}$ there suggests a sizable positive \bar{u} quark polarization relative to that expected from the global analysis of polarized DIS




Fig. 5 E_T^e distribution for W^- (top row) and W^+ (bottom row) candidates (black), background contributions, and the sum of background and $W \rightarrow e + \nu_e$ Monte Carlo (MC) signals (red dashed) in pp collisions at STAR.




Fig. 6 Single-spin asymmetry A_L for $W^\pm + Z^0$ production as a function of η_e in pp collisions at 510 GeV at STAR using 2011+2012 data [26].

data. These results provide the first clear evidence that the flavor symmetry is also broken in the polarized light sea quark, where $\Delta\bar{u}$ is positive and $\Delta\bar{d}$ is negative. The very recent DSSV⁺⁺ global analysis including the new RHIC data, in particular the STAR 2012 W A_L results,

shows a significant shift from negative to positive for the best-fit value of $\Delta\bar{u}$ with an integral of $0.05 < x < 1$ [29]. A first unbiased global determination of polarized parton distribution by the NNPDF group [19], also confirmed the above conclusions, even though they did not include any SIDIS data.

The PHENIX experiment also published its measurements on the single-spin asymmetry A_L for W^\pm production as a function of η_e in pp collisions at 510 GeV using 2011, 2012, and 2013 data [30], which cover only the mid-rapidity range, and the A_L results are consistent with the STAR results mentioned above in the same rapidity range. RHIC-STAR completed a long successful longitudinally polarized $p + p$ run at $\sqrt{s} = 510$ GeV in 2013 with an integrated luminosity of more than 300 pb^{-1} ; the new results will be available very soon and will further constrain the flavor-separated antiquark contributions to the proton spin.

The longitudinal spin transfer D_{LL} of Λ and $\bar{\Lambda}$ hyperons in pp collisions is expected to be sensitive to the helicity distribution function of strange (anti)quarks and to the polarized fragmentation functions [31, 32]. A first proof-of-principle measurement of D_{LL} at mid-rapidity has been performed with STAR data taken in 2005 [33]. In 2009, an eightfold larger data sample was collected at STAR; the D_{LL} analysis of this data sample extended the hyperon transverse momentum with better precision. Figure 8 shows the obtained D_{LL} results for Λ and $\bar{\Lambda}$




Fig. 7 Single-spin asymmetry A_L for W^\pm production as a function of η_e in pp collisions at 510 GeV at PHENIX using 2011, 2012, and 2013 data [30].




Fig. 8 Longitudinal spin transfer D_{LL} for Λ and $\bar{\Lambda}$ in polarized proton–proton collisions at $\sqrt{s} = 200$ GeV for positive η vs. p_T . The curves correspond to D_{LL} predictions for $\Lambda + \bar{\Lambda}$ based on different models of the polarized fragmentation functions [31].

versus p_T for $0 < \eta < 1.2$ in pp collisions at 200 GeV [34].

The data extend out to $p_T = 6$ GeV/ c with a statistical uncertainty of about 0.04. The model evaluations of D_{LL} [31] for the combination of Λ and $\bar{\Lambda}$ are consistent with the data and span a range of values that is similar to the experimental uncertainties.

4 Summary and outlook

RHIC, currently the only polarized hadron–hadron collider in the world, continues its efforts to deepen our understanding of the nucleon spin structure. New results from 2009 data on the jet double-spin asymmetry A_{LL} at RHIC-STAR in polarized proton–proton collisions at 200 GeV suggest a sizable gluon spin contribution comparable to the quark spin contribution itself. Future jet measurements will double the available data set at 200

GeV and also allow for a high statistics measurement at 500 GeV to provide constraints on Δg at lower x . Recent measurement of the single-spin asymmetry A_L for W^\pm production from 2011+2012 data at STAR, with a data sample about six times larger than that used for the first W measurement, have demonstrated new constraints on the antiquark helicity distributions $\Delta \bar{u}$ and $\Delta \bar{d}$. Future measurements, in particular the much larger data sample taken in 2013 at RHIC, will further constrain the gluon polarization and the flavor-separated antiquark contributions to the proton spin.

Acknowledgements The work was supported by the National Natural Science Foundation of China, China, under Grant Nos. 11175106 and 11222551, and the Natural Science Foundation of Shandong Province, China, under Grant No. ZR2013JQ001.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

1. For a review, see, e.g., C. A. Aidala, S. D. Bass, D. Hasch, and G. K. Mallot, The spin structure of the nucleon, *Rev. Mod. Phys.* 85, 655 (2013), arXiv: 1209.2803 [hep-ph]
2. G. Bunce, N. Saito, J. Soffer, and W. Vogelsang, Prospects for spin physics at RHIC, *Annu. Rev. Nucl. Part. Sci.* 50, 525 (2000), arXiv: hep-ph/0007218
3. D. de Florian and W. Vogelsang, Resummed cross-section for jet production at hadron colliders, *Phys. Rev. D* 76, 074031 (2007), arXiv: 0704.1677 [hep-ph]
4. A. Mukherjee and W. Vogelsang, Jet production in (un)polarized pp collisions: Dependence on jet algorithm,

- Phys. Rev. D* 86, 094009 (2012), arXiv: 1209.1785 [hep-ph]
5. E. C. Aschenauer, et al., The RHIC SPIN program: Achievements and future opportunities, arXiv: 1501.01220 [nucl-ex]
 6. J. Blümlein and H. Böttcher, QCD analysis of polarized deep inelastic scattering data, *Nucl. Phys. B* 841, 205 (2010), arXiv: 1005.3113
 7. D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, Global analysis of helicity Parton densities and their uncertainties, *Phys. Rev. Lett.* 101, 072001 (2008), arXiv: 0804.0422 [hep-ph]
 8. D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, Extraction of spin-dependent Parton densities and their uncertainties, *Phys. Rev. D* 80, 034030 (2009), arXiv: 0904.3821
 9. E. Leader, A. V. Sidorov, and D. B. Stamenov, Determination of polarized PDFs from a QCD analysis of inclusive and semi-inclusive deep inelastic scattering data, *Phys. Rev. D* 82, 114018 (2010), arXiv: 1010.0574
 10. R. D. Ball, et al. (NNPDF Collaboration), Unbiased determination of polarized parton distributions and their uncertainties, *Nucl. Phys. B* 874, 36 (2013), arXiv: 1303.7236
 11. M. Cacciari, G. P. Salam, and G. Soyez, The anti-k(t) jet clustering algorithm, *JHEP* 0804, 063 (2008), arXiv: 0802.1189 [hep-ph]
 12. G. C. Blazey, J. R. Dittmann, S. D. Ellis, V. D. Elvira, K. Frame, S. Grinstein, R. Hirsosky, R. Piegaiia, H. Schellman, R. Snihur, V. Sorin, and D. Zeppenfeld, Run II jet physics: Proceedings of the Run II QCD and Weak Boson Physics Workshop, arXiv: hep-ex/0005012
 13. B. I. Abelev, et al. (STAR Collaboration), Longitudinal double-spin asymmetry and cross section for inclusive jet production in polarized proton collisions at $\sqrt{s} = 200$ GeV, *Phys. Rev. Lett.* 97, 252001 (2006)
 14. B. I. Abelev, et al. (STAR Collaboration), Longitudinal double-spin asymmetry for inclusive jet production in $p + p$ collisions at $\sqrt{s} = 200$ GeV, *Phys. Rev. Lett.* 100, 232003 (2008)
 15. L. Adamczyk, et al. (STAR Collaboration), Longitudinal and transverse spin asymmetries for inclusive jet production at mid-rapidity in polarized $p + p$ collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. D* 86, 032006 (2012)
 16. B. I. Abelev, et al. (STAR Collaboration), Longitudinal double-spin asymmetry and cross section for inclusive neutral pion production at midrapidity in polarized proton collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. D* 80, 111108 (2009)
 17. L. Adamczyk, et al. (STAR Collaboration), Neutral pion cross section and spin asymmetries at intermediate pseudorapidity in polarized proton collisions at $\sqrt{s} = 200$ GeV, *Phys. Rev. D* 89, 012001 (2014), arXiv: 1309.1800 [nucl-ex]
 18. L. Adamczyk, et al. (STAR Collaboration), Precision measurement of the longitudinal double-spin asymmetry for inclusive jet production in polarized proton collisions at $\sqrt{s} = 200$ GeV, *Phys. Rev. Lett.* 115, 092002 (2015), arXiv: 1405.5134 [hep-ex]
 19. E. R. Nocera, et al. (NNPDF Collaboration), A first unbiased global determination of polarized PDFs and their un-
- certainties, *Nucl. Phys. B* 887, 276 (2014), arXiv: 1406.5539 [hep-ph]
20. A. Adare, et al. (PHENIX Collaboration), Inclusive double-helicity asymmetries in neutral-pion and eta-meson production in $p + p$ collisions at $\sqrt{s} = 200$ GeV, *Phys. Rev. D* 90, 012007 (2014), arXiv: 1402.6296 [hep-ex]
 21. D. de Florian, R. Sassot, M. Stratmann, and W. Vogelsang, Evidence for polarization of gluons in the proton, *Phys. Rev. Lett.* 113, 012001 (2014), arXiv: 1404.4293 [hep-ph]
 22. W. C. Chang and J. C. Peng, Flavor structure of the nucleon sea, *Prog. Part. Nucl. Phys.* 79, 95 (2014), arXiv: 1406.1260 [hep-ph]
 23. M. Aggarwal, et al. (STAR Collaboration), Measurement of the parity-violating longitudinal single-spin asymmetry for W^\pm boson production in polarized proton-proton collisions at $\sqrt{s} = 500$ GeV, *Phys. Rev. Lett.* 106, 062002 (2011), arXiv: 1009.0326
 24. A. Adare, et al. (PHENIX Collaboration), Cross section and parity-violating spin asymmetries of W^\pm boson production in polarized $p + p$ collisions at $\sqrt{s} = 500$ GeV, *Phys. Rev. Lett.* 106, 062001 (2011)
 25. L. Adamczyk, et al. (STAR Collaboration), Measurement of the $W \rightarrow e\nu$ and $Z/\gamma^* \rightarrow e+e-$ production cross sections at mid-rapidity in proton-proton collisions at $\sqrt{s} = 500$ GeV, *Phys. Rev. D* 85, 092010 (2012)
 26. L. Adamczyk, et al. (STAR Collaboration), Measurement of longitudinal spin asymmetries for weak boson production in polarized proton-proton collisions at RHIC, *Phys. Rev. Lett.* 113, 072301 (2014), arXiv: 1404.6880 [nucl-ex]
 27. D. de Florian and W. Vogelsang, Helicity parton distributions from spin asymmetries in W-boson production at RHIC, *Phys. Rev. D* 81, 094020 (2010)
 28. P. M. Nadolsky and C. P. Yuan, Single-spin physics with weak bosons at RHIC, *Nucl. Phys. B* 666(1), 31 (2003)
 29. E. C. Aschenauer, A. Bazilevsky, K. Boyle, K. O. Eysler, R. Fatemi, C. Gagliardi, M. Grosse-Perdekamp, J. Lajoie, Zhongbo Kang, Y. Kovchegov, et al., The RHIC spin program: Achievements and future opportunities, arXiv: 1304.0079
 30. A. Adare, et al. (PHENIX Collaboration), Measurement of parity-violating spin asymmetries in W^\pm production at midrapidity in longitudinally polarized $p+p$ collisions, arXiv: 1504.07451 [hep-ex]
 31. D. de Florian, M. Stratmann, and W. Vogelsang, Polarized Λ -baryon production in pp collisions, *Phys. Rev. Lett.* 81, 530 (1998)
 32. Q. H. Xu, Z. T. Liang, and E. Sichteremann, Anti-Lambda polarization in high energy pp collisions with polarized beams, *Phys. Rev. D* 73, 077503 (2006)
 33. B. I. Abelev, et al. (STAR Collaboration), Longitudinal spin transfer to Λ and $\bar{\Lambda}$ hyperons in polarized proton-proton collisions at $\sqrt{s} = 200$ GeV, *Phys. Rev. D* 80, 111102 (2009)
 34. J. Deng, et al. (STAR Collaboration), Longitudinal spin transfer to Λ and $\bar{\Lambda}$ in polarized proton-proton collisions at $\sqrt{s} = 200$ GeV at STAR, *Physics of Particle and Nuclei* 45(1), 73 (2014)