

MINI-REVIEW

The delay time of gravitational wave – gamma-ray burst associations

Bing Zhang

Department of Physics and Astronomy, University of Nevada, Las Vegas, NV 89154, USA

E-mail: zhang@physics.unlv.edu

Received June 22, 2019; accepted July 2, 2019

The first gravitational wave (GW) – gamma-ray burst (GRB) association, GW170817/GRB 170817A, had an offset in time, with the GRB trigger time delayed by ~ 1.7 s with respect to the merger time of the GW signal. We generally discuss the astrophysical origin of the delay time, Δt , of GW-GRB associations within the context of compact binary coalescence (CBC) – short GRB (sGRB) associations and GW burst – long GRB (IGRB) associations. In general, the delay time should include three terms, the time to launch a clean (relativistic) jet, Δt_{jet} ; the time for the jet to break out from the surrounding medium, Δt_{bo} ; and the time for the jet to reach the energy dissipation and GRB emission site, Δt_{GRB} . For CBC-sGRB associations, Δt_{jet} and Δt_{bo} are correlated, and the final delay can be from 10 ms to a few seconds. For GWB-IGRB associations, Δt_{jet} and Δt_{bo} are independent. The latter is at least ~ 10 s, so that Δt of these associations is at least this long. For certain jet launching mechanisms of IGRBs, Δt can be minutes or even hours long due to the extended engine waiting time to launch a jet. We discuss the cases of GW170817/GRB 170817A and GW150914/GW150914-GBM within this theoretical framework and suggest that the delay times of future GW/GRB associations will shed light into the jet launching mechanisms of GRBs.

Keywords gravitational waves, gamma-ray bursts: general

Contents

1	Introduction	1
2	Delay time of GW-GRB associations	2
2.1	General consideration	2
2.2	Δt_{jet}	2
2.3	Δt_{bo}	3
2.4	Δt_{GRB}	4
2.5	Burst duration T_{burst}	4
3	Different GW-GRB association models	5
3.1	CBC-sGRB associations	5
3.2	GWB-IGRB associations	6
4	Case studies	7
4.1	GW170817/GRB 170817A association	7
4.2	GW150914/GW150914-GBM association	7
5	Summary and discussion	7
	Acknowledgements	8
	References	8

after the merger and lasted for ~ 2 s. It consists of two pulses [3, 4], each lasting for ~ 1 s. Earlier, a controversial γ -ray signal, GW150914-GBM, was claimed by the Fermi GBM team to follow the first black hole - black hole (BH-BH) merger GW event, GW150914, with a delay of ~ 0.4 s [5–7].

The LIGO/Virgo third observational run (O3) started on April 1, 2019 and will last for one year. It is highly expected that more GW-GRB associations will be detected. At least NS-NS mergers and NS-BH mergers with a mass ratio $q \equiv m_1/m_2 < 5$ ($m_1 > m_2$) are expected to produce sGRBs¹. If some BH-BH mergers can make GRBs, more robust cases than GW150914/GW150914-GBM should be identified. Finally, under certain conditions, core collapse events that make long GRBs (IGRBs) may have a strong enough GW signal to be detected as a GW burst (GWB) by LIGO/Virgo GW detectors (e.g., [9]). It is possible that GWB-IGRB associations may be detected in the future.

The delay time of a GRB with respect to the GW signal cannot only be used to constrain fundamental physics (e.g., [10, 11]), but also carries important information about GRB physics, including jet launching mechanism, jet breakout from the surrounding medium, jet dissipation, and GRB radiation mechanism. All these are closely

1 Introduction

The first neutron star – neutron star (NS-NS) merger gravitational wave source GW170817 [1] was followed by a short gamma-ray burst (GRB) 170817A [2–4]. The short GRB (sGRB) triggered the Fermi GBM at $\Delta t \sim 1.7$ s

¹NS-BH mergers with $q > 5$ would not produce a GRB since the NS would not be tidally disrupted before being swallowed by the BH as a whole (e.g., [8]).

*arXiv: 1905.00781v1.

related to the unknown composition of the GRB jet, which is subject to intense debate in the field of GRBs ([12] for a comprehensive discussion). The origin of the ~ 1.7 s delay in GW170817/GRB 170817A has been discussed in the literature (e.g., [4, 11, 13–16]). In this mini-review, we systematically investigate several physical processes that contribute to the observed time delay Δt (Section 2). This is discussed within the context of compact binary coalescence (CBC)-sGRB associations and GWB-lGRB associations (Section 3). The case studies for GW170817/GRB 170817A and GW150914/GRB 150914-GBM are presented in Section 4. The results are summarized in Section 5 with some discussion.

2 Delay time of GW-GRB associations

2.1 General consideration

We assume that both GWs and photons travel with speed of light and only discuss the astrophysical origin of Δt . For discussions on how to use Δt to constrain physics beyond the standard model, see Refs. [10, 11] and references therein.

In general, the observed delay time due to an astrophysical origin should consist of three terms (Fig. 1), i.e.,

$$\Delta t = (\Delta t_{\text{jet}} + \Delta t_{\text{bo}} + \Delta t_{\text{GRB}})(1 + z), \quad (1)$$

where Δt_{jet} is the time for the engine to launch a relativistic jet, Δt_{bo} is the time for the jet to penetrate through

and break out from the surrounding medium (the ejecta in the CBC scenario and the stellar envelope in the core collapse scenario), and Δt_{GRB} is the time after breakout for the jet to reach the energy dissipation radius where the observed γ -rays are emitted. The $(1 + z)$ factor is the cosmological time dilation factor, which we will ignore in the rest of the discussion. All three time intervals are measured in the rest frame of the Earth observer (with the $(1 + z)$ correction). Since the engine is at rest with respect to the observer (ignoring proper motion of both the source and Earth) and since the jet is propagating with a non-relativistic speed in the surrounding medium, Δt_{jet} and Δt_{bo} are also essentially the times $\Delta \hat{t}_{\text{jet}}$ and $\Delta \hat{t}_{\text{bo}}$ measured in the rest frame of the central engine, which we call the “lab frame”.

2.2 Δt_{jet}

The jet launching time Δt_{jet} depends on the type of the central engine and the jet launching mechanism of GRBs. In the literature, a GRB jet can be launched either through accretion (onto a BH or a NS) or a magnetic mechanism. The latter applies to an NS (magnetar) engine. A relativistic jet is launched either as magnetic bubbles generated from differential rotation of the NS or through magnetic dipole spindown of a rapidly spinning magnetar.

For an accreting central engine, Δt_{jet} can be decomposed into three terms:

$$\Delta t_{\text{jet,acc}} = \Delta t_{\text{wait}} + \Delta t_{\text{acc}} + \Delta t_{\text{clean}}, \quad (2)$$




Fig. 1 A cartoon picture of a generic GW-GRB association. There are four steps: (1) A CBC or a core collapse event makes a bright GW signal, either in the form of a “chirp” or a “GWB”. This marks the zero time point T_0 . The event produces a hyper-accreting BH or a rapidly spinning magnetar; (2) After Δt_{jet} , a clean jet is launched from the central engine; (3) After another time interval Δt_{bo} , the jet breaks out from the surrounding medium; (4) After another time interval Δt_{GRB} , the jet reaches the GRB radius where γ -rays are emitted. Depending on the jet composition, there could be three possible sites: R_{ph} , R_{IS} , and R_{ICMART} , which correspond to three different durations of Δt_{GRB} .

where Δt_{wait} is the waiting time for a specific accretion model to operate; Δt_{acc} is the timescale to form the accretion disk and to start accretion; and Δt_{clean} is the timescale to launch a relativistic jet since accretion starts, which requires that mass loading is low enough, i.e., the mass loading parameter

$$\mu_0 \equiv \eta \Gamma_0 (1 + \sigma_0) \equiv L_{\text{jet}}(t) / \dot{M}_{\text{jet}}(t) c^2 \gg 1, \quad (3)$$

where η is the dimensionless enthalpy of the jet at the central engine, $\Gamma_0 \sim 1$ is the initial Lorentz factor of the jet, and σ_0 is the ratio between the magnetic energy density and matter energy density (including internal energy) at the central engine. Here $L_{\text{jet}}(t)$ is the time-dependent jet luminosity, and $\dot{M}_{\text{jet}}(t)$ is the time-dependent mass loading rate in the jet.

For the magnetic engine model, the jet launching time can be decomposed as

$$\Delta t_{\text{jet,mag}} = \Delta t_{\text{wait}} + \Delta t_{\text{B}} + \Delta t_{\text{clean}}, \quad (4)$$

where Δt_{wait} is again the waiting time for a specific magnetic model to operate, Δt_{B} is the timescale to establish a strong magnetic field through differential rotation, and Δt_{clean} is again the timescale for the environment to become clean enough to launch a relativistic jet.

A few more words about Δt_{wait} : Since the GRB jet launching mechanism is not identified, different jet launching models make different assumptions. When we discuss a particular model, Δt_{wait} is defined as the waiting time when the conditions for that mechanism to operate are satisfied. For example, for a model invoking a hyper-accreting BH to launch a jet, Δt_{wait} is the waiting time for a BH to form, which is the lifetime of a hypermassive neutron star (HMNS) or a supramassive neutron star (SMNS) before collapsing. Within this model, no jet is launched if the central object is an NS. On the other hand, under the same physical condition but for the jet launching model invoking a differentially rotating NS, a jet is directly launched during the HMNS phase, so that $\Delta t_{\text{wait}} = 0$. For another example, in the magnetar model, if the jet launching mechanism is through magnetic spindown, the early brief accretion phase may be regarded as Δt_{wait} . On the other hand, in the model invoking a hyper-accreting NS, $\Delta t_{\text{wait}} = 0$ for the same physical condition. See Section 3 for a detailed discussion on different jet-launching models.

Several timescales are related to the dynamical timescale of the system

$$t_{\text{dyn}} = 2\pi \left(\frac{R^3}{GM} \right)^{1/2} \simeq 1.8 \text{ ms } R_{6.5}^{3/2} \left(\frac{M}{3M_{\odot}} \right)^{-1/2}, \quad (5)$$

where M and R are the mass (normalized to $3M_{\odot}$, where M_{\odot} is the solar mass) and radius of the central engine (convention $Q_n = Q/10^n$ in cgs units adopted throughout the paper). In the accretion model, the disk forms within t_{dyn} , and the accretion starts within the viscous timescale

$\sim \alpha^{-1} t_{\text{dyn}} = 10\alpha^{-1} t_{\text{dyn}}$, where $\alpha \sim 0.1$ is the dimensionless viscosity constant. As a result, $\Delta t_{\text{acc}} \sim n t_{\text{dyn}}$, where $n \sim 10$. Similarly, in the magnetar model, magnetic amplification also takes a few dynamical timescales, i.e., $\Delta t_{\text{B}} \sim n t_{\text{dyn}}$, where $n \sim$ a few.

In both models, the timescale for the jet to become clean, Δt_{clean} is defined by the degree of mass loading. For a new born, hot central engine (either the hot accretion disk or the central magnetar), the dominant mass-loading mechanism is through the neutrino wind, with the mass-loading rate of an unmagnetized flow defined by [17]

$$\dot{M}_{\nu} = (2.5 \times 10^{-5} M_{\odot}/\text{s}) L_{\nu,52}^{5/3} \left(\frac{\epsilon_{\nu}}{10 \text{ MeV}} \right)^{10/3} R_6^{5/3} \left(\frac{M}{3M_{\odot}} \right)^{-2}, \quad (6)$$

where L_{ν} is the $\nu/\bar{\nu}$ luminosity, and ϵ_{ν} is the typical energy of $\nu/\bar{\nu}$. A magnetized engine will suppress mass loading by limiting entry of protons into the jet [18]. A detailed treatment of mass loading for the black hole and magnetar central engines have been carried out by Lei *et al.* [18] and Metzger *et al.* [19], respectively.

2.3 Δt_{bo}

After a clean jet with $\mu_0 \gg 1$ is launched, it has to propagate through the dense medium surrounding the engine. For the case of CBC-sGRB associations, the surrounding medium is mostly the ejecta launched right before the merger. For the case of GWB-IGRB associations, the surrounding medium is the in-falling stellar envelope of the progenitor star.

In order to launch a successful jet, a critical value of the jet energy needs to be reached (e.g., [20]). In the following, we assume that such a condition is satisfied, which is justified by the observations of GRB 170817A that show evidence of a successful jet [21, 22]. Very generally, the jet breakout timescale can be written as

$$\Delta t_{\text{bo}} = \frac{R_{\text{out}} - R_{\text{in}}}{(\beta_{\text{jet,h}} - \beta_{\text{out}})c}, \quad (7)$$

where R_{out} and β_{out} are the radius and dimensionless velocity (in the rest frame of the central engine) of the outer boundary of the surrounding medium, R_{in} is the radius of the central engine where the jet is launched, and $\beta_{\text{jet,h}}$ is the dimensionless speed of the jet head propagating inside the medium. Notice that $\beta_{\text{jet,h}}$ is much smaller than the termination dimensionless speed of the jet, β_{jet} , due to the high density of the surrounding medium. Whereas $\beta_{\text{jet}} \sim 1$ for a relativistic jet, $\beta_{\text{jet,h}} = 0.1\beta_{\text{jet,h,-1}}$ is typically non-relativistic.

For the case of CBC-sGRB associations, the dynamical ejecta moves outward with a dimensionless speed $\beta_{\text{out}} = \beta_{\text{ej}} = 0.1\beta_{\text{ej,-1}}$. The outer boundary of the ejecta²⁾

²⁾Note that the ejecta has a velocity profile, with the outer boundary defined by the fastest layer in the ejecta. For an order-of-magnitude treatment, we adopt the average speed of the ejecta.

is defined as $R_{\text{out}} = R_{\text{ej}} = \beta_{\text{ej}} c (\Delta t_{\text{jet}} + \Delta t_{\text{tidal}})$, where Δt_{tidal} is the time interval between the epoch when the neutron star is tidally disrupted (i.e., the dynamical ejecta is launched) and the epoch of coalescence. This is typically of the order of milliseconds [23].

In order to break out the ejecta, the jet head needs to propagate faster than the ejecta. Let the jet head advance with a dimensionless speed $\beta'_{\text{jet,h}}$ in the ejecta frame. Its lab-frame dimensionless speed reads

$$\beta_{\text{jet,h}} = \frac{\beta'_{\text{jet,h}} + \beta_{\text{ej}}}{1 + \beta'_{\text{jet,h}} \beta_{\text{ej}}}. \quad (8)$$

Equation (7) can be then written as

$$\Delta t_{\text{bo}}(\text{CBC/sGRB}) \simeq \frac{\beta_{\text{ej}} (\Delta t_{\text{jet}} + \Delta t_{\text{tidal}}) - R_{\text{in}}/c}{\beta_{\text{jet,h}} - \beta_{\text{ej}}}, \quad (9)$$

When $\Delta t_{\text{jet}} \gg \Delta t_{\text{tidal}}$, one has $R_{\text{out}} \gg R_{\text{in}}$, so that $\Delta t_{\text{bo}} \simeq \frac{\beta_{\text{ej}}}{\beta_{\text{jet,h}} - \beta_{\text{ej}}} \Delta t_{\text{jet}} \simeq (\beta_{\text{ej}}/\beta'_{\text{jet,h}}) \Delta t_{\text{jet}} \propto \Delta t_{\text{jet}}$ (see also Geng *et al.* [24]). In this case, the first two terms of Eq. (1) are correlated with each other. Since different bursts likely have different β_{ej} and $\beta'_{\text{jet,h}}$, the positive correlation between Δt_{jet} and Δt_{bo} should have a broad scatter.

For the case of GWB-lGRB associations, R_{out} is the outer boundary of the stellar envelope, R_* , which may be regarded as a constant during jet propagation, even if it is slowly shrinking due to fallback. Since $R_* \gg R_{\text{in}}$, the breakout time in this case is

$$\Delta t_{\text{bo}}(\text{GWB/lGRB}) \simeq \frac{R_*}{\beta_{\text{jet,h}} c}. \quad (10)$$

2.4 Δt_{GRB}

Since the jet travels with a relativistic speed after breaking out the surrounding medium (the brief acceleration phase ignored), the observer-frame time $\Delta t_{\text{GRB}} = (1 - \beta \cos \theta) \Delta \hat{t}_{\text{GRB}}$, where $\Delta \hat{t}_{\text{GRB}}$ is the lab-frame duration for the jet to travel to the GRB emission radius, β is the Lorentz factor and dimensionless speed of the relativistic jet, and θ is the angle between the jet direction and the line of sight. Since $\Delta \hat{t}_{\text{GRB}} \simeq R_{\text{GRB}}/c$, one has

$$\Delta t_{\text{GRB}} \simeq (1 - \beta \cos \theta) \frac{R_{\text{GRB}}}{c} \simeq \frac{R_{\text{GRB}}}{\Gamma^2 c}. \quad (11)$$

The last approximation is valid when $\theta \sim (0-1/\Gamma)$ with Γ being the Lorentz factor along the line of sight, which is relevant for a relativistically moving outflow with the $1/\Gamma$ cone covering the line-of-sight. For GW170817/GRB 170817A, simple arguments have ruled out the scenario invoking a top-hat jet beaming away from the line of sight (e.g., [4, 13, 25]).

The GRB emission radius is not identified. In the literature, there are at least three sites that have been suggested to emit γ -rays, which are

- The photosphere radius (e.g., [26–28])

$$R_{\text{ph}} \simeq (6 \times 10^{12} \text{ cm}) L_{w,52} \Gamma_2^{-3}, \quad (12)$$

where L_w is the isotropic-equivalent “wind” luminosity in the line-of-sight direction. Here the low-enthalpy-regime ($\eta < \eta_*$ in the notation of Mészáros & Rees [26]) has been adopted, which is relevant for weak GRBs associated with CBCs at a large viewing angle, such as GW170817/GRB 170817A.

- The internal shock radius [29, 30]

$$R_{\text{IS}} = \Gamma^2 c \delta t \simeq (3 \times 10^{12} \text{ cm}) \Gamma_2^2 \delta t_{-2}, \quad (13)$$

where δt is the variability timescale in the lightcurve, which is typically 10s of milliseconds.

- The internal collision-induced magnetic reconnection and turbulence (ICMART) radius [31, 32]

$$R_{\text{ICMART}} = \Gamma^2 c t_{\text{pulse}} \simeq (3 \times 10^{14} \text{ cm}) \Gamma_2^2 t_{\text{pulse}}, \quad (14)$$

where t_{pulse} is the duration of the broad pulses in the GRB lightcurve, which is typically seconds.

Which radius is relevant for GRB emission depends on the composition of the jet. For a matter dominated fireball, a quasi-thermal emission from the photosphere and a synchrotron emission component from the internal shock are expected [26, 33–35]. For a Poynting flux dominated outflow, both the photosphere and internal shock emission components are suppressed, and the GRB emission site is at a large radius R_{ICMART} [31].

Putting three cases together, one has

$$\Delta t_{\text{GRB}} = \begin{cases} (200 \text{ s})(1 - \beta \cos \theta) L_{w,52} \Gamma_2^{-3}, & R_{\text{GRB}} = R_{\text{ph}} \\ (1 - \beta \cos \theta) \Gamma^2 \delta t, & R_{\text{GRB}} = R_{\text{IS}} \\ (1 - \beta \cos \theta) \Gamma^2 t_{\text{pulse}}, & R_{\text{GRB}} = R_{\text{ICMART}} \end{cases} \simeq \begin{cases} (20 \text{ ms}) L_{w,52} \Gamma_2^{-5}, & R_{\text{GRB}} = R_{\text{ph}}, \\ \delta t, & R_{\text{GRB}} = R_{\text{IS}}, \\ t_{\text{pulse}}, & R_{\text{GRB}} = R_{\text{ICMART}}, \end{cases} \quad (15)$$

where the second part of the equation makes the assumption that there is relativistic moving materials along the line of sight.

2.5 Burst duration T_{burst}

It is relevant to discuss the true duration, T_{burst} , of a GRB here. Note that observationally defined duration T_{90} is the lower limit of T_{burst} , since it is limited by the detector’s sensitivity.

GRBs usually show highly variable lightcurves, sometimes displaying multiple broad “pulses” with rapidly varying spikes superposed on top [36]. Some GRBs only have one or two broad pulses. For GRBs with multiple pulses, the total duration T_{burst} is defined by the duration of the central engine activity. The duration of a broad

pulse, t_{pulse} , has two different interpretations. For models that invoke a small emission radius (R_{ph} or R_{IS}), one has $R_{\text{GRB}}/(\Gamma^2 c) \ll t_{\text{pulse}}$, so t_{pulse} has to be interpreted as the duration of one episode of the central engine activity. The broad pulses may be attributed to the modulation at the central engine, e.g., the interaction between the jet and the stellar envelope in the long GRB model (e.g., [37]). The hard-to-soft evolution of the peak energy E_p across broad pulses posed a challenge to such an interpretation [38]. Within the ICMART model [31] or the large-radius internal shock model [39], t_{pulse} is interpreted as emission duration of one fluid shell as it expands in space, with the peak time defined as either the time when the shell reaches the maximum dissipation, or when the synchrotron spectral peak sweeps across the observational band [32, 40]. For bursts with one single broad pulse, the burst duration is defined by

$$T_{\text{burst}} = t_{\text{pulse}} \simeq \frac{R_{\text{GRB}}}{\Gamma^2 c}, \quad (16)$$

which is the same expression as Δt_{GRB} (Eq. (11)).

3 Different GW-GRB association models

With the above preparation, in the following we discuss the delay times for different GW-GRB association systems in different models. The results are summarized in Table 1.

3.1 CBC-sGRB associations

BH-NS mergers with a moderate mass ratio ($q < 5$, [8]) are expected to be associated with sGRBs. The central engine is a BH, and the jet launching mechanism is accretion. One has $\Delta t_{\text{wait}} = 0$ s. The dynamical timescale (Eq. (5)) is \sim ms, so Δt_{acc} is \sim 10 ms. A hyper-accreting black hole engine is considered clean, especially if energy

is tapped via a global magnetic field [18]. One can take $\Delta t_{\text{clean}} \sim 0$ s. Overall, one would expect $\Delta t_{\text{jet}} \sim 10$ ms. Assuming $\beta'_{\text{jet,h}} \sim 0.1$, $\beta_{\text{ej}} \sim 0.1$, and $R_{\text{in}} \sim 3$ times of the Schwarzschild radius, one gets $\Delta t_{\text{bo}} \sim \Delta t_{\text{jet}} + \Delta t_{\text{tidal}}$, which is also ~ 10 ms. Finally, Δt_{GRB} depends on jet composition and dissipation mechanism (Eq. (15)). For a $\nu\bar{\nu}$ -annihilation driven fireball, Δt_{GRB} is usually very short ($< \text{ms}$) (unless Γ is extremely low) thanks to the small emission radius. For a Poynting-flux-dominated jet, $\Delta t_{\text{GRB}} \sim t_{\text{pulse}}$ can be up to the duration of the sGRB itself.

NS-NS mergers are more complicated. Depending on the equation of state and the total mass in the merger, there could be four different outcomes (e.g., [41]): a promptly formed BH, a differential-rotation-supported hypermassive NS (HMNS) followed by collapse, a uniform-rotation-supported supramassive NS (SMNS) followed by collapse, or a stable NS (SNS). The prompt BH case is similar to the BH-NS merger case. For the HMNS case (which forms a BH within $\sim (0.1-1)$ s after the merger) and the SMNS/SNS case (which forms a long-lived NS), we consider both an accretion-powered jet and a magnetically powered jet, respectively.

The HMNS/BH accretion model assumes that the jet is launched when the BH is formed. This model introduces a waiting time $\Delta t_{\text{wait}} \sim \Delta t_{\text{HMNS}} \sim (0.1-1)$ s, where Δt_{HMNS} is the lifetime of the HMNS. The jet launching time Δt_{jet} is dominated by Δt_{wait} , and the jet breakout time Δt_{bo} scales up with Δt_{jet} proportionally. The third time Δt_{GRB} again can be from shorter than ms (fireball photosphere emission) to the duration of the burst itself (Poynting jet).

The HMNS/BH magnetic model assumes that a jet can be launched during the HMNS phase, so that $\Delta t_{\text{wait}} = 0$ s. The magnetic field amplification time Δt_{B} is again several times of t_{dyn} , i.e., ~ 10 ms. The “clean” time t_{clean} is quite uncertain. Rosswog *et al.* [42] performed magnetohydrodynamic (MHD) simulations of NS-NS mergers in

Table 1 Δt in different GW-GRB association scenarios.

System	Engine	Jet mechanism	Δt_{jet}			Δt_{bo}	Δt_{GRB}	Δt
			Δt_{wait}	$\Delta t_{\text{acc}}/\Delta t_{\text{B}}$	Δt_{clean}			
BH-NS	BH	accretion	~ 0 s	~ 10 ms	~ 0 s	(10–100) ms	$< \text{ms}$ to $\sim \text{s}$	(0.01–few) s
NS-NS	BH	accretion	~ 0 s	~ 10 ms	~ 0 s	(10–100) ms	$< \text{ms}$ to $\sim \text{s}$	(0.01–few) s
NS-NS	HMNS/BH	accretion	(0.1–1) s	~ 10 ms	~ 0 s	(0.1–1) s	$< \text{ms}$ to $\sim \text{s}$	(0.1–few) s
NS-NS	HMNS/BH	magnetic	~ 0 s	~ 10 ms	(0–1) s	(0.01–1) s	$< \text{ms}$ to $\sim \text{s}$	(0.01–few) s
NS-NS	SMNS/SNS	accretion	~ 0 s	~ 10 ms	(0–0.1) s	(10–100) ms	$< \text{ms}$ to $\sim \text{s}$	(0.01–few) s
NS-NS	SMNS/SNS	magnetic	~ 0 s	~ 10 ms	(0–10) s	(0.01–10) s	$< \text{ms}$ to $\sim \text{s}$	(0.01–10) s
Type I collapsar	BH	accretion	(0–several) s	~ 10 ms	~ 0 s	(10–50) s	$< \text{ms}$ to $\sim \text{s}$	(10–50) s
Type II collapsar	BH	accretion	(10^2-10^4) s	~ 10 ms	~ 0 s	(10–50) s	$< \text{ms}$ to $\sim \text{s}$	(10^2-10^4) s
core collapse	magnetar	accretion	~ 0 s	~ 10 ms	(0–10) s	(10–50) s	$< \text{ms}$ to $\sim \text{s}$	(10–50) s
core collapse	magnetar	magnetic	~ 0 s	~ 10 ms	(0–10) s	(10–50) s	$< \text{ms}$ to $\sim \text{s}$	(10–50) s
core collapse	magnetar	spindown	$(1-10^3)$ s	N/A	~ 10 s	(10–50) s	$< \text{ms}$ to $\sim \text{s}$	$(10-10^3)$ s

the HMNS phase and claimed that magnetic fields can be amplified to several times of 10^{17} G within $\Delta t_B \sim 10$ ms. They suggested that a relativistic short GRB jet can be launched along the axis of the binary orbit during this period of time, even though their numerical simulations were not able to resolve the baryon loading process. Within this scenario, Δt_{clean} may be as short as 0 s, so that Δt_{jet} is of the order of $\Delta t_B \sim 10$ ms. The breakout time Δt_{bo} is correspondingly short. On the other hand, a recent numerical simulation [43] suggested that no relativistic jet is launched before 100 ms due to strong baryon loading. If this is the case, Δt_{clean} may be at least ~ 0.1 s for this model. Both Δt_{jet} and Δt_{bo} are increased correspondingly. Again Δt_{GRB} is negligible in the fireball model and can be of the order of the burst duration in the Poynting jet model.

Finally, the SMNS/SNS models assume that such systems can power sGRBs [44–46]. The observational evidence of this class of models is the so-called “internal X-ray plateau” (a lightcurve plateau followed by a sharp drop of flux which can be only interpreted as “internal” dissipation of a central engine outflow) observed following a good fraction of sGRBs [47–49]. The plateau is best interpreted as internal dissipation of a post-merger massive magnetar wind [50–53] during the magnetic spindown phase [54]. The sGRB needs to be produced by the massive NS, likely shortly after the merger.

The SMNS/SNS accretion model assumes that a relativistic sGRB jet can be launched from a magnetar via hyper-accretion [46, 55]. Within this scenario, $\Delta t_{\text{wait}} = 0$ s, and accretion starts after $\Delta t_{\text{acc}} \sim 10$ ms. Baryon loading in this model has not been well-studied, and it is assumed that Δt_{clean} is short, e.g., 0–100 ms. As a result, both Δt_{jet} and Δt_{bo} are tens of ms. Δt_{GRB} is again from $< \text{ms}$ (fireball) to ~ 1 s (Poynting jet).

The SMNS/SNS magnetic model is similar to the HMNS/BH magnetic model, except that Δt_{clean} can be longer (no longer limited by Δt_{HMNS}). Within the most optimistic model suggested by Rosswog *et al.* [42], Δt_{clean} can be as short as ~ 0 s. On the other hand, according to the calculation of Metzger *et al.* [19], baryon mass loss rate in a proto-NS could be initially very high, e.g., $\sigma_0 < 1$ before ~ 2 s and $\sigma_0 < 100$ before ~ 10 s. Within this scenario, the outflow is initially non-relativistic, and a clean jet capable of producing a sGRB is launched only after $\Delta t_{\text{clean}} \sim (1\text{--}10)$ s. A longer Δt_{clean} leads to longer Δt_{jet} and Δt_{bo} , which can exceed Δt_{GRB} even for the Poynting jet case. The total delay time Δt could be dominated by Δt_{clean} .

3.2 GWB-IGRB associations

Within the core collapse model of long GRBs, the violent collapsing process may leave behind a central object with a significant quadrupole moment to generate a GW burst (e.g., [9]). For these systems, unlike CBCs, it is not

straightforward to define the epoch of significant GW radiation (i.e., the peak of GW burst signal). The strongest GW radiation is likely produced during the core collapse phase (e.g., [56]). The rapidly proto-NS (magnetar) may carry a significant quadrupole moment and radiate GW emission as well [54, 57, 58]. Finally, after the NS collapses to a BH, the neutrino-dominated accretion flow (NDAF) into the BH may also radiate GW, even if with a lower amplitude (e.g., [59]). In the following discussion, we assume that the GWB emission peaks at the core collapse time.

One can consider two general categories of models: core collapse events leading to a BH engine (which is usually called the “collapsar” model), and core collapse events leading to a magnetar engine (which we call the “magnetar” model). The progenitor of both types of engines can be either an isolated single star or a binary system whose merger induces the core collapse of the merged star.

Two types of collapsars have been discussed in the literature: Type I collapsar model [60, 61] invokes the collapse of the iron core of a rapidly rotating helium star, forming a short-lived NS that subsequently collapses in a few seconds. Type II collapsar model [62], on the other hand, invokes a long-lived NS, which continues to accrete fall-back materials for an extended period of time before collapsing to a BH minutes or even hours later. The progenitor stars of Type II collapsars are more common, so that these events may have a higher rate than Type I collapsars [62]. For either case, since a BH is required to launch a GRB jet, there is a waiting time Δt_{wait} that marks the duration of the proto-NS phase, ranging from several seconds (Type I collapsar) to hours (Type II collapsar). This term is likely the dominating term in Δt_{jet} . Unlike CBC-sGRB associations, the jet breakout time Δt_{bo} is independent of Δt_{jet} when Δt_{jet} is smaller than the free-fall timescale of a massive star $t_{\text{ff}} = [3\pi/(32G\bar{\rho})]^{1/2} \sim 180 \text{ s } (\bar{\rho}/100 \text{ g} \cdot \text{cm}^{-3})^{-1/2}$ ($\bar{\rho}$ is the mean density of the stellar envelope), and is set by the size of the progenitor and the jet head speed. The widely accepted progenitor system of lGRBs is Wolf–Rayet stars [63]. Taking $R_* \sim 10^{11}$ cm and $\beta_{\text{jet,h}} \sim 0.1$, one gives $\Delta t_{\text{bo}} \sim (10\text{--}50)$ s based on Eq. (10), which is longer than Δt_{GRB} . The final Δt is \sim tens of seconds for Type I collapsar and $\sim (10^2\text{--}10^4)$ s for Type II collapsar.

For the magnetar model, a long GRB may be produced via one of the following three mechanisms: accretion, magnetic due to differential rotation, and magnetic due to spindown. The three mechanisms differ mainly in Δt_{jet} . For the accretion mechanism (e.g., [55]) and magnetic mechanism (e.g., [64–66]), $\Delta t_{\text{wait}} \sim 0$ s. The jet launching time Δt_{jet} is mostly controlled by Δt_{clean} , which may range from milliseconds (for most optimistic scenarios, e.g., [64, 66]) to ~ 10 s [19]. In any case, Δt_{bo} would contribute significantly to the total Δt . For the spindown model [57, 67], the assumption is that a GRB jet is launched as the magnetar spins down. This would be

after the early accretion phase. As a result, one should introduce an early Δt_{wait} for the duration of the accretion phase, which would be typically $1\text{--}10^3$ s. This is based on the observed duration of long GRBs (which are interpreted as the accretion time for most models) and an estimate of the free-fall timescale of a massive star $t_{\text{ff}} \sim 180$ s, which is the minimum timescale for accretion. As a result, this model would have a longer $\Delta t \sim (10\text{--}10^3)$ s than other magnetar models, with the delay mostly contributed from Δt_{wait} .

4 Case studies

4.1 GW170817/GRB 170817A association

This is the only robust GW-GRB association case. The GRB trigger time is delayed by ~ 1.7 s with respect to the binary NS merger time [2]. The duration of GRB 170817A is ~ 2 s, with two pulses each lasting ~ 1 s [3, 4]. Since this was a relatively weak burst, the true duration of each component (t_{pulse}) could be longer than 1 s.

According to the theory discussed above, there are two possible interpretations to the ~ 1.7 s delay.

The first scenario invokes a matter/radiation-dominated fireball. Within this scenario, the sGRB is most likely emission from the photosphere. Using Eq. (15) and noticing $L \sim 10^{47}$ erg·s $^{-1}$ for GRB 170817A [3, 4], one has $\Delta t_{\text{GRB}} \sim (20 \text{ ms}) L_{47} \Gamma_1^{-5}$. Forcing $\Delta t_{\text{GRB}} \sim 2$ s, one requires very low Lorentz factor $\Gamma \sim 4$. This was the suggested ‘‘cocoon breakout model’’ (or mildly relativistic, wide angle outflow model) of GRB 170817A shortly after the discovery of the event (e.g., [68, 69]), which was later disfavored by the discovery of superluminal motion of the radio afterglow of the source [21, 22]. Modeling of radio afterglow of the source suggests that the outflow Lorentz factor decays with time as $\Gamma \sim 4(t/10 \text{ day})^{-0.29}$ [70], which means that at $t < 10$ d, one has $\Gamma \gg 4$. This suggests that within this scenario, one has $\Delta t_{\text{GRB}} \ll \Delta t$, and the delay should be attributed to Δt_{jet} and Δt_{bo} . One way to make a long Δt_{jet} is to introduce a long $\Delta t_{\text{wait}} \sim 1$ s, which is regarded as the duration of the HMNS phase. Such a waiting time was indeed introduced in some of the numerical simulations (e.g., [71, 72]), and was suggested from kilonova modeling as well [73, 74]. A long waiting time is also the necessary condition to make a significant cocoon emission component [24]. This scenario has to assume that no relativistic jet is launched during the HMNS phase [75, 76], in contrast to some previous sGRB models (e.g., [42]). The issue of this scenario is that one has to interpret the delay time $\Delta t \sim 1.7$ s and the duration of the burst $T_{90} \sim 2$ s using two different mechanisms: while Δt is mostly controlled by Δt_{wait} , T_{90} has to be defined by the duration of the central engine (accretion timescale). The similar values of the two timescales have to be explained as a coincidence.

The second scenario, as advocated by Zhang *et al.* [4],

attributes Δt mostly to Δt_{GRB} . This is motivated by the intriguing fact that $\Delta t \sim 1.7$ s and $T_{90} \sim 2$ s are comparable. Based on Eq. (15), for a Poynting-flux-dominated outflow, $\Delta t_{\text{GRB}} \sim t_{\text{pulse}}$. If one takes the first pulse only and considers the weak nature of GRB 170817A (the true pulse duration should be longer than what is observed), one has $\Delta t_{\text{GRB}} > 1$ s, which occupies most of the observed Δt . Within this scenario, both Δt_{jet} and Δt_{bo} are short (say, $\ll 0.5$ s), which suggests a negligible Δt_{wait} . Within this scenario, there is no need to introduce an HMNS. The engine could be a BH, an HMNS with a lifetime shorter than 100 ms, an SMNS or even an SNS. One prediction of such a scenario is that Δt should be correlated with the burst duration (if the bursts have 1–2 simple pulses like GRB 170817A). For example, if the next NS-NS-merger-associated sGRB has a shorter duration (e.g., 0.5 s), the delay time Δt should be also correspondingly shorter. A smaller Δt_{wait} also suggests a less significant cocoon emission, even though the outflow is still a structured jet [24].

4.2 GW150914/GW150914-GBM association

Since observationally the case is not robust, this association may not be physical. On the other hand, if the association is real, the delay timescale $\Delta t \sim 0.4$ s places great constraints on the proposed models.

Most proposed models to interpret GW150914-GBM invoke substantial matter around the merging site. Loeb [77] invoked two BHs formed during the collapse of a massive star. Accretion after the merger powers the putative GRB. Putting aside other criticisms to the model (e.g., [78, 79]), Δt in such core collapse model should be at least Δt_{bo} , which is 10s of seconds. The observed $\Delta t \sim 0.4$ s therefore essentially rules out the model, unless a contrived jet launching time is introduced [80]. The same applies to other models that invoke a massive star as the progenitor of the putative GRB (e.g., [81]). The reactivated accretion disk model [82] is not subject to this constraint. However, it is likely that the reactivation happens way before the merger itself [83].

The charged BH merger model (Zhang [84], see [85]) for a more general discussion of charged CBC signals and Dai [86] for related signals) does not invoke a matter envelope surrounding the merger system. Within the framework discussed in this paper, both Δt_{jet} and Δt_{bo} are ~ 0 s, and Δt is dominated by Δt_{GRB} (see [84] for detailed discussion). The difficulty of this model is the origin of the enormous charge needed to account for the GRB.

5 Summary and discussion

We have discussed various physical processes that may cause a time delay Δt of a GRB associated with a GW event. The conclusions can be summarized as follows:

- In general, there are three timescales, i.e., Δt_{jet} , Δt_{bo}

and Δt_{GRB} , that will contribute to the observed Δt . Since the GRB jet launching mechanism is poorly understood, different scenarios make different assumptions. The assumptions introduced in different scenarios are sometimes contradictory (e.g., regarding whether a BH is needed to launch a GRB jet). The results are summarized in Table 1. With the GW information (which sets the fiducial time), one can in principle test these scenarios with a sample of GW/GRB associations in the future.

- We considered both CBC-sGRB associations and GWB-IGRB associations. For the former, different models point towards a similar range of Δt : from 10 ms to a few seconds. The 1.7 s delay of GW170817/GRB 170817A falls into this range, so this duration alone cannot be used to diagnose the jet launching mechanism. On the other hand, a statistical sample of GW/GRB associations can be in principle used to test the two scenarios with and without a significant intrinsic central engine waiting time: If Δt of different events are independent of the duration of the sGRB and especially the duration of the GRB pulses t_{pulse} , then it would be likely controlled by Δt_{jet} , in particular, Δt_{wait} of the central engine. Such a Δt_{wait} may be attributed to the lifetime of an HMNS, and a BH engine is needed to power a sGRB. If, on the other hand, Δt is always roughly proportional to t_{pulse} , as is the case of GW170817/GRB 170817A association, then Δt is likely dominated by Δt_{GRB} , with negligible Δt_{jet} and Δt_{bo} . The jet composition in this case is likely Poynting-flux-dominated, and the launch of a GRB jet may not necessarily require the formation of a BH.
- The GWB-IGRB associations all should have a longer delay, with Δt at least defined by the jet propagation and breakout time Δt_{bo} , which is at least ~ 10 s. In some models, such as the Type II collapsar model and the magnetar spindown model, there could be an additional waiting time Δt_{wait} before the presumed jet launching mechanism starts to operate. In these cases, the delay can be as long as minutes to even hours. Detecting GWB-IGRB associations with such a long delay (e.g., $\gg 100$ s) would point towards these specific jet launching mechanisms or a progenitor star much larger in size than a Wolf-Rayet star.

Acknowledgements I thank Wei-Hua Lei, Robert Mochkovitch, and Bin-Bin Zhang for discussion on the origin of Δt of GW170817/GRB170817A association.

References

1. B. P. Abbott, R. Abbott, T. D. Abbott, F. Acernese, K. Ackley, et al., GW170817: Observation of gravitational waves from a binary neutron star inspiral, *Phys. Rev. Lett.* 119(16), 161101 (2017)
2. B. P. Abbott, R. Abbott, T. D. Abbott, F. Acernese, K. Ackley, et al., Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A, *Astrophys. J.* 848(2), L13 (2017)
3. A. Goldstein, P. Veres, E. Burns, M. S. Briggs, R. Hamburg, et al., An ordinary short gamma-ray burst with extraordinary implications: *Fermi*-GBM detection of GRB 170817A, *Astrophys. J.* 848(2), L14 (2017)
4. B. B. Zhang, B. Zhang, H. Sun, W. H. Lei, H. Gao, Y. Li, L. Shao, Y. Zhao, Y. D. Hu, H. J. Lü, X. F. Wu, X. L. Fan, G. Wang, A. J. Castro-Tirado, S. Zhang, B. Y. Yu, Y. Y. Cao, and E. W. Liang, A peculiar low-luminosity short gamma-ray burst from a double neutron star merger progenitor, *Nat. Commun.* 9(1), 447 (2018)
5. V. Connaughton, E. Burns, A. Goldstein, L. Blackburn, M. S. Briggs, et al., *Fermi* GBM observations of LIGO gravitational-wave event GW150914, *Astrophys. J.* 826(1), L6 (2016)
6. V. Connaughton, E. Burns, A. Goldstein, L. Blackburn, M. S. Briggs, et al., On the interpretation of the *Fermi*-GBM transient observed in coincidence with LIGO gravitational-wave event GW150914, *Astrophys. J.* 853(1), L9 (2018)
7. J. Greiner, J. M. Burgess, V. Savchenko, and H. F. Yu, On the *Fermi*-GBM event 0.4 s after GW150914, *Astrophys. J.* 827(2), L38 (2016)
8. M. Shibata, K. Kyutoku, T. Yamamoto, and K. Taniguchi, Gravitational waves from black hole-neutron star binaries: Classification of waveforms, *Phys. Rev. D* 79(4), 044030 (2009)
9. S. Kobayashi and P. Mészáros, Gravitational radiation from gamma-ray burst progenitors, *Astrophys. J.* 589(2), 861 (2003)
10. J.-J. Wei, B.-B. Zhang, X.-F. Wu, H. Gao, P. Mészáros, B. Zhang, Z.-G. Dai, S.-N. Zhang, and Z.-H. Zhu, Multimessenger tests of the weak equivalence principle from GW170817 and its electromagnetic counterparts, *J. Cosmol. Astropart. Phys.* 2017(11), 035 (2017)
11. I. M. Shoemaker and K. Murase, Constraints from the time lag between gravitational waves and gamma rays: Implications of GW170817 and GRB 170817A, *Phys. Rev. D* 97(8), 083013 (2018)
12. B. Zhang, *The Physics of Gamma-Ray Bursts*, Cambridge: Cambridge University Press, 2018
13. J. Granot, D. Guetta, and R. Gill, Lessons from the short GRB 170817A: The first gravitational-wave detection of a binary neutron star merger, *Astrophys. J.* 850(2), L24 (2017)
14. P. Veres, P. Mészáros, A. Goldstein, N. Fraija, V. Connaughton, E. Burns, R. D. Preece, R. Hamburg, C. A. Wilson-Hodge, M. S. Briggs, and D. Kocevski, Gamma-ray burst models in light of the GRB 170817A-GW170817 connection, arXiv: 1802.07328 (2018)
15. D. B. Lin, T. Liu, J. Lin, X. G. Wang, W. M. Gu, and E. W. Liang, First electromagnetic pulse associated with a gravitational-wave event: Profile, duration, and delay, *Astrophys. J.* 856(2), 90 (2018)

16. O. S. Salafia, G. Ghisellini, G. Ghirlanda, and M. Colpi, Interpreting GRB170817A as a giant flare from a jet-less double neutron star merger, *Astron. Astrophys.* 619, A18 (2018)
17. Y. Z. Qian and S. E. Woosley, Nucleosynthesis in neutrino-driven winds (I): The physical conditions, *Astrophys. J.* 471(1), 331 (1996)
18. W. H. Lei, B. Zhang, and E. W. Liang, Hyperaccreting black hole as gamma-ray burst central engine (I): Baryon loading in gamma-ray burst jets, *Astrophys. J.* 765(2), 125 (2013)
19. B. D. Metzger, D. Giannios, T. A. Thompson, N. Bucciantini, and E. Quataert, The protomagnetar model for gamma-ray bursts, *Mon. Not. R. Astron. Soc.* 413(3), 2031 (2011)
20. P. C. Duffell, E. Quataert, D. Kasen, and H. Klion, Jet dynamics in compact object mergers: GW170817 likely had a successful jet, *Astrophys. J.* 866(1), 3 (2018)
21. K. P. Mooley, A. T. Deller, O. Gottlieb, E. Nakar, G. Hallinan, S. Bourke, D. A. Frail, A. Horesh, A. Corsi, and K. Hotokezaka, Superluminal motion of a relativistic jet in the neutron-star merger GW170817, *Nature* 561(7723), 355 (2018)
22. G. Ghirlanda, O. S. Salafia, Z. Paragi, M. Giroletti, J. Yang, et al., Compact radio emission indicates a structured jet was produced by a binary neutron star merger, *Science* 363(6430), 968 (2019)
23. M. Shibata and K. Taniguchi, Merger of black hole and neutron star in general relativity: Tidal disruption, torus mass, and gravitational waves, *Phys. Rev. D* 77(8), 084015 (2008)
24. J. J. Geng, B. Zhang, A. Kölligan, R. Kuiper, and Y. F. Huang, Propagation of a short GRB jet in the ejecta: Jet launching delay time, jet structure, and GW170817/GRB 170817A, arXiv: 1904.02326 (2019)
25. K. Ioka and T. Nakamura, Can an off-axis gamma-ray burst jet in GW170817 explain all the electromagnetic counterparts? *Prog. Theor. Exp. Phys.* 2018(4), 043E02 (2018)
26. P. Mészáros and M. J. Rees, Steep slopes and preferred breaks in gamma-ray burst spectra: The role of photospheres and comptonization, *Astrophys. J.* 530(1), 292 (2000)
27. M. J. Rees and P. Mészáros, Dissipative photosphere models of gamma-ray bursts and X-ray flashes, *Astrophys. J.* 628(2), 847 (2005)
28. A. Pe'er and F. Ryde, A theory of multicolor blackbody emission from relativistically expanding plasmas, *Astrophys. J.* 732(1), 49 (2011)
29. M. J. Rees and P. Mészáros, Unsteady outflow models for cosmological gamma-ray bursts, *Astrophys. J.* 430, L93 (1994)
30. S. Kobayashi, T. Piran, and R. Sari, Can internal shocks produce the variability in gamma-ray bursts? *Astrophys. J.* 490(1), 92 (1997)
31. B. Zhang and H. Yan, The internal-collision-induced magnetic reconnection and turbulence (ICMART) model of gamma-ray bursts, *Astrophys. J.* 726(2), 90 (2011)
32. Z. L. Uhm and B. Zhang, Toward an understanding of GRB prompt emission mechanism (I): The origin of spectral lags, *Astrophys. J.* 825(2), 97 (2016)
33. F. Daigne and R. Mochkovitch, The expected thermal precursors of gamma-ray bursts in the internal shock model, *Mon. Not. R. Astron. Soc.* 336(4), 1271 (2002)
34. A. Pe'er, P. Mészáros, and M. J. Rees, The observable effects of a photospheric component on GRB and XRF prompt emission spectrum, *Astrophys. J.* 642(2), 995 (2006)
35. B. Zhang and A. Pe'er, Evidence of an initially magnetically dominated outflow in GRB 080916C, *Astrophys. J.* 700(2), L65 (2009)
36. H. Gao, B. B. Zhang, and B. Zhang, Stepwise filter correlation method and evidence of superposed variability components in gamma-ray burst prompt emission light curves, *Astrophys. J.* 748(2), 134 (2012)
37. B. J. Morsony, D. Lazzati, and M. C. Begelman, The origin and propagation of variability in the outflows of long-duration gamma-ray bursts, *Astrophys. J.* 723(1), 267 (2010)
38. W. Deng and B. Zhang, Low energy spectral index and e_p evolution of quasi-thermal photosphere emission of gamma-ray bursts, *Astrophys. J.* 785(2), 112 (2014)
39. Ž. Bošnjak and F. Daigne, Spectral evolution in gamma-ray bursts: Predictions of the internal shock model and comparison to observations, *Astron. Astrophys.* 568, A45 (2014)
40. Z. L. Uhm, B. Zhang, and J. Racusin, Toward an understanding of GRB prompt emission mechanism (II): Patterns of peak energy evolution and their connection to spectral lags, *Astrophys. J.* 869(2), 100 (2018)
41. L. Baiotti and L. Rezzolla, Binary neutron star mergers: A review of Einstein's richest laboratory, *Rep. Prog. Phys.* 80(9), 096901 (2017)
42. S. Rosswog, E. Ramirez-Ruiz, and M. B. Davies, High-resolution calculations of merging neutron stars (III): Gamma-ray bursts, *Mon. Not. R. Astron. Soc.* 345(4), 1077 (2003)
43. R. Ciolfi, W. Kastaun, J. Vijay Kalinani, and B. Giacomazzo, The first 100 ms of a long-lived magnetized neutron star formed in a binary neutron star merger, arXiv: 1904.10222 (2019)
44. Z. G. Dai, X. Y. Wang, X. F. Wu, and B. Zhang, X-ray flares from postmerger millisecond pulsars, *Science* 311(5764), 1127 (2006)
45. W. H. Gao and Y. Z. Fan, Short-living supermassive magnetar model for the early X-ray flares following short GRBs, *Chin. J. Astron. Astrophys.* 6(5), 513 (2006)
46. B. D. Metzger, E. Quataert, and T. A. Thompson, Short-duration gamma-ray bursts with extended emission from protomagnetar spin-down, *Mon. Not. R. Astron. Soc.* 385(3), 1455 (2008)

47. A. Rowlinson, P. T. O'Brien, N. R. Tanvir, B. Zhang, P. A. Evans, N. Lyons, A. J. Levan, R. Willingale, K. L. Page, O. Onal, D. N. Burrows, A. P. Beardmore, T. N. Ukwatta, E. Berger, J. Hjorth, A. S. Fruchter, R. L. Tunnicliffe, D. B. Fox, and A. Cucchiara, The unusual X-ray emission of the short Swift GRB 090515: Evidence for the formation of a magnetar? *Mon. Not. R. Astron. Soc.* 409(2), 531 (2010)
48. A. Rowlinson, P. T. O'Brien, B. D. Metzger, N. R. Tanvir, and A. J. Levan, Signatures of magnetar central engines in short GRB light curves, *Mon. Not. R. Astron. Soc.* 430(2), 1061 (2013)
49. H. J. Lü, B. Zhang, W. H. Lei, Y. Li, and P. D. Lasky, The millisecond magnetar central engine in short GRBs, *Astrophys. J.* 805(2), 89 (2015)
50. B. Zhang, Early X-ray and optical afterglow of gravitational wave bursts from mergers of binary neutron stars, *Astrophys. J.* 763(1), L22 (2013)
51. H. Gao, B. Zhang, and H. J. Lü, Constraints on binary neutron star merger product from short GRB observations, *Phys. Rev. D* 93(4), 044065 (2016)
52. H. Sun, B. Zhang, and H. Gao, X-ray counterpart of gravitational waves due to binary neutron star mergers: Light curves, luminosity function, and event rate density, *Astrophys. J.* 835, 7 (2017)
53. Y. Q. Xue, X. C. Zheng, Y. Li, W. N. Brandt, B. Zhang, B. Luo, B. B. Zhang, F. E. Bauer, H. Sun, B. D. Lehmer, X. F. Wu, G. Yang, X. Kong, J. Y. Li, M. Y. Sun, J. X. Wang, and F. Vito, A magnetar-powered X-ray transient as the aftermath of a binary neutron-star merger, *Nature* 568(7751), 198 (2019)
54. B. Zhang and P. Mészáros, Gamma-ray burst afterglow with continuous energy injection: Signature of a highly magnetized millisecond pulsar, *Astrophys. J.* 552(1), L35 (2001)
55. D. Zhang and Z. G. Dai, Hyperaccreting disks around magnetars for gamma-ray bursts: Effects of strong magnetic fields, *Astrophys. J.* 718(2), 841 (2010)
56. C. D. Ott, The gravitational-wave signature of core-collapse supernovae, *Class. Quantum Gravity* 26(6), 063001 (2009)
57. V. V. Usov, Millisecond pulsars with extremely strong magnetic fields as a cosmological source of γ -ray bursts, *Nature* 357(6378), 472 (1992)
58. A. Corsi and P. Mészáros, Gamma-ray burst afterglow plateaus and gravitational waves: Multi-messenger signature of a millisecond magnetar? *Astrophys. J.* 702(2), 1171 (2009)
59. T. Liu, C. Y. Lin, C. Y. Song, and A. Li, Comparison of gravitational waves from central engines of gamma-ray bursts: Neutrino-dominated accretion flows, Blandford–Znajek mechanisms, and millisecond magnetars, *Astrophys. J.* 850(1), 30 (2017)
60. S. E. Woosley, Gamma-ray bursts from stellar mass accretion disks around black holes, *Astrophys. J.* 405, 273 (1993)
61. A. I. MacFadyen and S. E. Woosley, Collapsars: Gamma-ray bursts and explosions in “failed supernovae”, *Astrophys. J.* 524(1), 262 (1999)
62. A. I. MacFadyen, S. E. Woosley, and A. Heger, Supernovae, jets, and collapsars, *Astrophys. J.* 550(1), 410 (2001)
63. S. E. Woosley and J. S. Bloom, The supernova–gamma-ray burst connection, *Astronomy & Astrophysics* 44(1), 507 (2006)
64. W. Kluźniak and M. Ruderman, The central engine of gamma-ray bursters, *Astrophys. J.* 505(2), L113 (1998)
65. Z. G. Dai and T. Lu, γ -ray bursts and afterglows from rotating strange stars and neutron stars, *Phys. Rev. Lett.* 81(20), 4301 (1998)
66. M. A. Ruderman, L. Tao, and W. Kluźniak, A central engine for cosmic gamma-ray burst sources, *Astrophys. J.* 542(1), 243 (2000)
67. V. V. Usov, On the nature of non-thermal radiation from cosmological-ray bursters, *Mon. Not. R. Astron. Soc.* 267(4), 1035 (1994)
68. M. M. Kasliwal, E. Nakar, L. P. Singer, D. L. Kaplan, D. O. Cook, et al., Illuminating gravitational waves: A concordant picture of photons from a neutron star merger, *Science* 358(6370), 1559 (2017)
69. K. P. Mooley, E. Nakar, K. Hotokezaka, G. Hallinan, A. Corsi, et al., A mildly relativistic wide-angle outflow in the neutron-star merger event GW170817, *Nature* 554(7691), 207 (2018)
70. E. Nakar and T. Piran, Implications of the radio and X-ray emission that followed GW170817, *Mon. Not. R. Astron. Soc.* 478(1), 407 (2018)
71. O. Gottlieb, E. Nakar, and T. Piran, The cocoon emission – an electromagnetic counterpart to gravitational waves from neutron star mergers, *Mon. Not. R. Astron. Soc.* 473(1), 576 (2018)
72. O. Bromberg, A. Tchekhovskoy, O. Gottlieb, E. Nakar, and T. Piran, The γ -rays that accompanied GW170817 and the observational signature of a magnetic jet breaking out of NS merger ejecta, *Mon. Not. R. Astron. Soc.* 475(3), 2971 (2018)
73. B. Margalit and B. D. Metzger, Constraining the maximum mass of neutron stars from multi-messenger observations of GW170817, *Astrophys. J.* 850(2), L19 (2017)
74. R. Gill, A. Nathanail, and L. Rezzolla, When did the remnant of GW170817 collapse to a black hole? arXiv: 1901.04138 (2019)
75. M. Ruiz, S. L. Shapiro, and A. Tsokaros, GW170817, general relativistic magnetohydrodynamic simulations, and the neutron star maximum mass, *Phys. Rev. D* 97(2), 021501 (2018)
76. L. Rezzolla, E. R. Most, and L. R. Weih, Using gravitational-wave observations and quasi-universal relations to constrain the maximum mass of neutron stars, *Astrophys. J.* 852(2), L25 (2018)
77. A. Loeb, electromagnetic counterparts to black hole mergers detected by LIGO, *Astrophys. J.* 819(2), L21 (2016)

78. S. E. Woosley, The progenitor of GW150914, *Astrophys. J.* 824(1), L10 (2016)
79. L. Dai, J. C. McKinney, and M. C. Miller, Energetic constraints on electromagnetic signals from double black hole mergers, *Mon. Not. R. Astron. Soc.* 470(1), L92 (2017)
80. D. D’Orazio and A. Loeb, Single progenitor model for GW150914 and GW170104, *Phys. Rev. D* 97(8), 083008 (2018)
81. A. Janiuk, M. Bejger, S. Charzyński, and P. Sukova, On the possible gamma-ray burst–gravitational wave association in GW150914, *New Astron.* 51, 7 (2017)
82. R. Perna, D. Lazzati, and B. Giacomazzo, Short gamma-ray bursts from the merger of two black holes, *Astrophys. J.* 821(1), L18 (2016)
83. S. S. Kimura, S. Z. Takahashi, and K. Toma, Evolution of an accretion disc in binary black hole systems, *Mon. Not. R. Astron. Soc.* 465(4), 4406 (2017)
84. B. Zhang, Mergers of charged black holes: Gravitational-wave events, short gamma-ray bursts, and fast radio bursts, *Astrophys. J.* 827(2), L31 (2016)
85. B. Zhang, Charged compact binary coalescence signal and electromagnetic counterpart of plunging black hole–neutron star mergers, *Astrophys. J.* 873(2), L9 (2019)
86. Z. G. Dai, Inspiral of a spinning black hole–magnetized neutron star binary: Increasing charge and electromagnetic emission, *Astrophys. J.* 873(2), L13 (2019)