

## RESEARCH ARTICLE

 $p + ip$ -wave pairing symmetry at type-II van Hove singularitiesYin-Xiang Li<sup>1,†</sup>, Xiao-Tong Yang<sup>1</sup><sup>1</sup>*Tin Ka-Ping College of Science, University of Shanghai for Science and Technology, Shanghai 200093, China*  
Corresponding author. E-mail: <sup>†</sup>[yinxiangl@usst.edu.cn](mailto:yinxiangl@usst.edu.cn)

Received November 20, 2020; accepted March 5, 2021

Based on the random phase approximation calculation in two-orbital honeycomb lattice model, we investigate the pairing symmetry of Ni-based transition-metal trichalcogenides by electron doping access to type-II van Hove singularities (vHs). We find that chiral even-parity  $d + id$ -wave ( $E_g$ ) state is suppressed by odd-parity  $p + ip$ -wave ( $E_u$ ) state when electron doping approaches the type-II vHs. The type-II vHs peak in density of states (DOS) enables to strengthen the ferromagnetic fluctuation, which is responsible for triplet pairing. The competition between antiferromagnetic and ferromagnetic fluctuation results in pairing phase transition from singlet to triplet pairing. The Ni-based transition-metal trichalcogenides provide a promising platform to unconventional superconductor emerging from electronic DOS.

**Keywords** type-II van Hove singularities, pairing symmetry, random phase approximation, unconventional superconductor

## 1 Introduction

In the past decades, novel topological states of quantum matters are the active and attractive topics in condensed matter physics. The discovery of quantum spin Hall in HgTe has boosted the search of two-dimensional (2D) topological insulator [1, 2]. After that, three-dimensional (3D) topological insulators and topological semimetals have also been verified in both theoretical calculations and experiments [3–9]. Meanwhile, topological superconductor [10–17] has also attracted tremendous attentions due to the development of topological insulator. The topological superconductor with particle–hole symmetry can host Majorana zero mode which may be potentially employed in realizing topological quantum computation [18, 19]. Especially, it is interesting that 2D chiral  $p+ip$  wave topological superconductor in magnetic vortex cores can host Majorana zero modes [20]. Recently, the outstanding studies are experimental evidences for Majorana bound states in an iron-based superconductor  $\text{FeTe}_{0.55}\text{Se}_{0.45}$  by scanning tunneling spectroscopy [13, 21, 22]. Thus, searching for intrinsic topological superconductor is the active and prominent field in condensed matter physics.

2D materials include not only quantum spin Hall insulators and Chern insulators, but also unconventional superconductor as doped twisted bilayer graphene [23–26]. Among them, ternary transition-metal phosphorus

trichalcogenide (TMPT) compounds  $\text{APX}_3$  ( $A = 3d$  transition metals;  $X =$  chalcogens) have attracted enormous attention due to antiferromagnetic (AF) ordering as a hint for significant electronic correlations [27–31]. By suppressing AF order with external pressure, superconductivity emerges in iron-based TMPT compounds such as  $\text{FePSe}_3$ , with the highest  $T_c$  found at about 5.5 K [32]. The crystal structure of the TMPT family  $\text{APX}_3$  consists of edge shared  $\text{AX}_6$  octahedral complexes and P2 dimers. Transition metal atoms are arranged in a hexagonal lattice. In the octahedral crystal field, five 3d orbitals of transition-metal atoms split into high-energy  $e_g$  orbitals and low-energy  $t_{2g}$  orbitals. For  $\text{FePX}_3$  with  $\text{Fe}^{2+}$  ions ( $d^6$ ), it is an ideal system to study the high-to-low spin-state transition by pressure. For the case of  $\text{NiPX}_3$  with  $\text{Ni}^{2+}$   $d^8$  filling configuration,  $t_{2g}$  bands are fully occupied while  $e_g$  bands are half filled and dominate the spectral weight near the Fermi level. Theoretical calculations suggest that charge doping can suppress magnetic order, and superconductivity can eventually be achieved for  $\text{NiPS}_3$  [33].

With increasing electron doping, the system far away from half filling and approaches type-II van Hove singularities (vHs) [34] along  $\Gamma$ – $M$  and  $\Gamma$ – $K$  high symmetry line. For 2D superconductor, a vHs in the density of states (DOS) was proposed to drive a substantial enhancement of interaction effects and to promote the unconventional superconductor [34–45]. In general, superconductors with type-I vHs (the saddle points locate at time reversal invariant momentums (TRIMs)) favor singlet pairing. For type-II vHs superconductors (the saddle points at general  $k$  points), the triplet pair can compete with singlet

\*arXiv: 2103.1375. This article can also be found at <http://journal.hep.com.cn/fop/EN/10.1007/s11467-021-1068-7>.



pair [34]. By random phase approximation (RPA) calculation in honeycomb lattice, we find that the  $p + ip$ -wave ( $E_u$ ) pairing is enhancing and then overcomes an I-wave ( $A_{2g}$ ) state and a chiral  $d$ -wave ( $E_g$ ) state when electron doping from half filling to type-II vHs.

In this paper, the pairing symmetry of Ni-based transition-metal trichalcogenide superconductor is studied near type-II vHs, away from half filling 0.14 eV. Within RPA pattern, we use two-sublattice two-orbital Hubbard model to calculate the pairing symmetry of 2D van der Waals (vdW) material NiPS<sub>3</sub>. We find that the chiral even-parity  $d + id$ -wave ( $E_g$ ) state is suppressed by odd-parity  $p + ip$ -wave ( $E_u$ ) state when electron doping approaches the type-II vHs. In the lower doping level, chiral even-parity  $d + id$ -wave ( $E_g$ ) is the dominant pairing in the system [46]. There exist pairing phase transition for this 2D superconductivity material, from singlet pairing to triplet pairing. The increasement of DOS from type-II vHs will strengthen the triplet pair. The pairing result from RPA is consistent with the analysis from spin susceptibility calculation. The peak of spin susceptibility in  $\Gamma$  implies ferromagnetic fluctuation will be responsible for triplet pairing. The Fermi surface nesting from intra  $\beta$  pockets promotes the instability of ferromagnetic fluctuation.

The paper is organized as follows. In Section 2, we present the band structure, DOS and Fermi surface from two-sublattice two-orbital tight-binding model based on  $e_g$  orbitals ( $d_{xz}$  and  $d_{yz}$  orbitals). We find that type II vHs are only 0.14 eV above the Fermi level. In Section 3, we show the formalism of RPA approach for superconductor pairing based on multi-orbital Coulomb interactions. In Section 4, we analyse the spin susceptibility and pairing symmetry when electron doping is closing to vHs. The triplet pairing  $p + ip$  ( $E_u$ ) is the leading superconductivity state, which is caused by ferromagnetic fluctuation. Finally, we summarize and discuss these results in Section 5.


## 2 Electronic structure

The nickel phosphorous trichalcogenides compounds NiPX<sub>3</sub> (X=S, Se) are 2D vdW materials, which consist of layered hexagonal structures [33]. Each layer is constructed by  $MX_6$  edge-shared octahedral complexes. In this octahedral environment, the five d orbitals of Ni atom are split into  $t_{2g}$  and  $e_g$  groups. The  $e_g$  orbitals are close to half-filling while the  $t_{2g}$  orbitals are fully filled. The physics near Fermi surface are mainly from the  $d_{xz}$  and  $d_{yz}$  orbitals. In order to capture the low-energy physics, we use two-sublattice two-orbital tight-binding model on honeycomb lattice. The corresponding tight-binding model [33] is given by

$$H_0 = \sum_{\mathbf{k}\alpha\beta} \sum_{\mu\nu\sigma} h_{\mu\nu}^{\alpha\beta}(\mathbf{k}) c_{\mathbf{k}\alpha\mu\sigma}^\dagger c_{\mathbf{k}\beta\nu\sigma} = \sum_{\mathbf{k}} \psi_{\mathbf{k}\sigma}^\dagger h(\mathbf{k}) \psi_{\mathbf{k}\sigma}, \quad (1)$$

with  $\psi_{\mathbf{k}\sigma}^\dagger = (c_{\mathbf{k}A1\sigma}^\dagger, c_{\mathbf{k}A2\sigma}^\dagger, c_{\mathbf{k}B1\sigma}^\dagger, c_{\mathbf{k}B2\sigma}^\dagger)$ . Here  $\alpha, \beta$  are the sublattice indices ( $A, B$ ) and  $\mu, \nu$  are the orbital indexes ( $d_{xz}, d_{yz}$ ).  $c_{\alpha\mu\sigma}^\dagger$  creates a spin  $\sigma$  electron with momentum  $\mathbf{k}$  in  $\mu$  orbital on  $\alpha$  sublattice. The matrix elements of  $h_{\mu\nu}^{\alpha\beta}(\mathbf{k})$  are provided in the Appendix of Reference [46]. It is interesting that the leading hopping parameter is third nearest neighbor (TNN) hopping term. The TNN Ni cations formed superexchange antiferromagnetic state is favored in NiPX<sub>3</sub> (X = S, Se) parent compounds.

In Fig. 1(a), we show the orbital resolved band dispersion ( $d_{xz}$  and  $d_{yz}$  orbitals) from the tight-binding model. At pristine filling, there are eight Dirac points protected by  $D_{3d}$  symmetry near the Fermi level. Due to charge conversation, hole pocket and electron pocket appears at  $K/2$  and  $K$  respectively. Based on the two-fold rotational symmetry along  $\Gamma$ - $M$ , the mixture of  $d_{xz}$  and  $d_{yz}$  orbitals can be found along  $\Gamma$ - $K$  and  $K$ - $M$ , but not  $\Gamma$ - $M$ . The strongest orbital mixture occurs near the Fermi level and also around the Dirac points ( $K$  and  $K/2$ ). In order to analyse saddle points above the Fermi surface, we calculate the corresponding DOS in Fig. 1(b). We find DOS peak above the Fermi level near 0.14 eV, which verify the existence of vHs. We plot related orbital resolved Fermi surface with  $\delta = 0.3$  and  $\delta = 0.35$  per Ni atom with respect to the half-filling in Figs. 1(c) and (d). For hexagonal symmetry in two-dimensional honeycomb lattice, there are six type-II vHs (saddle points not at TRIM points) along  $\Gamma$ - $M$  or  $\Gamma$ - $K$  direction. When changing the doping level around vHs, there accompany Lifshitz transition of Fermi surfaces. Two pockets around  $K/2$  make together to be-



**Fig. 1** The orbital characters of bands are represented by different colors. (a) Band dispersion of tight-binding model based on  $d_{xz}$  and  $d_{yz}$  orbitals of Ni atoms. (b) The corresponding density of states (DOS) for NiPS<sub>3</sub>. The van Hove singularities (vHs) are above the Fermi level near 0.14 eV. Orbital resolved Fermi surface with two different electron fillings,  $\delta = 0.3$  (c) and  $\delta = 0.35$  (d) per Ni atom with respect to the half-filling. The type-II vHs appear along  $\Gamma$ - $M$  line or  $\Gamma$ - $K$  line which are not at TRIM points.

come one Fermi surface in Fig. 1(c). The outer pocket around  $\Gamma$  is fusing with  $K$  pocket as shown in Fig. 1(d). The DOS peak is closely related with topology of Fermi surface, and also make great influence on superconductor pairing. Generally, triplet pairing could compete with singlet pairing in the system with type-II vHs [34], which also be verified by our RPA calculation in the following section.

### 3 Random phase approximation

Based on two-sublattice two-orbital tight-binding model, we consider onsite multi-orbital Hubbard interaction for superconductor pairing as

$$\begin{aligned}
 H_{int} = & U \sum_{i,\alpha} n_{i\alpha\uparrow} n_{i\alpha\downarrow} + U' \sum_{i,\alpha<\beta} n_{i\alpha} n_{i\beta} \\
 & + J_H \sum_{i,\alpha<\beta,\sigma\sigma'} c_{i\alpha\sigma}^\dagger c_{i\beta\sigma'}^\dagger c_{i\alpha\sigma'} c_{i\beta\sigma} \\
 & + J' \sum_{i,\alpha\neq\beta} c_{i\alpha\uparrow}^\dagger c_{i\alpha\downarrow}^\dagger c_{i\beta\downarrow} c_{i\beta\uparrow}, \quad (2)
 \end{aligned}$$

where  $n_{i,\alpha} = n_{i,\alpha,\uparrow} + n_{i,\alpha,\downarrow}$ .  $U$ ,  $U'$ ,  $J$ , and  $J'$  represent the intra- and inter-orbital repulsion, the Hund's rule and pair-hopping terms. We adopt Kanamori relations  $U = U' + 2J$  and  $J = J'$  in the next calculation, which is required by the lattice symmetry. Considering RPA approximation [47, 48], the multi-orbital susceptibility is defined as

$$\begin{aligned}
 \chi_{l_1 l_2 l_3 l_4}(\mathbf{q}, \tau) = & \frac{1}{N} \sum_{\mathbf{k}\mathbf{k}'} \langle T_\tau c_{l_3\sigma}^\dagger(\mathbf{k} + \mathbf{q}, \tau) c_{l_4\sigma}(\mathbf{k}, \tau) \\
 & \times c_{l_2\sigma'}^\dagger(\mathbf{k}' - \mathbf{q}, 0) c_{l_1\sigma'}(\mathbf{k}', 0) \rangle. \quad (3)
 \end{aligned}$$

In momentum-frequency space, the multi-orbital bare susceptibility is given by

$$\begin{aligned}
 \chi_{l_1 l_2 l_3 l_4}^0(\mathbf{q}, i\omega_n) = & -\frac{1}{N} \sum_{\mathbf{k}\mu\nu} a_\mu^{l_4}(\mathbf{k}) a_\mu^{l_2^*}(\mathbf{k}) a_\nu^{l_1}(\mathbf{k} + \mathbf{q}) \\
 & \times a_\nu^{l_3^*}(\mathbf{k} + \mathbf{q}) \frac{n_F(E_\mu(\mathbf{k})) - n_F(E_\nu(\mathbf{k} + \mathbf{q}))}{i\omega_n + E_\mu(\mathbf{k}) - E_\nu(\mathbf{k} + \mathbf{q})}, \quad (4)
 \end{aligned}$$

where  $\mu$  and  $\nu$  are the band indices,  $n_F$  is the usual Fermi distribution,  $l_i$  ( $i = 1, 2, 3, 4$ ) are the orbital indices. By diagonalizing the above multi-orbital tight-binding Hamiltonian, we obtain the  $l_i$  orbital component of the eigenvector for band  $\mu$  ( $a_\mu^{l_i}(\mathbf{k})$ ) and eigenvalue  $E_\mu(\mathbf{k})$ . After that, we take the multi-orbital Hubbard interactions into consideration for calculating RPA susceptibilities. The corresponding RPA spin and charge susceptibilities [41, 49, 50] are given by

$$\begin{aligned}
 \chi_s^{RPA}(\mathbf{q}) = & \chi^0(\mathbf{q}) [1 - \bar{U}^s \chi^0(\mathbf{q})]^{-1}, \\
 \chi_c^{RPA}(\mathbf{q}) = & \chi^0(\mathbf{q}) [1 + \bar{U}^c \chi^0(\mathbf{q})]^{-1}, \quad (5)
 \end{aligned}$$

where  $\bar{U}^s$  ( $\bar{U}^c$ ) is the spin (charge) interaction matrix

$$\begin{aligned}
 \bar{U}^{s/c} = & \begin{pmatrix} \bar{U}_A^{s/c} & 0 \\ 0 & \bar{U}_B^{s/c} \end{pmatrix}, \\
 \bar{U}_{A/B, l_1 l_2 l_3 l_4}^s = & \begin{cases} U & l_1 = l_2 = l_3 = l_4, \\ U' & l_1 = l_3 \neq l_2 = l_4, \\ J & l_1 = l_2 \neq l_3 = l_4, \\ J' & l_1 = l_4 \neq l_3 = l_2, \end{cases} \\
 \bar{U}_{A/B, l_1 l_2 l_3 l_4}^c = & \begin{cases} U & l_1 = l_2 = l_3 = l_4, \\ -U' + 2J & l_1 = l_3 \neq l_2 = l_4, \\ 2U' - J & l_1 = l_2 \neq l_3 = l_4, \\ J' & l_1 = l_4 \neq l_3 = l_2. \end{cases}
 \end{aligned}$$

In this process, we only consider the electron scattering from Fermi surfaces near the Fermi level. The effective Cooper scattering interaction is written as

$$\begin{aligned}
 \Gamma_{ij}(\mathbf{k}, \mathbf{k}') = & \sum_{l_1 l_2 l_3 l_4} a_{v_i}^{l_2^*}(\mathbf{k}) a_{v_i}^{l_3^*}(-\mathbf{k}) \\
 & \times Re \left[ \Gamma_{l_1 l_2 l_3 l_4}(\mathbf{k}, \mathbf{k}', \omega = 0) \right] a_{v_j}^{l_1}(\mathbf{k}') a_{v_j}^{l_4}(-\mathbf{k}'), \quad (6)
 \end{aligned}$$

where the momenta  $\mathbf{k}$  and  $\mathbf{k}'$  are confined in different FSs with  $\mathbf{k} \in C_i$  and  $\mathbf{k}' \in C_j$ . The orbital vertex function  $\Gamma_{l_1 l_2 l_3 l_4}$  in spin singlet and triplet channels [47–49] are

$$\begin{aligned}
 \Gamma_{l_1 l_2 l_3 l_4}^S(\mathbf{k}, \mathbf{k}', \omega) = & \left[ \frac{3}{2} \bar{U}^s \chi_s^{RPA}(\mathbf{k} - \mathbf{k}', \omega) \bar{U}^s \right. \\
 & \left. + \frac{1}{2} \bar{U}^s - \frac{1}{2} \bar{U}^c \chi_c^{RPA}(\mathbf{k} - \mathbf{k}', \omega) \bar{U}^c + \frac{1}{2} \bar{U}^c \right]_{l_1 l_2 l_3 l_4}, \\
 \Gamma_{l_1 l_2 l_3 l_4}^T(\mathbf{k}, \mathbf{k}', \omega) = & \left[ -\frac{1}{2} \bar{U}^s \chi_s^{RPA}(\mathbf{k} - \mathbf{k}', \omega) \bar{U}^s \right. \\
 & \left. + \frac{1}{2} \bar{U}^s - \frac{1}{2} \bar{U}^c \chi_c^{RPA}(\mathbf{k} - \mathbf{k}', \omega) \bar{U}^c + \frac{1}{2} \bar{U}^c \right]_{l_1 l_2 l_3 l_4}, \quad (7)
 \end{aligned}$$

where  $\chi_s^{RPA}$  and  $\chi_c^{RPA}$  are the RPA spin and charge susceptibility respectively. For spin singlet and triplet channels, the pairing vertex functions are symmetric and antisymmetric parts of the interaction,  $\Gamma_{ij}^{S/T}(\mathbf{k}, \mathbf{k}') = \frac{1}{2} [\Gamma_{ij}(\mathbf{k}, \mathbf{k}') \pm \Gamma_{ij}(\mathbf{k}, -\mathbf{k}')]$ . Then, the pairing strength function for superconductor [42, 49–53] is

$$\begin{aligned}
 \lambda[g(\mathbf{k})] = & \frac{\sum_{ij} \oint_{C_i} \frac{d\mathbf{k}'_{//}}{v_F(\mathbf{k})} \oint_{C_j} \frac{d\mathbf{k}'_{//}}{v_F(\mathbf{k}')} g(\mathbf{k}) \Gamma_{ij}^{S/T}(\mathbf{k}, \mathbf{k}') g(\mathbf{k}')}{(2\pi)^2 \sum_i \oint_{C_i} \frac{d\mathbf{k}'_{//}}{v_F(\mathbf{k})} [g(\mathbf{k})]^2}, \quad (8)
 \end{aligned}$$


where  $v_F(\mathbf{k}) = |\nabla_{\mathbf{k}} E_i(\mathbf{k})|$  is the Fermi velocity on a given Fermi surface sheet  $C_i$ .

## 4 Susceptibility and pairing symmetry


Based on the multi-orbital RPA method (weak coupling approach) [47, 48], we investigate the pairing symmetry of electron doped nickel phosphorous trichalcogenides compounds NiPS<sub>3</sub>. Due to the existence of type-II vHs peak in DOS near Fermi level, we mainly discuss the electron doping level up to  $\delta = 0.3$  and  $\delta = 0.35$  per Ni atom separately. In order to analyse the pairing symmetry results, we first calculate the bare susceptibility  $\chi_0$  and spin susceptibility  $\chi_{RPA}^s$  along high-symmetry lines at two different doping levels in Figs. 2(a) and (c). For bare susceptibility at  $\delta = 0.3$  per Ni atom doping level [blue dash-dot line in Fig. 2(a)], there is a prominent peak at  $\Gamma$ , a smooth plateau around  $M$  and a broaden peak at  $K/2$ . The first one is mainly contributed by the intrapocket nesting between rather flat bands in NNN pockets  $\beta$  [ $Q_1$  in Fig. 1(c)]. The intrapocket nesting  $Q_2$  between NNN pockets  $\alpha$  is responsible for the second peak around  $M$ . The third peak at  $K/2$  is ascribed to the intrapocket nesting  $Q_3$  between NN pockets  $\beta$ . Then, we consider the RPA spin susceptibility for superconducting instability with  $U = 0.3$  eV and  $J/U = 0.2$  in Fig. 2(a). All the mentioned above peaks are enhanced significantly. Especially, the sharp peak at  $\Gamma$  near divergence indicates the ferromagnetic fluctuation between unit cells is dominant. By checking the eigenvectors of susceptibility matrix corresponding to the largest eigenvalue, we find all the signs of eigenvectors are positive which implies ferromagnetic fluctuation in a unit cell. Compared with lower doping level at  $\delta = 0.1$  per Ni atom, the emergence of spin susceptibility peak at  $\Gamma$  means that the ferromagnetic fluctuation is competing with anti-

ferromagnetic fluctuation. Undoubtedly, the type-II vHs could strengthen the ferromagnetic fluctuation. In order to make the spin susceptibility clearly, we plot the corresponding susceptibility in 2D Brillouin zone. From 2D pattern in Fig. 2(b), it is clear that  $C_6$  symmetry is maintained, a sharp peak at  $\Gamma$  and another peaks around  $M$ . With further doping to  $\delta = 0.35$  per Ni atom, the peak at  $\Gamma$  becomes sharper and the peaks around  $M$  move toward  $\Gamma$  and  $K$  in Figs. 2(c) and (d). Due to the underestimation of interaction parameter in RPA method, we adopt intraorbital repulsive interaction parameter  $U = 0.3$  below the critical point  $U_c = 0.35$  (to avoid magnetic instability) and Hund's coupling  $J/U \leq 0.2$ .


In this section, we mainly focus on the doping level near type-II vHs. Based on the irreducible representation of  $D_{3d}$  point group in this material, we classify the pairing states into subgroups of this point group. Figure 3(a) shows the leading pairing strengths in singlet and triplet channels with different electron dopings at fixed  $U = 0.3$  and  $J/U = 0.2$ . From lower doping ( $\delta = 0.1$ ) to type-II vHs ( $\delta = 0.35$ ), the system exists pairing phase transition from singlet pair to triplet pair. In the low doping level, the nearly degenerate singlet pair states  $E_g(x^2 - y^2, xy)$  and  $A_{2g}$  overcome the triplet pair states  $E_u(x, y)$  and  $A_{2u}$ . When the doping level near type-II vHs, these singlet pairings are suppressed by triplet pair state  $E_u(x, y)$ . Undoubtedly, the type-II vHs peak in DOS enhances the strength of triplet pairing. This result has also been anticipated by above RPA susceptibility analysis that the dominating ferromagnetic fluctuation favors triplet pairing. In Fig. 3(b), we plot the pairing eigenvalues as a function of  $U$  with a fixed  $J/U = 0.2$  with  $\delta = 0.35$  per Ni atom. The leading pairing state is still  $E_u(x, y)$  and the pairing strength is increasing with increased interaction  $U$ . In Fig. 4, we plot the gap functions of leading two-fold degenerate pairing state  $E_u(x, y)$  with  $U = 0.3$  and  $J/U = 0.2$  at  $\delta = 0.3$  (a, b) and  $\delta = 0.35$  (c, d) per Ni atom respectively. For  $p_x$  ( $p_y$ ) with  $\delta = 0.3$ , the pairing nodes along  $y$  ( $x$ ) axis with mirror symmetry  $M_x$  ( $M_y$ ). The gap function on  $\beta$  pocket is comparable to that of  $\alpha$  pocket. The superconducting orders connected by nesting vector ( $Q_1$ )



**Fig. 2** Susceptibilities for different electron doping. The bare susceptibilities  $\chi_0$  and RPA spin susceptibilities  $\chi_{RPA}^s$  along high-symmetry lines for  $\delta = 0.3$  (a) and  $\delta = 0.35$  (c) per Ni atom (near type-II vHs). The corresponding RPA spin susceptibilities in 2D Brillouin zone are shown in (b) and (d) with  $U = 0.3$  and  $J/U = 0.2$ .



**Fig. 3** Leading pairing strengths in spin singlet [ $E_g$  ( $d$ -wave),  $A_{2g}$  ( $I$ -wave)] and triplet [ $E_u$  ( $p$ -wave),  $A_{2u}$  ( $f$ -wave)] channels for superconducting states with different electron dopings  $\delta$  at  $U = 0.3$  and  $J/U = 0.2$  in (a). The pairing strengths with  $\delta = 0.35$  per Ni atom at  $J/U = 0.2$  are plotted in (b).



**Fig. 4** The gap functions  $p_x$  and  $p_y$  of leading triplet pairings with  $U = 0.3$  and  $J/U = 0.2$  at  $\delta = 0.3$  (a, b) and  $\delta = 0.35$  per Ni atom (c, d).

between flat bands in NNN pockets  $\beta$  have the same sign. For gap functions with  $\delta = 0.35$  in Figs. 4(c) and (d), there have the similar phenomenon in system. From above RPA calculation, type-II vHs peak in DOS induces odd-parity  $p_x + ip_y$ -wave ( $E_u$ ) pairing state in electron doped nickel phosphorous trichalcogenides compounds  $\text{NiPS}_3$ .

## 5 Conclusion

In this paper, we have investigated the pairing symmetry for Ni-based transition metal trichalcogenide  $\text{NiPS}_3$  based on two-sublattice two-orbital Hubbard model. By applying multi-orbital RPA method, we find that the odd-parity  $p+ip$  ( $E_u$ ) pairing state overcomes chiral even-parity  $d+id$  ( $E_g$ ) state. The enhancement of ferromagnetic fluctuation induced by type-II vHs peak in DOS is responsible for triplet pairing  $E_u$ . The nesting vector  $Q_1$  between NNN pockets  $\beta$  results in the instability peak of RPA spin susceptibility at  $\Gamma$ . This implies triplet pairing is the leading state which is consistent with our RPA's pairing calculation. The competition between ferromagnetic and antiferromagnetic fluctuation makes the transition from singlet to triplet pairing while doping approaches the type-II vHs. The effect of electronic DOS on unconventional superconductor's pairing may be realized in the layered Ni-based transition-metal trichalcogenides.

## References

1. B. A. Bernevig, T. L. Hughes, and S. C. Zhang, Quantum spin Hall effect and topological phase transition in HgTe quantum wells, *Science* 314(5806), 1757 (2006)
2. M. König, S. Wiedmann, C. Brüne, A. Roth, H. Buhmann, L. W. Molenkamp, X. L. Qi, and S. C. Zhang, Quantum spin Hall insulator state in HgTe quantum wells, *Science* 318, 766 (2007)
3. H. J. Zhang, C. X. Liu, X. L. Qi, X. Dai, Z. Fang, and S. C. Zhang, Topological insulators in  $\text{Bi}_2\text{Se}_3$ ,  $\text{Bi}_2\text{Te}_3$  and  $\text{Sb}_2\text{Te}_3$  with a single Dirac cone on the surface, *Nat. Phys.* 5(6), 438 (2009)
4. Y. Xia, D. Qian, D. Hsieh, L. Wray, A. Pal, H. Lin, A. Bansil, D. Grauer, Y. S. Hor, R. J. Cava, and M. Z. Hasan, Observation of a large-gap topological-insulator class with a single Dirac cone on the surface, *Nat. Phys.* 5(6), 398 (2009)
5. L. Fu, Topological crystalline insulators, *Phys. Rev. Lett.* 106(10), 106802 (2011)
6. T. H. Hsieh, H. Lin, J. W. Liu, W. H. Duan, A. Bansil, and L. Fu, Topological crystalline insulators in the SnTe material class, *Nat. Commun.* 3(1), 982 (2012)
7. Z. J. Wang, H. M. Weng, Q. S. Wu, X. Dai, and Z. Fang, Three-dimensional Dirac semimetal and quantum transport in  $\text{Cd}_3\text{As}_2$ , *Phys. Rev. B* 88(12), 125427 (2013)
8. H. M. Weng, C. Fang, Z. Fang, B. A. Bernevig, and X. Dai, Weyl semimetal phase in noncentrosymmetric transition-metal monophosphides, *Phys. Rev. X* 5(1), 011029 (2015)
9. B. Q. Lv, N. Xu, H. M. Weng, J. Z. Ma, P. Richard, X. C. Huang, L. X. Zhao, G. F. Chen, C. E. Matt, F. Bisti, V. N. Strocov, J. Mesot, Z. Fang, X. Dai, T. Qian, M. Shi, and H. Ding, Observation of Weyl nodes in TaAs, *Nat. Phys.* 11(9), 724 (2015)
10. N. N. Hao and J. P. Hu, Topological phases in the single-layer FeSe, *Phys. Rev. X* 4(3), 031053 (2014)
11. X. X. Wu, S. S. Qin, Y. Liang, C. C. Le, H. Fan, and J. P. Hu,  $\text{CaFeAs}_2$ : A staggered intercalation of quantum spin Hall and high-temperature superconductivity, *Phys. Rev. B* 91(8), 081111 (2015)
12. X. X. Wu, S. S. Qin, Y. Liang, H. Fan, and J. P. Hu, Topological characters in  $\text{Fe}(\text{Te}_{1-x}\text{Se}_x)$  thin films, *Phys. Rev. B* 93(11), 115129 (2016)
13. D. F. Wang, L. Y. Kong, P. Fan, H. Chen, S. Y. Zhu, W. Y. Liu, L. Cao, Y. J. Sun, S. X. Du, J. Schneeloch, R. D. Zhong, G. D. Gu, L. Fu, H. Ding, and H. J. Gao, Evidence for Majorana bound states in an iron-based superconductor, *Science* 362(6412), 333 (2018)
14. Z. B. Yan, F. Song, and Z. Wang, Majorana corner modes in a high-temperature platform, *Phys. Rev. Lett.* 121(9), 096803 (2018)
15. Q. Y. Wang, C. C. Liu, Y. M. Lu, and F. Zhang, High-temperature Majorana corner states, *Phys. Rev. Lett.* 121(18), 186801 (2018)
16. R. X. Zhang, W. S. Cole, and S. Das Sarma, Helical hinge Majorana modes in iron-based superconductors, *Phys. Rev. Lett.* 122(18), 187001 (2019)
17. X. X. Wu, W. A. Benalcazar, Y. X. Li, R. Thomale, C. X. Liu, and J. P. Hu, Boundary-obstructed topological high- $T_c$  superconductivity in iron pnictides, *Phys. Rev. X* 10(4), 041014 (2020)
18. A. Y. Kitaev, Fault-tolerant quantum computation by anyons, *Ann. Phys.* 303(1), 2 (2003)

19. M. H. Freedman, *P/NP*, and the quantum field computer, *Proc. Natl. Acad. Sci. USA* 95(1), 98 (1998)
20. N. Read and D. Green, Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect, *Phys. Rev. B* 61(15), 10267 (2000)
21. L. Y. Kong, S. Y. Zhu, M. Papaj, H. Chen, L. Cao, H. Isobe, Y. Q. Xing, W. Y. Liu, D. F. Wang, P. Fan, Y. J. Sun, S. X. Du, J. Schneeloch, R. D. Zhong, G. D. Gu, L. Fu, H. J. Gao, and H. Ding, Half-integer level shift of vortex bound states in an iron-based superconductor, *Nat. Phys.* 15(11), 1181 (2019)
22. S. Y. Zhu, L. Y. Kong, L. Cao, H. Chen, M. Papaj, S. X. Du, Y. Q. Xing, W. Y. Liu, D. F. Wang, C. M. Shen, F. Z. Yang, J. Schneeloch, R. D. Zhong, G. D. Gu, L. Fu, Y. Y. Zhang, H. Ding, and H. J. Gao, Nearly quantized conductance plateau of vortex zero mode in an iron-based superconductor, *Science* 367(6474), 189 (2020)
23. Y. Cao, V. Fatemi, S. Fang, K. Watanabe, T. Taniguchi, E. Kaxiras, and P. Jarillo-Herrero, Unconventional superconductivity in magic-angle graphene superlattices, *Nature* 556(7699), 43 (2018)
24. Y. Cao, V. Fatemi, A. Demir, S. Fang, S. L. Tomarken, J. Y. Luo, J. D. Sanchez-Yamagishi, K. Watanabe, T. Taniguchi, E. Kaxiras, R. C. Ashoori, and P. Jarillo-Herrero, Correlated insulator behaviour at half-filling in magic-angle graphene superlattices, *Nature* 556(7699), 80 (2018)
25. M. Yankowitz, S. Chen, H. Polshyn, Y. Zhang, K. Watanabe, T. Taniguchi, D. Graf, A. F. Young, and C. R. Dean, Tuning superconductivity in twisted bilayer graphene, *Science* 363(6431), 1059 (2019)
26. X. Lu, P. Stepanov, W. Yang, M. Xie, M. A. Aamir, I. Das, C. Urgell, K. Watanabe, T. Taniguchi, G. Zhang, A. Bachtold, A. H. MacDonald, and D. K. Efetov, Superconductors, orbital magnets and correlated states in magic-angle bilayer graphene, *Nature* 574(7780), 653 (2019)
27. A. R. Wildes, V. Simonet, E. Ressouche, G. J. McIntyre, M. Avdeev, E. Suard, S. A. J. Kimber, D. Lancon, G. Pepe, B. Moubaraki, and T. J. Hicks, Magnetic structure of the quasi-two-dimensional antiferromagnet NiPS<sub>3</sub>, *Phys. Rev. B* 92(22), 224408 (2015)
28. B. L. Chittari, Y. Park, D. Lee, M. Han, A. H. MacDonald, E. Hwang, and J. Jung, Electronic and magnetic properties of single-layer MPX<sub>3</sub> metal phosphorous trichalcogenides, *Phys. Rev. B* 94(18), 184428 (2016)
29. N. Sivadas, M. W. Daniels, R. H. Swendsen, S. Okamoto, and D. Xiao, Magnetic ground state of semiconducting transition-metal trichalcogenide monolayers, *Phys. Rev. B* 91(23), 235425 (2015)
30. J. U. Lee, S. Lee, J. H. Ryoo, S. Kang, T. Y. Kim, P. Kim, C. H. Park, J. Park, and H. Cheong, Ising-type magnetic ordering in atomically thin FePS<sub>3</sub>, *Nano Lett.* 16(12), 7433 (2016)
31. C. T. Kuo, M. Neumann, K. Balamurugan, H. J. Park, S. Kang, H. W. Shiu, J. H. Kang, B. H. Hong, M. Han, T. W. Noh, and J. G. Park, Exfoliation and Raman spectroscopic fingerprint of few-layer NiPS<sub>3</sub> van der Waals crystals, *Sci. Rep.* 6(1), 20904 (2016)
32. Y. G. Wang, J. J. Ying, Z. Y. Zhou, J. L. Sun, T. Wen, Y. N. Zhou, N. N. Li, Q. Zhang, F. Han, Y. M. Xiao, P. Chow, W. G. Yang, V. V. Struzhkin, Y. S. Zhao, and H. K. Mao, Emergent superconductivity in an iron-based honeycomb lattice initiated by pressure-driven spin-crossover, *Nat. Commun.* 9(1), 1914 (2018)
33. Y. H. Gu, Q. Zhang, C. C. Le, Y. X. Li, T. Xiang, and J. P. Hu, Ni-based transition metal trichalcogenide monolayer: A strongly correlated quadruple-layer graphene, *Phys. Rev. B* 100(16), 165405 (2019)
34. H. Yao and F. Yang, Topological odd-parity superconductivity at type-II two-dimensional van Hove singularities, *Phys. Rev. B* 92(3), 035132 (2015)
35. R. Nandkishore, L. S. Levitov, and A. V. Chubukov, Chiral superconductivity from repulsive interactions in doped graphene, *Nat. Phys.* 8(2), 158 (2012)
36. W. S. Wang, Y. Y. Xiang, Q. H. Wang, F. Wang, F. Yang, and D. H. Lee, Functional renormalization group and variational Monte Carlo studies of the electronic instabilities in graphene near 1/4 doping, *Phys. Rev. B* 85(3), 035414 (2012)
37. M. L. Kiesel, C. Platt, W. Hanke, D. A. Abanin, and R. Thomale, Competing many-body instabilities and unconventional superconductivity in graphene, *Phys. Rev. B* 82(2), 020507 (2012)
38. T. X. Ma, F. Yang, H. Yao, and H. Q. Lin, Possible triplet *p+ip* superconductivity in graphene at low filling, *Phys. Rev. B* 90, 245114 (2014)
39. Z. Y. Meng, F. Yang, K. S. Chen, H. Yao, and H. Y. Kee, Evidence for spin-triplet odd-parity superconductivity close to type-II van Hove singularities, *Phys. Rev. B* 91(18), 184509 (2015)
40. X. Chen, Y. G. Yao, H. Yao, F. Yang, and J. Ni, Topological *p+ip* superconductivity in doped graphene-like single-sheet materials BC<sub>3</sub>, *Phys. Rev. B* 92(17), 174503 (2015)
41. L. D. Zhang, F. Yang, and Y. G. Yao, Itinerant ferromagnetism and *p+ip* superconductivity in doped bilayer silicene, *Phys. Rev. B* 92(10), 104504 (2015)
42. C. C. Liu, L. D. Zhang, W. Q. Chen, and F. Yang, Chiral spin density wave and *d+id* superconductivity in the magic-angle-twisted bilayer graphene, *Phys. Rev. Lett.* 121(21), 217001 (2018)
43. H. Isobe, N. F. Q. Yuan, and L. Fu, Unconventional superconductivity and density waves in twisted bilayer graphene, *Phys. Rev. X* 8(4), 041041 (2018)
44. D. Di Sante, X. Wu, M. Fink, W. Hanke, and R. Thomale, Triplet superconductivity in the Dirac semimetal germanene on a substrate, *Phys. Rev. B* 99(20), 201106 (2019)
45. X. X. Wu, M. Fink, W. Hanke, R. Thomale, and D. Di Sante, Unconventional superconductivity in a doped quantum spin Hall insulator, *Phys. Rev. B* 100(4), 041117 (2019)
46. Y. X. Li, X. X. Wu, Y. H. Gu, C. C. Le, S. S. Qin, R. Thomale, and J. P. Hu, Topological superconductivity in Ni-based transition metal trichalcogenides, *Phys. Rev. B* 100, 214513 (2019)

47. S. Graser, T. A. Maier, P. J. Hirschfeld, and D. J. Scalapino, Near-degeneracy of several pairing channels in multiorbital models for the Fe pnictides, *New J. Phys.* 11(2), 025016 (2009)
48. A. F. Kemper, T. A. Maier, S. Graser, H. P. Cheng, P. J. Hirschfeld, and D. J. Scalapino, Sensitivity of the superconducting state and magnetic susceptibility to key aspects of electronic structure in ferropnictides, *New J. Phys.* 12(7), 073030 (2010)
49. X. X. Wu, F. Yang, C. C. Le, H. Fan, and J. P. Hu, Triplet  $p_z$ -wave pairing in quasi-one-dimensional  $A_2Cr_3As_3$  superconductors ( $A = K, Rb, Cs$ ), *Phys. Rev. B* 92(10), 104511 (2015)
50. L. D. Zhang, F. Yang, and Y. G. Yao, Possible electric-field-induced superconducting states in doped silicene, *Sci. Rep.* 5(1), 8203 (2015)
51. Y. T. Kang, C. Lu, F. Yang, and D. X. Yao, Single-orbital realization of high-temperature  $s^\pm$  superconductivity in the square-octagon lattice, *Phys. Rev. B* 99(18), 184506 (2019)
52. L. D. Zhang, X. X. Wu, H. Fan, F. Yang, and J. P. Hu, Revisitation of superconductivity in  $K_2Cr_3As_3$  based on the six-band model, *Europhys. Lett.* 113(3), 37003 (2016)
53. F. Liu, C. C. Liu, K. H. Wu, F. Yang, and Y. G. Yao,  $d+id$  chiral superconductivity in bilayer silicene, *Phys. Rev. Lett.* 111(6), 066804 (2013)