

REVIEW

# Role of tegument proteins in herpesvirus assembly and egress

Haitao Guo<sup>1,2</sup>, Sheng Shen<sup>1,2</sup>, Lili Wang<sup>1,2</sup>, Hongyu Deng<sup>1</sup>✉

<sup>1</sup> CAS Key Laboratory of Infection and Immunity, Institute of Biophysics, Chinese Academy of Sciences, Beijing 100101, China

<sup>2</sup> Graduate School of the Chinese Academy of Sciences, Beijing 100080, China

✉ Correspondence: rrain6@yahoo.com

Received October 4, 2010 Accepted November 4, 2010

## ABSTRACT

**Morphogenesis and maturation of viral particles is an essential step of viral replication. An infectious herpesviral particle has a multilayered architecture, and contains a large DNA genome, a capsid shell, a tegument and an envelope spiked with glycoproteins. Unique to herpesviruses, tegument is a structure that occupies the space between the nucleocapsid and the envelope and contains many virus encoded proteins called tegument proteins. Historically the tegument has been described as an amorphous structure, but increasing evidence supports the notion that there is an ordered addition of tegument during virion assembly, which is consistent with the important roles of tegument proteins in the assembly and egress of herpesviral particles. In this review we first give an overview of the herpesvirus assembly and egress process. We then discuss the roles of selected tegument proteins in each step of the process, i.e., primary envelopment, deenvelopment, secondary envelopment and transport of viral particles. We also suggest key issues that should be addressed in the near future.**

**KEYWORDS** herpesvirus, tegument, assembly, egress, transport


## INTRODUCTION

The family *Herpesviridae* consists of three subfamilies including *Alphaherpesvirinae*, *Betaherpesvirinae* and *Gammapherpesvirinae* whose members infect a broad range of animal species including mammals, birds and reptiles (Davison et al., 2009). Until now, more than 130 herpesviruses have been identified, among which there are eight

human pathogens, including the *Alphaherpesvirinae* members herpes simplex virus type 1 (HSV-1), herpes simplex virus type 2 (HSV-2) and varicella zoster virus (VZV), the *Betaherpesvirinae* members human cytomegalovirus (HCMV), human herpesvirus type 6 (HHV-6) and human herpesvirus type 7 (HHV-7), and the *Gammapherpesvirinae* members Epstein-Barr virus (EBV) and Kaposi's sarcoma-associated herpesvirus (KSHV). HSV most commonly cause mucocutaneous infections, resulting in recurrent orolabial or genital lesions (Roizman et al., 2007). HCMV infection is responsible for approximately 8% of infectious mononucleosis cases and is also associated with inflammatory and proliferative diseases (Söderberg-Nauclér, 2006; Steiner, 2007). EBV and KSHV are associated with several malignancies, such as Burkitt's lymphoma and Kaposi's sarcoma (Chang et al., 1994; Kutok and Wang, 2006).

All members of the *Herpesviridae* family comprise two distinct stages in their life cycle: lytic replication and latency (Sunil-Chandra et al., 1992; Decker et al., 1996a, b; Dupin et al., 1999; Flaño et al., 2000). Members of the *Alphaherpesvirinae* subfamily establish latency in neurons, members of the *Betaherpesvirinae* subfamily establish latency in a range of nonneuronal cells, whereas members of *Gammapherpesvirinae* subfamily members establish latency mainly in B and T lymphocytes. Latency provides the viruses with advantages to escape host immune surveillance so as to establish lifelong persistent infection and thus contributes to transformation and development of malignancies. However, it is through lytic replication that viruses propagate and transmit among hosts to maintain viral reservoirs. Both viral latency and lytic replication play important roles in herpesvirus pathogenesis.

All herpesvirions have a characteristic multilayered architecture (Fig. 1). An infectious virion contains a double-stranded DNA genome, an icosahedral capsid shell, a thick, proteinaceous tegument compartment, and a lipid bilayer



**Figure 1. Diagram of herpesvirion structure.** The virion particle is approximately 200 nm in diameter. A linear double-stranded DNA genome is packaged within an icosahedral capsid to form the nucleocapsid. The capsid is embedded in a matrix known as the tegument layer, which contains many virus coded proteins. The tegument is itself surrounded by the envelope, a lipid membrane containing several viral glycoproteins.

envelope spiked with glycoproteins (Rixon, 1993; Roizman and Pellett, 2001; Liu and Zhou, 2006; Dai et al., 2008). As a unique structure of herpesviruses, the tegument plays important roles in multiple aspects of the viral life cycle, including translocation of nucleocapsids into the nucleus as well as virion assembly and egress (Subak-Sharpe and Dargan, 1998; Fuchs et al., 2002a). As components that are carried into a host cell during viral infection, tegument proteins can also transactivate the expression of viral immediate-early genes and cellular genes as well as modulate, host signal transduction and innate immunity (Bresnahan and Shenk, 2000; Castillo and Kowalik, 2002; Ishov et al., 2002; Zhu et al., 2002).

Virion assembly and egress is an essential stage of herpesviral lytic replication, hence contributes indirectly to herpesvirus pathogenesis. In this review we summarize the growing range of functions attributed to tegument proteins during the course of herpesvirus assembly and egress. We mainly focus on tegument proteins from the *Alphaherpesvirinae* subfamily, in particular HSV-1 and the related porcine pathogen pseudorabies virus (PrV), since they are the most extensively studied. Meanwhile, some well studied tegument proteins from the *Betaherpesvirinae* and *Gammaherpesvirinae* subfamilies are also discussed.

## OVERVIEW OF HERPESVIRUS ASSEMBLY AND EGRESS

During herpesvirus lytic infection, after attachment and penetration, capsids are transported to the nucleus via interaction with microtubules, docking at the nuclear pore where the viral genome is released into the nucleus (Döhner et al., 2002; Wolfstein et al., 2006), where transcription of viral immediate-early, early and late genes and genome replication take place. Then virion assembly and egress proceed from the nucleus to the cytoplasm. Several models have been

proposed to explain this process, among which the most widely accepted is the envelopment-deenvelopment-reenvelopment model: capsid formation and viral DNA packaging in the nucleus, primary envelopment at the inner nuclear membrane and deenvelopment at the outer nuclear membrane, association with tegument proteins and secondary envelopment (i.e., reenvelopment) in the cytoplasm, and budding of complete virions into Golgi body-derived compartments for egress via the cellular secretory pathway (Mettenleiter, 2002, 2004). In each stage of virion assembly and egress, tegument proteins are found to play important roles. There are also lines of evidence suggesting direct exit from the nucleus of unenveloped nucleocapsids via enlarged nuclear pores or transport of perinuclear enveloped nucleocapsids through a continuum of outer nuclear, rough ER and Golgi membranes (Wild et al., 2002; Leuzinger et al., 2005; Wild et al., 2005, 2009). In this review, the functional roles of tegument proteins will be discussed in the context of the envelopment-deenvelopment-reenvelopment model of virion assembly.

## TEGUMENT PROTEINS

Composition of tegument in all three *Herpesviridae* subfamilies has been studied. At least 26, 38 or 17 virus-encoded tegument components are recruited into an HSV-1, HCMV or EBV virion, respectively (Johannsen et al., 2004; Varnum et al., 2004; Zhu et al., 2005; Loret et al., 2008). Some of these proteins, such as pUL36, pUL47, pUL48 and pUL49, are classified as major, structurally significant components (Heine et al., 1974), whereas some, such as vhs and the protein kinase pUL13, are minor but nonetheless important components (Schek and Bachenheimer, 1985; Overton et al., 1992). Some tegument proteins are conserved in all herpesvirus members (Table 1), and their functions are progressively unrevealed.

As an indispensable part of mature virion, tegument plays important roles in herpesviral particle production. Much effort has been placed on elucidating the molecular interactions between various tegument proteins with the hope of understanding the mechanisms by which this complex virus is assembled. Tegument addition is thought to originate in the nucleus, then more tegument proteins are added to capsids in the cytoplasm following nuclear egress and finally at trans-Golgi network (TGN)-derived membranes during maturation budding (Mettenleiter, 2006). The complexity of virion assembly route clearly raises the issue that different tegument proteins may be incorporated into herpesviral particles at different stages of egress and that the composition of virions in the perinuclear space may not be the same as that of the mature extracellular viruses. For example, the UL31 and UL34 gene products of PrV are present in viral particles in the perinuclear space but are absent from virions at later stages of assembly and mature virions (Fuchs et al., 2002c). In contrast, the pUL36, pUL37, pUL46, pUL47, pUL48 and

**Table 1** Conservation of herpesvirus tegument proteins

HSV-1 tegument proteins	nomenclature of HSV-1 tegument homologs		
	HCMV	EBV	KSHV
UL7	UL103	BBRF2	ORF42
UL11	UL99	BBLF1	ORF38
UL13	UL97	BGLF4	ORF36
UL14	UL95	BGLF3	ORF34
UL16	UL94	BGLF2	ORF33
UL21	UL87	BcRF1	ORF24
UL23	NA	BXLF2	ORF21
UL36	UL48	BPLF1	ORF64
UL37	UL47	BOLF1	ORF63
UL51	UL71	BSRF1	ORF55
UL41	NA	NA	NA
UL46	NA	NA	NA
UL47	NA	NA	NA
UL48	NA	NA	NA
UL49	NA	NA	NA
UL50	NA	NA	NA
UL55	NA	NA	NA
US2	NA	NA	NA
US3	NA	NA	NA
US10	NA	NA	NA
US11	NA	NA	NA
RL1 (ICP34.5)	NA	NA	NA
RL2 (ICP0)	NA	NA	NA
RS1 (ICP4)	NA	NA	NA

NA: not applicable; HSV: herpes simplex virus.

pUL49 tegument proteins of PrV are found in only cytoplasmic enveloped virions and mature PrV particles and are therefore presumably not recruited into the virion during primary envelopment (Fuchs et al., 2002a; Klupp et al., 2002; Kopp et al., 2002). The PrV pUS3 protein kinase, on the other hand, is a component of both primary and mature enveloped virions (Granzow et al., 2004). However, some researchers showed conflicting data on the related HSV-1 homologs, in which HSV tegument proteins pUL36, pUL37, pUL48 and pUL49 have been found associated with intranuclear capsids or existing in primary enveloped virions (Mossman et al., 2000; Naldinho-Souto et al., 2006; Bucks et al., 2007; Padula et al., 2009). These results indicate that the sequence of tegument protein addition and the regions in which capsid acquires the tegument are very complicated. The inner tegument proteins may interact with the capsid while the outer tegument proteins may be recruited by interacting with the cytoplasmic tails of viral glycoproteins. Several molecular interactions have been implicated in linking the viral capsid, tegument and membrane during the envelopment process. Examples include pUL25

(capsid)-pUL36 (tegument), pUL49 (tegument)-gD (membrane), pUL48 (tegument)-gH (membrane), and pUL11 (membrane)-pUL16 (tegument/capsid) interactions (McNab et al., 1998; Mettenleiter, 2002, 2004; Collier et al., 2007). These interactions will be elaborated in the following sections.

## PRIMARY ENVELOPMENT AND DEENVELOPMENT

After capsid assembly, the nucleocapsids will bud at the inner nuclear membrane (NM). There, the viruses encounter the nuclear lamina, a rigid network of lamin proteins lining the inner NM, which is a major obstacle for budding of the nucleocapsids. Herpesviruses disrupt the nuclear lamina in order to bud into the perinuclear space and acquire the inner NM as the envelope, the so called primary envelopment (Scott and O'Hare, 2001; Muranyi et al., 2002; Leach et al., 2007; Mou et al., 2007; Mou et al., 2008). HSV-1 and PrV disrupt the lamina through two viral proteins, pUL31 and pUL34, which form a complex colocalized to the inner NM (Reynolds et al., 2002; Reynolds et al., 2004; Simpson-Holley et al., 2004; Simpson-Holley et al., 2005; Bjerke and Roller, 2006). Research suggested that pUL31 and pUL34 promote viral nuclear egress by several mechanisms, including (1) affecting maturation of viral replication intermediates so that capsids assemble adjacent to the nuclear envelope, making it convenient for primary egress (Simpson-Holley et al., 2004, 2005), (2) causing displacement of and conformational changes in lamins A/C and B (Reynolds et al., 2004; Simpson-Holley et al., 2004; Bjerke and Roller, 2006), and (3) mislocalizing the membrane lamin receptors, such as the lamin B receptor and emerin, which tether lamins to the inner NM (Scott and O'Hare, 2001; Leach et al., 2007; Morris et al., 2007; Mou et al., 2008). Amazingly, expression of PrV pUL31 and pUL34 in RK-13 cells resulted in formation of vesicles in the perinuclear space, indicating that they may compose the essential budding machinery (Klupp et al., 2007).

pUL31 and pUL34 are conserved in all three *Herpesviridae* subfamilies. In EBV, the pUL34 positional homolog BBRF1 and the pUL31 positional homolog BFLF2 also interact with each other. HSV-1 UL34 or EBV BBRF1 mutant viruses showed a phenotype of nuclear retention (Lake and Hutt-Fletcher, 2004; Farina et al., 2005; Gonnella et al., 2005; Calderwood et al., 2007). EBV BFLF2 functions in both viral DNA packaging and primary envelopment, as deletion of BFLF2 led to sequestration of nucleocapsids in the nucleus and production of more empty capsids without viral genomes (Granato et al., 2008).

Alphaherpesviruses also express a viral serine/threonine protein kinase, pUS3, that phosphorylates lamins and lamin receptor emerin, which are thought to be involved in maintaining nuclear integrity (Leach et al., 2007; Morris et al., 2007; Mou et al., 2008). In addition, both HSV-1 and PrV pUL31 and pUL34 have been shown to be phosphorylated in a pUS3-dependent manner, and all three proteins have been found to be incorporated into primary virions (Purves et al.,

1991; Granzow et al., 2004; Ryckman and Roller, 2004; Kato et al., 2005; Mou et al., 2007, 2009). pUS3 also regulates the localization of the pUL31/pUL34 complex, as deletion of pUS3 disrupts their localization from a smooth nuclear rim distribution to distinct foci (Wagenaar et al., 1995; Klupp et al., 2001a; Reynolds et al., 2002; Ryckman and Roller, 2004). These observations indicate that pUS3 might function in virion primary envelopment.

The other tegument protein kinase, pUL13, may also play a role in regulating the localization of pUL31 and pUL34 in HSV-1 either directly or indirectly through phosphorylation of pUS3 (Kato et al., 2006, 2008), indicating its participating in primary envelopment. The HCMV homolog of HSV-1 pUL13, pUL97, which is also a kinase, is incorporated into virions (Michel et al., 1996). pUL97 has been reported to play an important role in virion nuclear egress, as a marked decrease in the number of mature capsids was observed in the cytoplasm of cells infected with UL97 null mutant compared to the wild type virus (Wolf et al., 2001; Krosky et al., 2003). Deletion of HCMV UL97 causes an aggregation of tegument and other structural proteins in nuclear inclusion, suggesting that pUL97 kinase is required for the normal function of tegument proteins in the nucleus and may function in initiation of tegumentation or primary envelopment (Prichard et al., 2005). Although it is not clear where pUL97 is added onto the nucleocapsid, interaction between pUL97 and the HCMV major tegument protein pp65 is required for the incorporation of pUL97 into viral particles (Kamil and Coen, 2007; Chevillotte et al., 2009).

HSV-1 tegument proteins pUL36, pUL37, pUL48 and pUL49 have been found to be associated with intranuclear capsids or exist in primary enveloped virions, though their specific functions in nuclear egress are not clear yet (Mossman et al., 2000; Naldinho-Souto et al., 2006; Bucks et al., 2007; Padula et al., 2009). pUL36 contains a capsid binding domain and interacts with a minor capsid protein pUL25 previously found to participate in DNA encapsidation (McNab et al., 1998; Coller et al., 2007). Such an interaction is required for assembly of pUL36 onto capsid, indicating that pUL36 can bind to capsid directly. This result is consistent with those from electron microscope studies, which suggest presence of HSV-1 pUL36 in the inner most layer of the tegument, attached to the vertices of the capsid (McNabb and Courtney, 1992; Zhou et al., 1999). pUL36 is conserved throughout the *Herpesviridae*. It forms a complex with pUL37 as part of the capsid-associated inner tegument. Deletion of either of the genes encoding pUL36 or pUL37 blocks further tegumentation of the capsid in the cytoplasm (Desai, 2000; Desai et al., 2001; Klupp et al., 2001b; Fuchs et al., 2004; Leege et al., 2009; Roberts et al., 2009). By interacting with pUL48, pUL36 is a major factor for recruiting pUL48 onto capsids (Ko et al., 2010). However, in PrV, pUL36 and pUL48 only exist in mature virions, and are undetectable from nuclear or primary enveloped virions, indicating that these two tegument proteins are acquired in the cytoplasm

(Naldinho-Souto et al., 2006; Möhl et al., 2009). Deletion of pUL48 does not affect the capsid assembly and nuclear egress of PrV (Fuchs et al., 2002a). These conflicting results suggest that pUL36 and pUL48 of HSV-1 and PrV may function differently.

Deenvelopment is a step of herpesvirus nuclear egress whose details are poorly understood. pUL48 might play a role in this process for HSV-1, as large numbers of enveloped particles accumulate between the inner and outer NM in the absence of pUL48 (Mossman et al., 2000). In addition, HSV-1 gB and gH/gL have redundant or overlapping roles in fusion with the outer NM (Farnsworth et al., 2007b). Consistent with this notion, EBV, KSHV and PrV gB null mutants all exhibit defects in nuclear egress (Peeters et al., 1992; Lee and Longnecker, 1997; Krishnan et al., 2005). In the primary enveloped virions, pUS3 can phosphorylate the gB cytoplasmic domain and this modification is important for gB-mediated fusion of the virion envelope and the outer NM (Wisner et al., 2009). Thus, pUS3 plays important roles in both primary envelopment and deenvelopment.

## SECONDARY ENVELOPMENT

Most tegument proteins are recruited onto the nucleocapsid in the cytoplasm. The addition of outer tegument is accompanied with virion secondary envelopment. During this process, several molecular interactions, such as pUL11 (membrane)-pUL16 (tegument), pUL48 (tegument)-gH (membrane) and pUL49 (tegument)-gE (membrane), have been implicated in linking the viral capsid, tegument and membrane (Mettenleiter, 2002, 2004). pUL16 is conserved throughout the *Herpesviridae*, and plays a pivotal role in virion morphogenesis in the cytoplasm. pUL16 binds to the capsid by interacting with pUL21 which is also a tegument protein associated with capsid in both the nucleus and the cytoplasm (de Wind et al., 1992; Takakuwa et al., 2001; Klupp et al., 2005). pUL16 has another binding partner, named pUL11, which is a membrane-bound tegument protein (Loomis et al., 2003). Two distinct binding sites exist in pUL16 for pUL11 and pUL21, as covalent modification of its free cysteines with *N*-ethylmaleimide blocks its binding to pUL11 but not to pUL21 (Yeh et al., 2008; Harper et al., 2010). This observation suggests that pUL16 may interact with pUL21 and pUL11 in sequence and both interactions contribute to efficient incorporation of pUL16 into virions (Meckes et al., 2010). The pUL11 tegument protein is also conserved among all herpesviruses, and each homolog contains amino acid motifs that allow covalent modifications with two fatty acids, myristate and palmitate (MacLean et al., 1989; Loomis et al., 2001). These two modifications allow pUL11 accumulating at TGN-derived vesicles in the absence of other viral proteins (Loomis et al., 2001). In addition, there exist leucine-isoleucine and acidic cluster motifs, responsible for recovering pUL11 from plasma membrane to the internal

membrane. This might enable the recruitment of other membrane proteins (virus or host encoded) from the plasma membrane for secondary envelopment occurring around the TGN-vesicles (Loomis et al., 2001).

Similar to the conservation of pUL16 and pUL11, their interaction is also conserved among all herpesviruses. In HCMV, the pUL11 homolog pUL99 has been found to interact with the pUL16 homolog pUL94 (Liu et al., 2009). Consistent with the previously described model of capsid transport and budding, herpesviruses lacking pUL16, pUL21 or pUL11 (or their homologs) have defects in virus egress, resulting in decreased amounts of extracellular viruses produced and the accumulation of non-enveloped capsids in the cytoplasm (MacLean et al., 1989, 1992; Baines and Roizman, 1992; Baines et al., 1994; Kopp et al., 2003; Schimmer and Neubauer, 2003; Silva et al., 2003; Britt et al., 2004; Jones and Lee, 2004; Silva et al., 2005; Seo and Britt, 2006; Guo et al., 2009).

In murine gammaherpesvirus-68 (MHV-68), the homolog of pUL16 is ORF33. Our laboratory has confirmed that ORF33 indeed encodes a tegument protein for MHV-68. Through constructing and analyzing an ORF33 null mutant, we demonstrated that ORF33 is not required for viral genome replication or gene expression, but is essential for virion morphogenesis and egress. More interestingly, ORF33 null mutation caused a partial retention of nucleocapsids in the nucleus, and maturation of virions was arrested at a cytoplasmic stage of partially tegumented nucleocapsids, indicating that ORF33 participates in both primary envelopment of nucleocapsids and morphogenesis of virions in the cytoplasm (Guo et al., 2009). We also found that ORF33 localizes in the nucleus during early infection and interacts with another nuclear tegument protein (Guo et al., unpublished data). We hypothesize that such interaction may facilitate the nuclear budding or egress of nucleocapsids, although such an interaction is relatively redundant.

A number of interactions have been identified involving HSV-1 pUL48 and inner tegument proteins, outer tegument proteins or the cytoplasmic tails of glycoproteins, supporting the concept that pUL48 contributes to linking capsid and envelope during virion formation. For instance, pUL48 has been shown to interact with the outer tegument proteins pUL41, pUL46, pUL47 and pUL49, through a combination of yeast two-hybrid, *in vitro* pull-down and co-immunoprecipitation studies (Smibert et al., 1994; Elliott et al., 1995; Vittone et al., 2005). HSV-2 pUL48 has also been shown to copurify and colocalize with pUL46 during the course of infection (Kato et al., 2000). HSV-1 pUL46 is associated with cellular membrane independent of the presence of other viral proteins. Surprisingly, membrane association within the cell does not translate to association of this protein with the viral envelope, as pUL46 exhibits a strong affinity for the capsid of purified viral particles (Murphy et al., 2008). However, deletion of pUL46 in HSV-1 and PrV does not block virion assembly

(Zhang et al., 1991; Kopp et al., 2002). Although deletion of pUL47 causes an approximately 10-fold decrease in viral titer, pUL47 is not essential for virion assembly in the cytoplasm (Kopp et al., 2002). HSV-1 pUL49 interacts with pUL48, but this interaction is not required for incorporation of pUL49 into viral particles (O'Regan et al., 2007). Similar to pUL46, deletion of pUL49 in HSV-1 does not have a significant effect on virus assembly (Elliott et al., 2005). In the case of PrV, deletion of any of pUL46, pUL47, pUL48 or pUL49 does not prevent incorporation of the remaining three proteins into the tegument (Fuchs et al., 2002a; Michael et al., 2006). Furthermore, an increased incorporation of cellular actin into virions was observed when pUL46, pUL47 or pUL49 was deleted, suggesting that actin may fill the void left by the absence of these tegument proteins (del Rio et al., 2005; Michael et al., 2006). These observations indicate that redundancy exists in tegument assembly. In addition to tegument proteins mentioned above, HSV-1 pUL48 also interacts with the cytoplasmic tails of HSV-1 glycoprotein gB, gD and gH, as identified by pull-down, co-immunoprecipitation and chemical cross-linking assays (Zhu and Courtney, 1994; Gross et al., 2003; Kamen et al., 2005). However, the role of these interactions in HSV-1 assembly requires further investigation.

pUL49, besides an interaction with pUL48, also interacts with a number of viral envelope proteins. Several studies have shown that pUL49 interacts with the cytoplasmic tails of HSV-1 gE, gD and envelope protein pUS9 (Chi et al., 2005; Farnsworth et al., 2007a; O'Regan et al., 2007). In PrV, interactions of pUL49 with the cytoplasmic tail of both gE and gM were identified (Fuchs et al., 2002b). Furthermore, for both HSV-1 and PrV, gE or gM is sufficient for recruitment of pUL49 into virions through a direct interaction (Fuchs et al., 2002b; Michael et al., 2006; O'Regan et al., 2007; Stylianou et al., 2009).

Other tegument proteins promoting secondary envelopment have also been identified. ORF52 is a tegument protein conserved in gammaherpesviruses. The MHV-68 ORF52 localizes in TGN-derived vesicles in both transfection and viral infection. Furthermore, ORF52 was found to specifically function in virus budding into the vesicles in the cytoplasm (Bortz et al., 2007). The crystal structure of ORF52 has been determined at 2.1 Å resolution, revealing a dimeric association of this protein. The self-association was confirmed by co-immunoprecipitation and fluorescence resonance energy transfer experiments. Functional complementation assay demonstrated that both the N-terminal  $\alpha$ -helix and the conserved Arg<sup>95</sup> residue are critical for ORF52 function (Benach et al., 2007). The detailed mechanism responsible for involvement of ORF52 in secondary envelopment is currently under investigation in our laboratory.

In HCMV, the tegument protein pUL32 has been reported to be important in cytoplasmic maturation of virions, especially in virus egress (Tandon and Mocarski, 2008). pUL32

interacts with Bicaudal D1 (BicD1), a protein thought to play a role in trafficking within the secretory pathway (Indran et al., 2010). This interaction helps pUL32 to traffic to the virus assembly compartment, suggesting a primary contribution of this protein in the morphogenesis and/or cytoplasmic transport of progeny virion particles to sites of virion secondary envelopment.


## TRANSPORT AND RELEASE

How do nucleocapsids or partially tegumented nucleocapsids transport to the site where secondary envelopment takes place? Microtubule has been found to be involved in transporting non-enveloped viral particles. Several tegument proteins have been identified as candidates linking deenveloped nucleocapsids to the anterograde cellular microtubule-dependent molecular motor kinesin. In alphaherpesviruses, pUS11 was found to interact with both kinesin-1 and the kinesin-related protein PAT-1 (Diefenbach et al., 2002; Benboudjema et al., 2003). An association between pUL21 and microtubules was also detected (Takakuwa et al., 2001), raising the possibility that pUL21 could be involved with capsid transport to the TGN-derived vesicles, where pUL16 interacts with pUL11, facilitating the budding process by linking capsids to the membrane. In gammaherpesviruses, an interaction between KSHV tegument protein ORF45

and kinesin-2, determined by yeast two-hybrid and co-immunoprecipitation assays, has been reported (Sathish et al., 2009). Disrupting the interaction between KSHV ORF45 and kinesin-2 by a dominant negative mutant of ORF45 leads to a decreased final production of mature viruses. This result supports the idea that the association between a viral tegument protein and a cellular molecular motor is important for the intracellular movement of deenveloped nucleocapsids. The HSV-1 pUL36 is a multifunctional large tegument protein. As part of the inner tegument, pUL36 is also important for the transport of enveloped virions. A potential mechanism was proposed that, during viral infection, pUL36 resides on the surface of organelles which nucleocapsids bud into, and pUL36 would directly or indirectly recruit kinesin motors to the organelles to enable motility (Shanda and Wilson, 2008).

## CONCLUSIONS AND PERSPECTIVES

Through interacting with and manipulating the host micro-environment, tegument proteins are multifunctional during the complete cycle of herpesvirus lytic replication. In particular, they play key structural roles during virion primary envelopment, secondary envelopment and virion trafficking by forming a bridge between capsid or capsid-associated proteins and membrane-associated viral proteins or cellular



**Figure 2. Diagram summarizing the roles of selected tegument proteins in herpesvirus assembly and egress.** Viral genomes are packaged into preformed capsids in the nucleus. Through primary envelopment at the inner nuclear membrane and deenvelopment at the outer nuclear membrane, the nucleocapsid is transported into the cytoplasm from the nucleus. Acquisition of tegument proteins onto the nucleocapsid is completed in the cytoplasm. The immature virion is transported via the microtubules into a trans-Golgi derived vesicle containing viral glycoproteins. After transport of the vesicle to the cell surface, the vesicle and plasma membrane fuse, resulting in the egress of a mature, enveloped virion.

molecular motors (Fig. 2). The herpesvirion assembly and egress process is complex and dynamic, and so are the roles of tegument proteins. As a complex process, herpesvirion assembly and egress involves many protein–protein interactions with marked redundancy. Indeed, interactions between tegument proteins and other viral and cellular proteins are increasingly reported *in vitro* or *in vivo*. However, future research on tegument will require identifying herpesviral protein–protein interaction and herpesviral protein–cellular protein interaction maps during the course of infecting host cells and validating such interactions by viral genetics approach. This has been done to some extent with the *Alphaherpesvirinae* subfamily member VZV and the *Gammaherpesvirinae* subfamily members KSHV and EBV (Uetz et al., 2006; Calderwood et al., 2007; Rozen et al., 2008). Such information will improve our understanding of the biology of herpesvirus and the roles of tegument proteins. More importantly, by examining the importance of each pair of protein–protein interaction, useful therapeutic targets may be identified. The current antiviral strategy for treatment of herpesviruses employs inhibitors of viral DNA replication which have varying efficacies depending on the *Herpesvirinae* subfamily being treated. Therefore, new targets for therapeutic interventions through different mechanisms are useful, especially when combined with the viral DNA replication inhibitors.

The herpesvirion assembly and egress is also a dynamic process, and dissecting the roles of tegument proteins in such a dynamic process calls for integration of different technical approaches. Recently, combinations of molecular tags visible in light and electron microscopes have become particularly advantageous in the analysis of dynamic events at the cellular level (Martin et al., 2005; Gaietta et al., 2006; Lanman et al., 2008). Engineering such tags into herpesvirus tegument proteins as well as capsid/envelope proteins will enable optical live cell imaging and correlated ultrastructural analysis by electron microscopy, and help provide high-resolution information for the dynamic process of herpesvirion assembly and egress.

## ACKNOWLEDGEMENTS

We thank Dr. Z. Hong Zhou and members of the Deng' laboratory for helpful discussions. This work was supported by the "One Hundred Talents Program" from the Chinese Academy of Sciences and the National Protein Science Project (No. 2006CB910901) from the Ministry of Science and Technology.

## REFERENCES

Baines, J.D., Koyama, A.H., Huang, T., and Roizman, B. (1994). The UL21 gene products of herpes simplex virus 1 are dispensable for growth in cultured cells. *J Virol* 68, 2929–2936.

Baines, J.D., and Roizman, B. (1992). The UL11 gene of herpes simplex virus 1 encodes a function that facilitates nucleocapsid

envelopment and egress from cells. *J Virol* 66, 5168–5174.

Benach, J., Wang, L., Chen, Y., Ho, C.K., Lee, S., Seetharaman, J., Xiao, R., Acton, T.B., Montelione, G.T., Deng, H., *et al.* (2007). Structural and functional studies of the abundant tegument protein ORF52 from murine gammaherpesvirus 68. *J Biol Chem* 282, 31534–31541.

Benboudjema, L., Mulvey, M., Gao, Y., Pimplikar, S.W., and Mohr, I. (2003). Association of the herpes simplex virus type 1 Us11 gene product with the cellular kinesin light-chain-related protein PAT1 results in the redistribution of both polypeptides. *J Virol* 77, 9192–9203.

Bjerke, S.L., and Roller, R.J. (2006). Roles for herpes simplex virus type 1 UL34 and US3 proteins in disrupting the nuclear lamina during herpes simplex virus type 1 egress. *Virology* 347, 261–276.

Bortz, E., Wang, L., Jia, Q., Wu, T.T., Whitelegge, J.P., Deng, H., Zhou, Z.H., and Sun, R. (2007). Murine gammaherpesvirus 68 ORF52 encodes a tegument protein required for virion morphogenesis in the cytoplasm. *J Virol* 81, 10137–10150.

Bresnahan, W.A., and Shenk, T.E. (2000). UL82 virion protein activates expression of immediate early viral genes in human cytomegalovirus-infected cells. *Proc Natl Acad Sci U S A* 97, 14506–14511.

Britt, W.J., Jarvis, M., Seo, J.Y., Drummond, D., and Nelson, J. (2004). Rapid genetic engineering of human cytomegalovirus by using a lambda phage linear recombination system: demonstration that pp28 (UL99) is essential for production of infectious virus. *J Virol* 78, 539–543.

Bucks, M.A., O'Regan, K.J., Murphy, M.A., Wills, J.W., and Courtney, R.J. (2007). Herpes simplex virus type 1 tegument proteins VP1/2 and UL37 are associated with intranuclear capsids. *Virology* 361, 316–324.

Calderwood, M.A., Venkatesan, K., Xing, L., Chase, M.R., Vazquez, A., Holthaus, A.M., Ewence, A.E., Li, N., Hirozane-Kishikawa, T., Hill, D.E., *et al.* (2007). Epstein-Barr virus and virus human protein interaction maps. *Proc Natl Acad Sci U S A* 104, 7606–7611.

Castillo, J.P., and Kowalik, T.F. (2002). Human cytomegalovirus immediate early proteins and cell growth control. *Gene* 290, 19–34.

Chang, Y., Cesarman, E., Pessin, M.S., Lee, F., Culpepper, J., Knowles, D.M., and Moore, P.S. (1994). Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science* 266, 1865–1869.

Chevillotte, M., Landwehr, S., Linta, L., Frascaroli, G., Lüske, A., Buser, C., Mertens, T., and von Einem, J. (2009). Major tegument protein pp65 of human cytomegalovirus is required for the incorporation of pUL69 and pUL97 into the virus particle and for viral growth in macrophages. *J Virol* 83, 2480–2490.

Chi, J.H., Harley, C.A., Mukhopadhyay, A., and Wilson, D.W. (2005). The cytoplasmic tail of herpes simplex virus envelope glycoprotein D binds to the tegument protein VP22 and to capsids. *J Gen Virol* 86, 253–261.

Coller, K.E., Lee, J.I., Ueda, A., and Smith, G.A. (2007). The capsid and tegument of the alphaherpesviruses are linked by an interaction between the UL25 and VP1/2 proteins. *J Virol* 81, 11790–11797.

Dai, W., Jia, Q., Bortz, E., Shah, S., Liu, J., Atanasov, I., Li, X., Taylor, K.A., Sun, R., and Zhou, Z.H. (2008). Unique structures in a tumor herpesvirus revealed by cryo-electron tomography and microscopy. *J Struct Biol* 161, 428–438.

- Davison, A.J., Eberle, R., Ehlers, B., Hayward, G.S., McGeoch, D.J., Minson, A.C., Pellett, P.E., Roizman, B., Studdert, M.J., and Thiry, E. (2009). The order Herpesvirales. *Arch Virol* 154, 171–177.
- de Wind, N., Wagenaar, F., Pol, J., Kimman, T., and Berns, A. (1992). The pseudorabies virus homology of the herpes simplex virus UL21 gene product is a capsid protein which is involved in capsid maturation. *J Virol* 66, 7096–7103.
- Decker, L.L., Klamon, L.D., and Thorley-Lawson, D.A. (1996a). Detection of the latent form of Epstein-Barr virus DNA in the peripheral blood of healthy individuals. *J Virol* 70, 3286–3289.
- Decker, L.L., Shankar, P., Khan, G., Freeman, R.B., Dezube, B.J., Lieberman, J., and Thorley-Lawson, D.A. (1996b). The Kaposi sarcoma-associated herpesvirus (KSHV) is present as an intact latent genome in KS tissue but replicates in the peripheral blood mononuclear cells of KS patients. *J Exp Med* 184, 283–288.
- del Rio, T., DeCoste, C.J., and Enquist, L.W. (2005). Actin is a component of the compensation mechanism in pseudorabies virus virions lacking the major tegument protein VP22. *J Virol* 79, 8614–8619.
- Desai, P., Sexton, G.L., McCaffery, J.M., and Person, S. (2001). A null mutation in the gene encoding the herpes simplex virus type 1 UL37 polypeptide abrogates virus maturation. *J Virol* 75, 10259–10271.
- Desai, P.J. (2000). A null mutation in the UL36 gene of herpes simplex virus type 1 results in accumulation of unenveloped DNA-filled capsids in the cytoplasm of infected cells. *J Virol* 74, 11608–11618.
- Diefenbach, R.J., Miranda-Saksena, M., Diefenbach, E., Holland, D. J., Boadle, R.A., Armati, P.J., and Cunningham, A.L. (2002). Herpes simplex virus tegument protein US11 interacts with conventional kinesin heavy chain. *J Virol* 76, 3282–3291.
- Döhner, K., Wolfstein, A., Prank, U., Echeverri, C., Dujardin, D., Vallee, R., and Sodeik, B. (2002). Function of dynein and dynactin in herpes simplex virus capsid transport. *Mol Biol Cell* 13, 2795–2809.
- Dupin, N., Fisher, C., Kellam, P., Ariad, S., Tulliez, M., Franck, N., van Marck, E., Salmon, D., Gorin, I., Escande, J.P., *et al.* (1999). Distribution of human herpesvirus-8 latently infected cells in Kaposi's sarcoma, multicentric Castleman's disease, and primary effusion lymphoma. *Proc Natl Acad Sci U S A* 96, 4546–4551.
- Elliott, G., Hafezi, W., Whiteley, A., and Bernard, E. (2005). Deletion of the herpes simplex virus VP22-encoding gene (UL49) alters the expression, localization, and virion incorporation of ICP0. *J Virol* 79, 9735–9745.
- Elliott, G., Mouzakis, G., and O'Hare, P. (1995). VP16 interacts via its activation domain with VP22, a tegument protein of herpes simplex virus, and is relocated to a novel macromolecular assembly in coexpressing cells. *J Virol* 69, 7932–7941.
- Farina, A., Feederle, R., Raffa, S., Gonnella, R., Santarelli, R., Frati, L., Angeloni, A., Torrisi, M.R., Faggioni, A., and Delecluse, H.J. (2005). BFRF1 of Epstein-Barr virus is essential for efficient primary viral envelopment and egress. *J Virol* 79, 3703–3712.
- Farnsworth, A., Wisner, T.W., and Johnson, D.C. (2007a). Cytoplasmic residues of herpes simplex virus glycoprotein gE required for secondary envelopment and binding of tegument proteins VP22 and UL11 to gE and gD. *J Virol* 81, 319–331.
- Farnsworth, A., Wisner, T.W., Webb, M., Roller, R., Cohen, G., Eisenberg, R., and Johnson, D.C. (2007b). Herpes simplex virus glycoproteins gB and gH function in fusion between the virion envelope and the outer nuclear membrane. *Proc Natl Acad Sci U S A* 104, 10187–10192.
- Flaño, E., Husain, S.M., Sample, J.T., Woodland, D.L., and Blackman, M.A. (2000). Latent murine gamma-herpesvirus infection is established in activated B cells, dendritic cells, and macrophages. *J Immunol* 165, 1074–1081.
- Fuchs, W., Granzow, H., Klupp, B.G., Kopp, M., and Mettenleiter, T.C. (2002a). The UL48 tegument protein of pseudorabies virus is critical for intracytoplasmic assembly of infectious virions. *J Virol* 76, 6729–6742.
- Fuchs, W., Klupp, B.G., Granzow, H., Hengartner, C., Brack, A., Mundt, A., Enquist, L.W., and Mettenleiter, T.C. (2002b). Physical interaction between envelope glycoproteins E and M of pseudorabies virus and the major tegument protein UL49. *J Virol* 76, 8208–8217.
- Fuchs, W., Klupp, B.G., Granzow, H., and Mettenleiter, T.C. (2004). Essential function of the pseudorabies virus UL36 gene product is independent of its interaction with the UL37 protein. *J Virol* 78, 11879–11889.
- Fuchs, W., Klupp, B.G., Granzow, H., Osterrieder, N., and Mettenleiter, T.C. (2002c). The interacting UL31 and UL34 gene products of pseudorabies virus are involved in egress from the host-cell nucleus and represent components of primary enveloped but not mature virions. *J Virol* 76, 364–378.
- Gaietta, G.M., Giepmans, B.N., Deerinck, T.J., Smith, W.B., Ngan, L., Llopis, J., Adams, S.R., Tsien, R.Y., and Ellisman, M.H. (2006). Golgi twins in late mitosis revealed by genetically encoded tags for live cell imaging and correlated electron microscopy. *Proc Natl Acad Sci U S A* 103, 17777–17782.
- Gonnella, R., Farina, A., Santarelli, R., Raffa, S., Feederle, R., Bei, R., Granato, M., Modesti, A., Frati, L., Delecluse, H.J., *et al.* (2005). Characterization and intracellular localization of the Epstein-Barr virus protein BFLF2: interactions with BFRF1 and with the nuclear lamina. *J Virol* 79, 3713–3727.
- Granato, M., Feederle, R., Farina, A., Gonnella, R., Santarelli, R., Hub, B., Faggioni, A., and Delecluse, H.J. (2008). Deletion of Epstein-Barr virus BFLF2 leads to impaired viral DNA packaging and primary egress as well as to the production of defective viral particles. *J Virol* 82, 4042–4051.
- Granzow, H., Klupp, B.G., and Mettenleiter, T.C. (2004). The pseudorabies virus US3 protein is a component of primary and of mature virions. *J Virol* 78, 1314–1323.
- Gross, S.T., Harley, C.A., and Wilson, D.W. (2003). The cytoplasmic tail of Herpes simplex virus glycoprotein H binds to the tegument protein VP16 in vitro and in vivo. *Virology* 317, 1–12.
- Guo, H., Wang, L., Peng, L., Zhou, Z.H., and Deng, H. (2009). Open reading frame 33 of a gammaherpesvirus encodes a tegument protein essential for virion morphogenesis and egress. *J Virol* 83, 10582–10595.
- Harper, A.L., Meckes, D.G. Jr, Marsh, J.A., Ward, M.D., Yeh, P.C., Baird, N.L., Wilson, C.B., Semmes, O.J., and Wills, J.W. (2010). Interaction domains of the UL16 and UL21 tegument proteins of herpes simplex virus. *J Virol* 84, 2963–2971.
- Heine, J.W., Honess, R.W., Cassai, E., and Roizman, B. (1974). Proteins specified by herpes simplex virus. XII. The virion polypeptides of type 1 strains. *J Virol* 14, 640–651.
- Indran, S.V., Ballestas, M.E., and Britt, W.J. (2010). Bicaudal D1-dependent trafficking of human cytomegalovirus tegument protein

- pp150 in virus-infected cells. *J Virol* 84, 3162–3177.
- Ishov, A.M., Vladimirova, O.V., and Maul, G.G. (2002). Daxx-mediated accumulation of human cytomegalovirus tegument protein pp71 at ND10 facilitates initiation of viral infection at these nuclear domains. *J Virol* 76, 7705–7712.
- Johannsen, E., Luftig, M., Chase, M.R., Weicksel, S., Cahir-McFarland, E., Illanes, D., Sarracino, D., and Kieff, E. (2004). Proteins of purified Epstein-Barr virus. *Proc Natl Acad Sci U S A* 101, 16286–16291.
- Jones, T.R., and Lee, S.W. (2004). An acidic cluster of human cytomegalovirus UL99 tegument protein is required for trafficking and function. *J Virol* 78, 1488–1502.
- Kamen, D.E., Gross, S.T., Girvin, M.E., and Wilson, D.W. (2005). Structural basis for the physiological temperature dependence of the association of VP16 with the cytoplasmic tail of herpes simplex virus glycoprotein H. *J Virol* 79, 6134–6141.
- Kamil, J.P., and Coen, D.M. (2007). Human cytomegalovirus protein kinase UL97 forms a complex with the tegument phosphoprotein pp65. *J Virol* 81, 10659–10668.
- Kato, A., Tanaka, M., Yamamoto, M., Asai, R., Sata, T., Nishiyama, Y., and Kawaguchi, Y. (2008). Identification of a physiological phosphorylation site of the herpes simplex virus 1-encoded protein kinase Us3 which regulates its optimal catalytic activity in vitro and influences its function in infected cells. *J Virol* 82, 6172–6189.
- Kato, A., Yamamoto, M., Ohno, T., Kodaira, H., Nishiyama, Y., and Kawaguchi, Y. (2005). Identification of proteins phosphorylated directly by the Us3 protein kinase encoded by herpes simplex virus 1. *J Virol* 79, 9325–9331.
- Kato, A., Yamamoto, M., Ohno, T., Tanaka, M., Sata, T., Nishiyama, Y., and Kawaguchi, Y. (2006). Herpes simplex virus 1-encoded protein kinase UL13 phosphorylates viral Us3 protein kinase and regulates nuclear localization of viral envelopment factors UL34 and UL31. *J Virol* 80, 1476–1486.
- Kato, K., Daikoku, T., Goshima, F., Kume, H., Yamaki, K., and Nishiyama, Y. (2000). Synthesis, subcellular localization and VP16 interaction of the herpes simplex virus type 2 UL46 gene product. *Arch Virol* 145, 2149–2162.
- Klupp, B.G., Böttcher, S., Granzow, H., Kopp, M., and Mettenleiter, T.C. (2005). Complex formation between the UL16 and UL21 tegument proteins of pseudorabies virus. *J Virol* 79, 1510–1522.
- Klupp, B.G., Fuchs, W., Granzow, H., Nixdorf, R., and Mettenleiter, T.C. (2002). Pseudorabies virus UL36 tegument protein physically interacts with the UL37 protein. *J Virol* 76, 3065–3071.
- Klupp, B.G., Granzow, H., Fuchs, W., Keil, G.M., Finke, S., and Mettenleiter, T.C. (2007). Vesicle formation from the nuclear membrane is induced by coexpression of two conserved herpesvirus proteins. *Proc Natl Acad Sci U S A* 104, 7241–7246.
- Klupp, B.G., Granzow, H., and Mettenleiter, T.C. (2001a). Effect of the pseudorabies virus US3 protein on nuclear membrane localization of the UL34 protein and virus egress from the nucleus. *J Gen Virol* 82, 2363–2371.
- Klupp, B.G., Granzow, H., Mundt, E., and Mettenleiter, T.C. (2001b). Pseudorabies virus UL37 gene product is involved in secondary envelopment. *J Virol* 75, 8927–8936.
- Ko, D.H., Cunningham, A.L., and Diefenbach, R.J. (2010). The major determinant for addition of tegument protein pUL48 (VP16) to capsids in herpes simplex virus type 1 is the presence of the major tegument protein pUL36 (VP1/2). *J Virol* 84, 1397–1405.
- Kopp, M., Granzow, H., Fuchs, W., Klupp, B.G., Mundt, E., Karger, A., and Mettenleiter, T.C. (2003). The pseudorabies virus UL11 protein is a virion component involved in secondary envelopment in the cytoplasm. *J Virol* 77, 5339–5351.
- Kopp, M., Klupp, B.G., Granzow, H., Fuchs, W., and Mettenleiter, T.C. (2002). Identification and characterization of the pseudorabies virus tegument proteins UL46 and UL47: role for UL47 in virion morphogenesis in the cytoplasm. *J Virol* 76, 8820–8833.
- Krishnan, H.H., Sharma-Walia, N., Zeng, L., Gao, S.J., and Chandran, B. (2005). Envelope glycoprotein gB of Kaposi's sarcoma-associated herpesvirus is essential for egress from infected cells. *J Virol* 79, 10952–10967.
- Krosky, P.M., Baek, M.C., and Coen, D.M. (2003). The human cytomegalovirus UL97 protein kinase, an antiviral drug target, is required at the stage of nuclear egress. *J Virol* 77, 905–914.
- Kutok, J.L., and Wang, F. (2006). Spectrum of Epstein-Barr virus-associated diseases. *Annu Rev Pathol* 1, 375–404.
- Lake, C.M., and Hutt-Fletcher, L.M. (2004). The Epstein-Barr virus BFRF1 and BFLF2 proteins interact and coexpression alters their cellular localization. *Virology* 320, 99–106.
- Lanman, J., Crum, J., Deerinck, T.J., Gaietta, G.M., Schneemann, A., Sosinsky, G.E., Ellisman, M.H., and Johnson, J.E. (2008). Visualizing flock house virus infection in *Drosophila* cells with correlated fluorescence and electron microscopy. *J Struct Biol* 161, 439–446.
- Leach, N., Bjerke, S.L., Christensen, D.K., Bouchard, J.M., Mou, F., Park, R., Baines, J., Haraguchi, T., and Roller, R.J. (2007). Emerin is hyperphosphorylated and redistributed in herpes simplex virus type 1-infected cells in a manner dependent on both UL34 and US3. *J Virol* 81, 10792–10803.
- Lee, S.K., and Longnecker, R. (1997). The Epstein-Barr virus glycoprotein 110 carboxy-terminal tail domain is essential for lytic virus replication. *J Virol* 71, 4092–4097.
- Leege, T., Granzow, H., Fuchs, W., Klupp, B.G., and Mettenleiter, T.C. (2009). Phenotypic similarities and differences between UL37-deleted pseudorabies virus and herpes simplex virus type 1. *J Gen Virol* 90, 1560–1568.
- Leuzinger, H., Ziegler, U., Schraner, E.M., Fraefel, C., Glauser, D.L., Heid, I., Ackermann, M., Mueller, M., and Wild, P. (2005). Herpes simplex virus 1 envelopment follows two diverse pathways. *J Virol* 79, 13047–13059.
- Liu, F., and Zhou, Z.H. (2006). Comparative virion structures of human herpesviruses. In *Human herpesviruses: biology, therapy, and immunoprophylaxis*, G.C.-F. A. Arvin, P. Moore, et al., ed. (Cambridge, United Kingdom, Cambridge University Press), pp. 27–42.
- Liu, Y., Cui, Z., Zhang, Z., Wei, H., Zhou, Y., Wang, M., and Zhang, X. E. (2009). The tegument protein UL94 of human cytomegalovirus as a binding partner for tegument protein pp28 identified by intracellular imaging. *Virology* 388, 68–77.
- Loomis, J.S., Bowzard, J.B., Courtney, R.J., and Wills, J.W. (2001). Intracellular trafficking of the UL11 tegument protein of herpes simplex virus type 1. *J Virol* 75, 12209–12219.
- Loomis, J.S., Courtney, R.J., and Wills, J.W. (2003). Binding partners for the UL11 tegument protein of herpes simplex virus type 1. *J Virol* 77, 11417–11424.
- Loret, S., Guay, G., and Lippé, R. (2008). Comprehensive characterization of extracellular herpes simplex virus type 1 virions. *J Virol*

- 82, 8605–8618.
- MacLean, C.A., Clark, B., and McGeoch, D.J. (1989). Gene UL11 of herpes simplex virus type 1 encodes a virion protein which is myristylated. *J Gen Virol* 70, 3147–3157.
- MacLean, C.A., Dolan, A., Jamieson, F.E., and McGeoch, D.J. (1992). The myristylated virion proteins of herpes simplex virus type 1: investigation of their role in the virus life cycle. *J Gen Virol* 73, 539–547.
- Martin, B.R., Giepmans, B.N., Adams, S.R., and Tsien, R.Y. (2005). Mammalian cell-based optimization of the biarsenical-binding tetracysteine motif for improved fluorescence and affinity. *Nat Biotechnol* 23, 1308–1314.
- McNab, A.R., Desai, P., Person, S., Roof, L.L., Thomsen, D.R., Newcomb, W.W., Brown, J.C., and Homa, F.L. (1998). The product of the herpes simplex virus type 1 UL25 gene is required for encapsidation but not for cleavage of replicated viral DNA. *J Virol* 72, 1060–1070.
- McNabb, D.S., and Courtney, R.J. (1992). Characterization of the large tegument protein (ICP1/2) of herpes simplex virus type 1. *Virology* 190, 221–232.
- Meckes, D.G. Jr, Marsh, J.A., and Wills, J.W. (2010). Complex mechanisms for the packaging of the UL16 tegument protein into herpes simplex virus. *Virology* 398, 208–213.
- Mettenleiter, T.C. (2002). Herpesvirus assembly and egress. *J Virol* 76, 1537–1547.
- Mettenleiter, T.C. (2004). Budding events in herpesvirus morphogenesis. *Virus Res* 106, 167–180.
- Mettenleiter, T.C. (2006). Intriguing interplay between viral proteins during herpesvirus assembly or: the herpesvirus assembly puzzle. *Vet Microbiol* 113, 163–169.
- Michael, K., Klupp, B.G., Mettenleiter, T.C., and Karger, A. (2006). Composition of pseudorabies virus particles lacking tegument protein US3, UL47, or UL49 or envelope glycoprotein E. *J Virol* 80, 1332–1339.
- Michel, D., Pavić, I., Zimmermann, A., Haupt, E., Wunderlich, K., Heuschmid, M., and Mertens, T. (1996). The UL97 gene product of human cytomegalovirus is an early-late protein with a nuclear localization but is not a nucleoside kinase. *J Virol* 70, 6340–6346.
- Möhl, B.S., Böttcher, S., Granzow, H., Kuhn, J., Klupp, B.G., and Mettenleiter, T.C. (2009). Intracellular localization of the pseudorabies virus large tegument protein pUL36. *J Virol* 83, 9641–9651.
- Morris, J.B., Hofemeister, H., and O'Hare, P. (2007). Herpes simplex virus infection induces phosphorylation and delocalization of emerin, a key inner nuclear membrane protein. *J Virol* 81, 4429–4437.
- Mossman, K.L., Sherburne, R., Lavery, C., Duncan, J., and Smiley, J. R. (2000). Evidence that herpes simplex virus VP16 is required for viral egress downstream of the initial envelopment event. *J Virol* 74, 6287–6299.
- Mou, F., Forest, T., and Baines, J.D. (2007). US3 of herpes simplex virus type 1 encodes a promiscuous protein kinase that phosphorylates and alters localization of lamin A/C in infected cells. *J Virol* 81, 6459–6470.
- Mou, F., Wills, E., and Baines, J.D. (2009). Phosphorylation of the U(L)31 protein of herpes simplex virus 1 by the U(S)3-encoded kinase regulates localization of the nuclear envelopment complex and egress of nucleocapsids. *J Virol* 83, 5181–5191.
- Mou, F., Wills, E.G., Park, R., and Baines, J.D. (2008). Effects of lamin A/C, lamin B1, and viral US3 kinase activity on viral infectivity, virion egress, and the targeting of herpes simplex virus U(L)34-encoded protein to the inner nuclear membrane. *J Virol* 82, 8094–8104.
- Muranyi, W., Haas, J., Wagner, M., Krohne, G., and Koszinowski, U. H. (2002). Cytomegalovirus recruitment of cellular kinases to dissolve the nuclear lamina. *Science* 297, 854–857.
- Murphy, M.A., Bucks, M.A., O'Regan, K.J., and Courtney, R.J. (2008). The HSV-1 tegument protein pUL46 associates with cellular membranes and viral capsids. *Virology* 376, 279–289.
- Naldinho-Souto, R., Browne, H., and Minson, T. (2006). Herpes simplex virus tegument protein VP16 is a component of primary enveloped virions. *J Virol* 80, 2582–2584.
- O'Regan, K.J., Murphy, M.A., Bucks, M.A., Wills, J.W., and Courtney, R.J. (2007). Incorporation of the herpes simplex virus type 1 tegument protein VP22 into the virus particle is independent of interaction with VP16. *Virology* 369, 263–280.
- Overton, H.A., McMillan, D.J., Klavinskis, L.S., Hope, L., Ritchie, A.J., and Wong-kai-in, P. (1992). Herpes simplex virus type 1 gene UL13 encodes a phosphoprotein that is a component of the virion. *Virology* 190, 184–192.
- Padula, M.E., Sydnor, M.L., and Wilson, D.W. (2009). Isolation and preliminary characterization of herpes simplex virus 1 primary enveloped virions from the perinuclear space. *J Virol* 83, 4757–4765.
- Peeters, B., de Wind, N., Hooisma, M., Wagenaar, F., Gielkens, A., and Moormann, R. (1992). Pseudorabies virus envelope glycoproteins gp50 and gII are essential for virus penetration, but only gII is involved in membrane fusion. *J Virol* 66, 894–905.
- Prichard, M.N., Britt, W.J., Daily, S.L., Hartline, C.B., and Kern, E.R. (2005). Human cytomegalovirus UL97 Kinase is required for the normal intranuclear distribution of pp65 and virion morphogenesis. *J Virol* 79, 15494–15502.
- Purves, F.C., Spector, D., and Roizman, B. (1991). The herpes simplex virus 1 protein kinase encoded by the US3 gene mediates posttranslational modification of the phosphoprotein encoded by the UL34 gene. *J Virol* 65, 5757–5764.
- Reynolds, A.E., Liang, L., and Baines, J.D. (2004). Conformational changes in the nuclear lamina induced by herpes simplex virus type 1 require genes U(L)31 and U(L)34. *J Virol* 78, 5564–5575.
- Reynolds, A.E., Wills, E.G., Roller, R.J., Ryckman, B.J., and Baines, J.D. (2002). Ultrastructural localization of the herpes simplex virus type 1 UL31, UL34, and US3 proteins suggests specific roles in primary envelopment and egress of nucleocapsids. *J Virol* 76, 8939–8952.
- Rixon, F.J. (1993). Structure and assembly of herpesviruses. *Semin Virol* 4, 135–144.
- Roberts, A.P., Abaitua, F., O'Hare, P., McNab, D., Rixon, F.J., and Pasdeloup, D. (2009). Differing roles of inner tegument proteins pUL36 and pUL37 during entry of herpes simplex virus type 1. *J Virol* 83, 105–116.
- Roizman, B., Knipe, D.M., and Whitely, R.J. (2007). Herpes Simplex Virus. In *Fields Virology*, D.M. Knipe, Howley, P.M., ed. (Philadelphia, Lippincott Williams&Wilkins), pp. 2501–2601.
- Roizman, B., and Pellett, P.E. (2001). Herpesviridae: a brief introduction, In *Fields virology*, D.M.K.a.P.M. Howley, ed. (Philadelphia, PA, Lippincott-Raven Publishers), pp. 2381–2398.
- Rozen, R., Sathish, N., Li, Y., and Yuan, Y. (2008). Virion-wide protein

- interactions of Kaposi's sarcoma-associated herpesvirus. *J Virol* 82, 4742–4750.
- Ryckman, B.J., and Roller, R.J. (2004). Herpes simplex virus type 1 primary envelopment: UL34 protein modification and the US3-UL34 catalytic relationship. *J Virol* 78, 399–412.
- Sathish, N., Zhu, F.X., Yuan, Y., and Farzan, M. (2009). Kaposi's sarcoma-associated herpesvirus ORF45 interacts with kinesin-2 transporting viral capsid-tegument complexes along microtubules. *PLoS Pathog* 5, e1000332.
- Schek, N., and Bachenheimer, S.L. (1985). Degradation of cellular mRNAs induced by a virion-associated factor during herpes simplex virus infection of Vero cells. *J Virol* 55, 601–610.
- Schimmer, C., and Neubauer, A. (2003). The equine herpesvirus 1 UL11 gene product localizes to the trans-golgi network and is involved in cell-to-cell spread. *Virology* 308, 23–36.
- Scott, E.S., and O'Hare, P. (2001). Fate of the inner nuclear membrane protein lamin B receptor and nuclear lamins in herpes simplex virus type 1 infection. *J Virol* 75, 8818–8830.
- Seo, J.Y., and Britt, W.J. (2006). Sequence requirements for localization of human cytomegalovirus tegument protein pp28 to the virus assembly compartment and for assembly of infectious virus. *J Virol* 80, 5611–5626.
- Shanda, S.K., and Wilson, D.W. (2008). UL36p is required for efficient transport of membrane-associated herpes simplex virus type 1 along microtubules. *J Virol* 82, 7388–7394.
- Silva, M.C., Schröder, J., and Shenk, T. (2005). Human cytomegalovirus cell-to-cell spread in the absence of an essential assembly protein. *Proc Natl Acad Sci U S A* 102, 2081–2086.
- Silva, M.C., Yu, Q.C., Enquist, L., and Shenk, T. (2003). Human cytomegalovirus UL99-encoded pp28 is required for the cytoplasmic envelopment of tegument-associated capsids. *J Virol* 77, 10594–10605.
- Simpson-Holley, M., Baines, J., Roller, R., and Knipe, D.M. (2004). Herpes simplex virus 1 U(L)31 and U(L)34 gene products promote the late maturation of viral replication compartments to the nuclear periphery. *J Virol* 78, 5591–5600.
- Simpson-Holley, M., Colgrove, R.C., Nalepa, G., Harper, J.W., and Knipe, D.M. (2005). Identification and functional evaluation of cellular and viral factors involved in the alteration of nuclear architecture during herpes simplex virus 1 infection. *J Virol* 79, 12840–12851.
- Smibert, C.A., Popova, B., Xiao, P., Capone, J.P., and Smiley, J.R. (1994). Herpes simplex virus VP16 forms a complex with the virion host shutoff protein vhs. *J Virol* 68, 2339–2346.
- Söderberg-Nauclér, C. (2006). Does cytomegalovirus play a causative role in the development of various inflammatory diseases and cancer? *J Intern Med* 259, 219–246.
- Steininger, C. (2007). Clinical relevance of cytomegalovirus infection in patients with disorders of the immune system. *Clin Microbiol Infect* 13, 953–963.
- Stylianou, J., Maringer, K., Cook, R., Bernard, E., and Elliott, G. (2009). Virion incorporation of the herpes simplex virus type 1 tegument protein VP22 occurs via glycoprotein E-specific recruitment to the late secretory pathway. *J Virol* 83, 5204–5218.
- Subak-Sharpe, J.H., and Dargan, D.J. (1998). HSV molecular biology: general aspects of herpes simplex virus molecular biology. *Virus Genes* 16, 239–251.
- Sunil-Chandra, N.P., Efsthathiou, S., and Nash, A.A. (1992). Murine gammaherpesvirus 68 establishes a latent infection in mouse B lymphocytes in vivo. *J Gen Virol* 73, 3275–3279.
- Takakuwa, H., Goshima, F., Koshizuka, T., Murata, T., Daikoku, T., and Nishiyama, Y. (2001). Herpes simplex virus encodes a virion-associated protein which promotes long cellular processes in over-expressing cells. *Genes Cells* 6, 955–966.
- Tandon, R., and Mocarski, E.S. (2008). Control of cytoplasmic maturation events by cytomegalovirus tegument protein pp150. *J Virol* 82, 9433–9444.
- Uetz, P., Dong, Y.A., Zeretzke, C., Atzler, C., Baiker, A., Berger, B., Rajagopala, S.V., Roupelieva, M., Rose, D., Fossum, E., *et al.* (2006). Herpesviral protein networks and their interaction with the human proteome. *Science* 311, 239–242.
- Varnum, S.M., Strelow, D.N., Monroe, M.E., Smith, P., Auberry, K.J., Pasa-Tolic, L., Wang, D., Camp, D.G. 2nd, Rodland, K., Wiley, S., *et al.* (2004). Identification of proteins in human cytomegalovirus (HCMV) particles: the HCMV proteome. *J Virol* 78, 10960–10966.
- Vittone, V., Diefenbach, E., Triffett, D., Douglas, M.W., Cunningham, A.L., and Diefenbach, R.J. (2005). Determination of interactions between tegument proteins of herpes simplex virus type 1. *J Virol* 79, 9566–9571.
- Wagenaar, F., Pol, J.M., Peeters, B., Gielkens, A.L., de Wind, N., and Kimman, T.G. (1995). The US3-encoded protein kinase from pseudorabies virus affects egress of virions from the nucleus. *J Gen Virol* 76, 1851–1859.
- Wild, P., Engels, M., Senn, C., Tobler, K., Ziegler, U., Schraner, E.M., Loepfe, E., Ackermann, M., Mueller, M., and Walther, P. (2005). Impairment of nuclear pores in bovine herpesvirus 1-infected MDBK cells. *J Virol* 79, 1071–1083.
- Wild, P., Schraner, E.M., Cantieni, D., Loepfe, E., Walther, P., Müller, M., and Engels, M. (2002). The significance of the Golgi complex in envelopment of bovine herpesvirus 1 (BHV-1) as revealed by cryobased electron microscopy. *Micron* 33, 327–337.
- Wild, P., Senn, C., Manera, C.L., Sutter, E., Schraner, E.M., Tobler, K., Ackermann, M., Ziegler, U., Lucas, M.S., and Kaech, A. (2009). Exploring the nuclear envelope of herpes simplex virus 1-infected cells by high-resolution microscopy. *J Virol* 83, 408–419.
- Wisner, T.W., Wright, C.C., Kato, A., Kawaguchi, Y., Mou, F., Baines, J.D., Roller, R.J., and Johnson, D.C. (2009). Herpesvirus gB-induced fusion between the virion envelope and outer nuclear membrane during virus egress is regulated by the viral US3 kinase. *J Virol* 83, 3115–3126.
- Wolf, D.G., Courcelle, C.T., Prichard, M.N., and Mocarski, E.S. (2001). Distinct and separate roles for herpesvirus-conserved UL97 kinase in cytomegalovirus DNA synthesis and encapsidation. *Proc Natl Acad Sci U S A* 98, 1895–1900.
- Wolfstein, A., Nagel, C.H., Radtke, K., Döhner, K., Allan, V.J., and Sodeik, B. (2006). The inner tegument promotes herpes simplex virus capsid motility along microtubules in vitro. *Traffic* 7, 227–237.
- Yeh, P.C., Meckes, D.G. Jr, and Wills, J.W. (2008). Analysis of the interaction between the UL11 and UL16 tegument proteins of herpes simplex virus. *J Virol* 82, 10693–10700.
- Zhang, Y., Sirko, D.A., and McKnight, J.L. (1991). Role of herpes simplex virus type 1 UL46 and UL47 in alpha TIF-mediated transcriptional induction: characterization of three viral deletion mutants. *J Virol* 65, 829–841.
- Zhou, Z.H., Chen, D.H., Jakana, J., Rixon, F.J., and Chiu, W. (1999).

- Visualization of tegument-capsid interactions and DNA in intact herpes simplex virus type 1 virions. *J Virol* 73, 3210–3218.
- Zhu, F.X., Chong, J.M., Wu, L., and Yuan, Y. (2005). Virion proteins of Kaposi's sarcoma-associated herpesvirus. *J Virol* 79, 800–811.
- Zhu, F.X., King, S.M., Smith, E.J., Levy, D.E., and Yuan, Y. (2002). A Kaposi's sarcoma-associated herpesviral protein inhibits virus-mediated induction of type I interferon by blocking IRF-7 phosphorylation and nuclear accumulation. *Proc Natl Acad Sci U S A* 99, 5573–5578.
- Zhu, Q., and Courtney, R.J. (1994). Chemical cross-linking of virion envelope and tegument proteins of herpes simplex virus type 1. *Virology* 204, 590–599.