

## ERRATUM

# Small GTPases and cilia

Yujie Li<sup>1</sup>, Jinghua Hu<sup>1,2,3</sup>✉

<sup>1</sup> Department of Nephrology and Hypertension, Mayo Clinic, Rochester, Minnesota, USA

<sup>2</sup> Translational PKD Center, Mayo Clinic, Rochester, Minnesota, USA

<sup>3</sup> Department of Biochemistry and Molecular Biology, Mayo Clinic, Rochester, Minnesota, USA

✉ Correspondence: Hu.jinghua@mayo.edu

Received January 3, 2011 Accepted January 6, 2011

---

Erratum to: Protein Cell 2011, 1(1): 13–25

DOI 10.1007/s13238-011-1004-7

Two references were missed on left column of page 16.

The original sentence is:

Arf3 has been shown to interact with the X-linked retinitis pigmentosa protein (RP2), which is demonstrated to act as the GAP for Arf3 (Grayson et al., 2002).

The corrected sentence is:

Arf3 has been shown to interact with the X-linked retinitis pigmentosa protein (RP2), which is demonstrated to act as the GAP for Arf3 (Bartolini et al., 2002; Grayson et al., 2002; Veltel et al., 2008).

### REFERENCES

Bartolini, F., Bhamidipati, A., Thomas, S., Schwahn, U., Lewis, S.A., and Cowan, N.J. (2002). Functional overlap between retinitis pigmentosa 2 protein and the tubulin-specific chaperone cofactor C. *J Biol Chem* 277, 14629–14634.

Veltel, S., Gasper, R., Eisenacher, E., and Wittinghofer, A. (2008). The retinitis pigmentosa 2 gene product is a GTPase-activating protein for Arf-like 3. *Nat Struct Mol Biol* 15, 373–380.

The online version of the original article can be found at  
<http://dx.doi.org/10.1007/s13238-011-1004-7>