

## REVIEW

# Signaling control of the constitutive androstane receptor (CAR)

Hui Yang, Hongbing Wang✉

Department of Pharmaceutical Sciences, University of Maryland School of Pharmacy, 20 Penn Street, Baltimore, MD 21201, USA

✉ Correspondence: hwang@rx.umaryland.edu (H. Wang)

Received November 24, 2013 Accepted December 7, 2013

## ABSTRACT

The constitutive androstane receptor (CAR, NR1I3) plays a crucial role in the regulation of drug metabolism, energy homeostasis, and cancer development through modulating the transcription of its numerous target genes. Different from prototypical nuclear receptors, CAR can be activated by either direct ligand binding or ligand-independent (indirect) mechanisms both initiated with nuclear translocation of CAR from the cytoplasm. In comparison to the well-defined ligand-based activation, indirect activation of CAR appears to be exclusively involved in the nuclear translocation through mechanisms yet to be fully understood. Accumulating evidence reveals that without activation, CAR forms a protein complex in the cytoplasm where it can be functionally affected by multiple signaling pathways. In this review, we discuss recent progresses in our understanding of the signaling regulation of CAR nuclear accumulation and activation. We expect that this review will also provide greater insight into the similarity and difference between the mechanisms of direct vs. indirect human CAR activation.

**KEYWORDS** constitutive androstane receptor, nuclear translocation, phosphorylation, signaling regulation

## INTRODUCTION

The constitutive androstane receptor (CAR), a member of the nuclear receptor superfamily (subfamily 1, group I, member 3 [NR1I3]), plays an important role in coordinating cellular responses to the stimulation of both exogenous and endogenous chemicals by regulating the expression of its target genes (Qatanani and Moore, 2005; Stanley et al.,

2006; Plant, 2007). Originally cloned as a constitutively activated receptor without a clearly defined biological function, the importance of CAR in xenobiotic metabolism was first appreciated when CAR was functionally linked to the long-known phenobarbital-mediated induction of hepatic cytochrome P450 (CYP) 2B gene family (Honkakoski et al., 1998; Kawamoto et al., 1999). Encouraged by these findings, numerous investigations have been carried out to explore the role of CAR in xenobiotic metabolism, detoxification, and clearance (Maglich et al., 2002; Tolson and Wang, 2010). In humans, two functional enhancer modules, namely the phenobarbital-responsive enhancer module (PBREM) and the xenobiotic-responsive enhancer module (XREM), have been identified upstream of the *CYP2B6* gene and functionally characterized as the CAR binding sites in response to chemical stimuli (Honkakoski et al., 1998; Wang et al., 2003). CAR is also known to control the inductive expression of other CYP enzymes such as CYP3A4 (Goodwin et al., 2002), CYP2Cs (Ferguson et al., 2002; Gerbal-Chaloin et al., 2002), CYP2A6 (Wortham et al., 2007), and to a lesser extent CYP1A1 and CYP1A2 (Yoshinari et al., 2010), which contribute to the metabolism of approximately 75% of clinically used drugs and the detoxification of numerous environmental chemicals (Johansson and Ingelman-Sundberg, 2010). Further studies have extended CAR target genes including those encoding phase II enzymes such as the uridine diphosphate glucuronosyltransferase (UGT) isoforms (i.e., UGT1A1, UGT1A6, and UGT1A9) (Sugatani et al., 2005; Osabe et al., 2008; Buckley and Klaassen, 2009), glutathione S-transferases and sulfotransferases (Maglich et al., 2002; Yanagiba et al., 2009), as well as efflux and uptake drug transporters such as multidrug resistance-associated proteins (MRPs) (Cherrington et al., 2002, 2003), multidrug resistance protein 1 (MDR1) (Burk et al., 2005a, 2005b; Cervený et al., 2007),

and organic anion-transporting polypeptide 1 (OATP1) (Ding et al., 2006; Osabe et al., 2008). In addition to its broad spectrum of target genes, CAR also senses numerous xenobiotics and endobiotics as activators or deactivators and translates chemical stimulation into coordinated metabolism, detoxification, and clearance in the liver. Up-regulation of these drug-metabolizing enzymes or drug transporters by CAR activators may accelerate the biotransformation of co-administered drugs, usually leading to decreased therapeutic efficacy, enhanced toxicity, or increased bioactivation of prodrugs. For instance, recent studies in our lab have demonstrated that activation of CAR can enhance the bioactivation of cyclophosphamide (CPA) and facilitate CPA-based chemotherapeutic activity in leukemia cells (Wang et al., 2013). Understanding the role of CAR in mediating variable drug responsiveness and drug-drug interactions has become an intense focus of both academic and industrial research efforts and may lead to enhanced prediction of drug-drug interactions and xenobiotic-induced cytotoxicity.


Other than the well-established roles of CAR in the regulation of drug metabolism and transport, where it functions as a xenobiotic sensor, emerging evidence strongly suggests that CAR also modulates various hepatic functions that control diverse physiological and pathophysiological conditions, including energy metabolism, insulin signaling, cell proliferation, and tumor development (Fig. 1). In mice, selective activation of CAR significantly alleviated high fat diet-induced obesity and type 2 diabetes via a combined inhibition of lipogenesis, fatty acid synthesis, and gluconeogenesis, as well as the increase of energy expenditure in brown adipose tissues (Dong et al., 2009; Gao et al., 2009; Masuyama and Hiramatsu, 2012). Particularly, CAR influences energy homeostasis by suppressing the expression of phosphoenolpyruvate carboxykinase (PEPCK), glucose-6-phosphatase (G6Pase) (Kachaylo et al., 2012), sterol regulatory element-binding protein 1c (Roth et al., 2008), acetyl-CoA carboxylase 1, fatty acid synthase (FAS), and stearoyl-CoA desaturase-1 (SCD-1) (Du et al., 2008). The essential role of CAR in phenobarbital- and 1,4-bis[2-(3,5-dichloropyridyloxy)] benzene (TCPOBOP)-induced tumor promotion was initially established by using CAR knockout and wild-type mice (Yamamoto et al., 2004; Huang et al., 2005). In this regard, the known tumor promoters stimulated cancer progression by a CAR-dependent perturbation of the expression of the growth arrest and DNA damage-inducible 45 beta (GADD45B) (Columbano et al., 2005), the murine double minute 2 (mdm2) (Huang et al., 2005), as well as the newly identified tubulin alpha 8 (TUBA8) (Kamino et al., 2011a). In contrast to these observed roles of murine CAR in tumor development, activation of human (h) CAR by the selective activator, 6-(4-chlorophenyl)imidazo[2,1-b][1,3]thiazole-5-carbaldehydeO-(3,4-dichlorobenzyl)oxime (CITCO), appears to be associated with cell cycle arrest and enhanced apoptosis in human brain tumor stem cells,

illustrating an anti-cancer potential (Chakraborty et al., 2011). Moreover, the enhanced cell proliferation by phenobarbital in the liver of wild-type mice was completely abrogated in the double-humanized CAR and pregnane X receptor (PXR) mouse model (Ross et al., 2010). Although the underlying mechanisms of the significant species differences of CAR in tumor development are largely unknown, such variances might be attributed to the divergent regulation of differential genes governing DNA synthesis, cell proliferation, apoptosis, and migration by hCAR vs. its rodent counterparts (Ross et al., 2010; Kamino et al., 2011b; Takizawa et al., 2011). Collectively, findings from these initial basic investigations hold the potential to advance CAR from a well-known xenobiotic sensor to an endobiotic modulator that may eventually become a promising drug target for metabolic disorders as well as cancer therapy.

Unlike PXR, the closest relative of CAR in the nuclear receptor superfamily tree, CAR is constitutively activated in nearly all immortalized cells and spontaneously accumulated in nuclei of these cells prior to chemical stimulated activation (Kawamoto et al., 1999). Moreover, CAR is featured as a nuclear receptor that could be transactivated through either the classical direct ligand binding or a mutedly defined ligand-independent indirect mechanism (Kawamoto et al., 1999; Maglich et al., 2003). These characteristics make the studies of CAR activation extremely challenging and pose major difficulties for evaluating drug-mediated CAR activation *in vitro*. This review is aimed to highlight the recent advances in our understanding of the molecular mechanisms behind drug-mediated nuclear translocation and activation of CAR, with a particular focus centered on signaling pathways that contribute to indirect activation of CAR.

## ACTIVATION OF CAR

As a so-called orphan receptor, CAR can be activated by a broad array of xenobiotic chemicals, often at micromolar concentrations, which differs from the classical steroid-hormone receptors which respond to endogenous ligands at nanomolar concentrations (Giguere, 1999; Tzameli and Moore, 2001). Structurally, however, CAR shares common functional features with other typical nuclear receptors, including a highly variable N-terminal AF1 domain, a DNA binding domain (DBD), a ligand-binding domain (LBD), and a C-terminal AF2 domain. The highly conserved DBD contains unique structures that can recognize and bind to specific promoter regions in target genes, namely xenobiotic response elements such as the aforementioned PBREM and XREM in the CYP2B6 promoter. Response elements binding to CAR are usually composed of two direct repeats of the consensus hexameric sequence of AG(G/T)TCA spaced by three to four nucleotides (i.e., DR3 or DR4) (Makinen et al., 2002). In the nucleus, CAR only binds to its response elements after forming heterodimers with the retinoid X receptor (RXR). The X-ray crystal structure of the hCAR/RXR LBDs



**Figure 1. Schematic illustration of biological functions of CAR.** The size of hollow arrows indicates the abundance of available evidence for each function of CAR. Up and down black arrows symbolize increased and decreased gene expression, respectively.

reveals that CAR contains a single-turn Helix X that restricts the conformational freedom of the C-terminal AF2, and a relatively small ligand binding pocket (Xu et al., 2004). Such features permit CAR to interact with co-activator proteins and maintain a constitutively activated status once translocated into the nucleus. Importantly, although CAR shares several common characteristics with classical nuclear receptors, increasing evidence suggests that CAR can be activated by both direct ligand-binding and ligand-independent mechanisms (Kawamoto et al., 1999; Maglich et al., 2003). To this end, it appears that CAR activation is a multi-step process and most identified CAR activators may not directly bind to the receptor.

**Direct activation**

Owing to the constitutive activation of CAR, the initial search for CAR ligands has resulted with the identification of androstenol and androstanol as inverse-agonists of CAR (Forman et al., 1998). Mechanistically, these androstanes convert CAR from constitutive to basal activity by disrupting the salt bridge that locks the H12 helix in its active conformation, promoting co-activator release from the LBD without interfering CAR/RXR dimerization or DNA binding (Shan et al., 2004). Subsequent studies uncovered TCPOBOP, the most potent known member of the phenobarbital-like class of CYP2B inducers, as the first agonist of mouse (m) CAR, in

that TCPOBOP dose-dependently restores mCAR activity following inhibition by the inverse agonists (Tzamelis et al., 2000). Notably, mutation of key residues inside the mCAR ligand-binding pocket entirely eliminated the stimulatory effect of TCPOBOP, as well as the inhibitory effect of androstanes, without affecting the constitutive activity of CAR (Tzamelis et al., 2000). These results clearly establish CAR as a xenobiotic responsive modular protein that can be activated/deactivated by binding with agonistic and antagonistic ligands, respectively.

Importantly, CAR exhibits remarkable species selectivity in its ligand binding and activation profiles, which makes direct extrapolation of findings from mouse to human extremely risky. For example, TCPOBOP and estradiol activate mouse but not human CAR, and pharmacological concentrations of androstanol, progesterone, and testosterone repress mouse but not human CAR (Handschin and Meyer, 2003; Maglich et al., 2003). The first selective hCAR agonist, the imidazothiazole derivative-CITCO, came through a combination of *in vitro* and cell-based screening in 2003 (Maglich et al., 2003). CITCO selectively binds to hCAR and activates CAR target genes in human primary hepatocytes (Maglich et al., 2003; Ferguson et al., 2005; Faucette et al., 2006). Recent evidence also reveals that CITCO can efficiently enhance recruitment of co-activators to the LBD of hCAR by competing with antagonists such as PK11195 (Li et al., 2008) and metformin (Yang et al., 2013). However,

CITCO only moderately enhances the constitutively activated hCAR (less than 2-fold), compared with that of TCPOBOP for mCAR (5- to 10-fold) in cell-based luciferase reporter assays (Tzamelis et al., 2000; Maglich et al., 2003). Moreover, CITCO also activates human PXR and induces PXR target genes at higher concentrations, leaving direct comparison of human CAR and PXR target genes yet challenging (Maglich et al., 2003). To date, there is no pure hCAR agonist reported. Other chemicals exhibiting agonistic effects on hCAR such as the antimalarial artemisinin, the psychoactive diazepam and the anti-fungal myclobutani, are also associated with potent activation of hPXR (Burk et al., 2005b; Li et al., 2009). Overall, ligand-dependent direct activation of CAR may still rely on its initial step of translocating CAR into the nucleus. Once inside the nucleus, CITCO bound hCAR adopts a conformation similar to the constitutively active apo-CAR and maintains the intrinsically high constitutive activity.

#### Indirect activation

The hallmark feature that differentiates CAR from classical nuclear receptors lies in its ligand-independent nuclear accumulation and constitutive activation once expressed inside the nucleus of cells. To date, numerous CAR activators have been identified, including clinically used drugs, environmental chemicals, and endogenous steroid metabolites (Qatanani and Moore, 2005; Li and Wang, 2010; Molnar et al., 2013). Most of these activators however do not bind directly to CAR; instead activating CAR by stimulating its nuclear translocation in a ligand-independent manner (Li et al., 2009). For example, the typical CYP2B inducer and CAR activator phenobarbital does not bind directly to CAR but induces CAR transcriptional activation exclusively via nuclear translocation (Kawamoto et al., 1999; Moore et al., 2000). Notably, constitutive activation of CAR is not always a beneficial feature. In this regard, CAR activation can enhance the metabolism and toxicity of some drugs, such as acetaminophen (Zhang et al., 2002), and potentially increase tumor propensity by stimulating cell proliferation (Takizawa et al., 2011). To accommodate such potential adversity, CAR is primarily located in the cytoplasm prior to activation in primary hepatocytes and intact liver *in vivo* (Kawamoto et al., 1999; Li et al., 2009). In this native hepatocyte environment, CAR is spontaneously sequestered in the cytoplasm as a multi-protein complex including the heat shock protein 90 (Hsp90), cytoplasmic CAR retention protein, protein phosphatase 1 regulatory subunit 16A, and potentially other yet unidentified proteins (Kobayashi et al., 2003; Yoshinari et al., 2003; Sueyoshi et al., 2008). Upon the stimulation of phenobarbital-type indirect activators or CITCO/TCPOBOP-like direct ligand-binding, CAR disassociates from the cytoplasmic localized protein complex and moves into the nucleus. It was believed that this process is regulated by protein kinase-mediated phosphorylation/dephosphorylation of CAR. A

major breakthrough came with the identification of the conserved threonine (Thr)-38 of human CAR as the primary residue that governs nuclear translocation and activation of CAR (Mutoh et al., 2009). Dephosphorylation of the Thr-38 appears to be essential for CAR translocation regardless of exposure to direct or indirect activators (Mutoh et al., 2009). The exact molecular mechanisms controlling Thr-38 CAR phosphorylation/dephosphorylation remain to be fully understood. However, several kinase signaling pathways have recently been suggested to be important in the phosphorylation of CAR.

#### Protein phosphatase 2A (PP2A)

The role of protein kinase-based signaling pathways in controlling phenobarbital-mediated induction of CYP450s had been proposed, even before CAR was recognized as the fundamental target of phenobarbital. Early studies showed that both activation of protein kinase A (PKA) by elevated intracellular cyclic adenosine monophosphate (cAMP) and the inhibition of protein phosphatases PP1 and PP2A by okadaic acid (OA) resulted in complete repression of phenobarbital-inducible CYP gene transactivation in primary rat hepatocytes (Sidhu and Omiecinski, 1995, 1997). Although the transcription factor(s) that drive the phenobarbital induction event was/were yet to be determined, these results indicated that both PKA and protein phosphatase pathways exert marked roles in modulating the signaling of phenobarbital-mediated CYP induction. After establishing CAR as the critical DNA-binding protein required for phenobarbital response, Negishi and colleagues demonstrated that OA pretreatment was sufficient to inhibit phenobarbital-mediated nuclear translocation of CAR and induction of Cyp2b10 in primary mouse hepatocytes, suggesting that CAR nuclear accumulation is most likely regulated by a dephosphorylation-sensitive signaling cascade (Kawamoto et al., 1999). Further studies from the same research group revealed that CAR exists as a complex with Hsp90 in the cytoplasm of non-induced mouse liver hepatocytes. More importantly, phenobarbital treatment recruited PP2A to the protein complex, which led to the dephosphorylation of CAR (Yoshinari et al., 2003).

Realizing the importance of phosphorylation/dephosphorylation in CAR nuclear translocation and activation, the next significant question to be answered was which amino acid residue(s) is/are responsible for such chemical-stimulated signaling. Serial-deletion and site-directed mutagenesis of CAR led to the identification of a leucine-rich motif (LXXLXXL) close to the C-terminal region, namely the xenobiotic response sequence (XRS), as the potential functional unit which dictates the nuclear translocation of CAR in response to various phenobarbital-type inducers (Zelko et al., 2001; Xia and Kemper, 2007). Nevertheless, these residues were not direct targets of either PP2A or PKA. A real breakthrough in this regard came with the

diligent work by Mutoh et al. in 2009, in which the Thr-38 residue of hCAR was established as the primary determinant for chemical-mediated phosphorylation/ dephosphorylation of CAR, while dephosphorylation of Thr-38 is a prerequisite for CAR translocation into the nucleus (Mutoh et al., 2009). Consistent with earlier observations, treatment with OA increased the phosphorylation of CAR at Thr-38 and sequestered CAR in the cytoplasm of mouse primary hepatocytes (Mutoh et al., 2013).

### Extracellular signal-regulated kinase (ERK) and p38 mitogen-activated protein kinase (MAPK)

Accumulating evidence has demonstrated that expression of various CYP enzymes was significantly repressed during liver regeneration, infection or inflammation, suggesting cellular signaling molecules such as growth hormones and cytokines may play a role in the expression of xenobiotic-metabolizing CYPs (Bauer et al., 2004; Koike et al., 2007). In particular, two independent studies provided strong evidence to show that phenobarbital-dependent activation of the rat CYP2B1 promoter was repressed by the presence of epidermal growth factor (EGF) but promoted by U0126, a known inhibitor of the MEK-ERK signaling pathway (Bauer et al., 2004; Joannard et al., 2006). Encouraged by these observations, Negishi and coworkers provided further mechanistic evidence suggesting ERK is an endogenous signal, regulating CAR phosphorylation and nuclear translocation, by which U0126-mediated Cyp2b10 induction via ERK1/2 deactivation was completely abrogated in CAR knockout mice (Koike et al., 2007). Moreover, co-immunoprecipitation experiments revealed that activated ERK1/2 co-precipitated only with the Thr-38 phosphorylated CAR, where the C-terminal located XRS appears to be essential for this interaction (Osabe and Negishi, 2011). This interaction was significantly increased after EGF exposure while treatment with U0126 decreased the level of CAR phosphorylation at Thr-38 and eventually released CAR into the nucleus (Osabe and Negishi, 2011).

An outstanding phenomenon observed was that ectopic expression of hCAR in HepG2 cells does not convey optimal induction of CYP2B6 compared to what was observed in human primary hepatocytes; many other cellular signals have been shown to regulate the activation of CAR. Recently, the p38 MAPK was identified as a required factor optimizing CAR activation and CYP2B6 induction in liver cells (Saito et al., 2013). In human primary hepatocytes, p38 MAPK is highly activated, which significantly differs from that in human hepatoma cell lines, including HepG2 cells. Activation of p38 MAPK by anisomycin robustly potentiated induction of CYP2B6 mRNA by CAR activators in HepG2 cells to levels that were comparable to what was observed in ligand-treated human primary hepatocytes. The potential significance of p38 MAPK in chemical-elicited CAR activation was also indirectly supported by the facts that


phenobarbital-mediated induction of CYP2B1 was stimulated in the liver of diabetic rats where p38 MAPK was activated by the disease itself (Yoshida et al., 1996); while CYP induction by phenobarbital was attenuated in tumor-bearing rats where p38 MAPK was down-regulated (Numazawa et al., 2005). Nonetheless, a definite role of p38 MAPK in CAR activation has yet to be established, given that activation of p38 MAPK appears to enhance some but not all target genes of hCAR.

### Epidermal growth factor receptor (EGFR)

Previous studies have shown that phenobarbital-induced CYP2B gene transactivation could be effectively repressed by growth factors, such as EGF and insulin-like growth factor (IGF) (Bauer et al., 2004; Kietzmann et al., 1999; Thasler et al., 2006). EGFR is a member of the ErbB family of receptors that coordinates extracellular signals, such as EGF, to cellular signaling cascades and eventually promotes cell proliferation (Di Fiore et al., 1990). Recent studies by Mutoh et al., identified EGFR as a phenobarbital-responsive receptor that mediates CAR dephosphorylation and activation in mouse primary hepatocytes (Mutoh et al., 2013). As shown in this study, phenobarbital antagonizes EGF-stimulated EGFR phosphorylation and activation; abrogation of EGFR signaling further induces the dephosphorylation of the downstream receptor for activated C kinase 1 (RACK1) at the residue of Tyr-52. The dephosphorylated RACK1 then directly recruits PP2A to the cytosol localized CAR protein complex, where it dephosphorylates and releases CAR into the nucleus (Fig. 2). More importantly, this study provides the first evidence that phenobarbital can directly bind to EGFR at pharmacologically relevant concentrations. Given that phenobarbital is often referred to as an "orphan compound" without a known direct target, EGFR may represent one of the molecular targets that initiates phenobarbital-mediated cellular responses, including CAR activation. On the other hand, phenobarbital may not function as a prototypical EGFR inhibitor, such as gefitinib and erlotinib, which can antagonize EGFR-mediated cell proliferation and tumor development (Nakajima et al., 2012; Shin et al., 2013). In fact, phenobarbital itself is a potent tumor promoter in rodent animals via a CAR-dependent mechanism (Huang et al., 2005; Yamamoto et al., 2004). Therefore, it is reasonable to speculate that phenobarbital might be an atypical antagonist of EGFR, which only selectively inhibits certain downstream events of EGFR signaling.

### AMP activated protein kinase (AMPK)

AMPK is an enzyme that functions as an energy sensor by regulating cellular energy metabolism and homeostasis. AMPK plays an important role in fatty acid oxidation, glucose uptake, and hepatic lipogenesis by reacting to the fluctuation of the cellular AMP:ATP ratio (Hardie et al., 2012; Inoki et al., 2012). Recent studies suggested that AMPK is involved in



**Figure 2. Antagonistic effect of phenobarbital on EGFR signaling and CAR activation.** Arrows indicate activation and the blunt arrow represents deactivation. (This figure was adopted from Mutoh et al., 2013, Science Signaling).

CAR-regulated *CYP2B* gene induction by phenobarbital-type inducers, but the precise role of AMPK in the activation of CAR remains controversial. Studies from Meyer and colleagues showed that AMPK activator 5-AMINO-1- $\beta$ -D-ribofuranosyl-imidazole-4-carboxamide (AICAR), or expression of a constitutively active form of AMPK, mimicked phenobarbital-mediated induction of *CYP2B6* in hepatoma cell lines (Rencurel et al., 2005). On the other hand, liver-specific deletion of AMPK catalytic subunits in mice impaired the inductive expression of *Cyp2b10* and *Cyp3a11*, but did not inhibit the nuclear accumulation of CAR induced by phenobarbital (Rencurel et al., 2006). Therefore, the authors presumed the existence of another control step of CAR signaling independent of translocation. However, an *in vivo* study conversely showed that AICAR and metformin induced CAR nuclear translocation but failed to induce hepatic *CYP2B* genes in mice and rats, suggesting AMPK activation is not sufficient for *CYP2B* induction (Shindo et al., 2007).


In another study, AICAR was shown to prevent nuclear translocation of CAR and repress phenobarbital-induced *CYP2B* expression in rat primary hepatocytes (Kanno et al., 2010). In the same study, metformin and the constitutively active form of AMPK, however, enhanced PBREM-driven transactivation by phenobarbital, suggesting AICAR inhibits CAR translocation in an AMPK-independent manner. Most recently, we have shown that metformin dramatically repressed phenobarbital/CITCO-induced *CYP2B6* expression through inhibiting dephosphorylation and nuclear translocation of CAR (Yang et al., 2013). Consistent with this observation, our data also demonstrated that AICAR mimicked the effect of metformin on *CYP2B6* suppression, and such repression was partially but concentration-dependently restored by co-treatment with compound C (6-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-3-pyridin-4-yl-pyrazolo[1,5-a]pyrimidine, a known inhibitor of AMPK). Although sequence alignments of the conserved Thr-38 region of CAR revealed no consensus AMPK site, signaling molecules downstream of the AMPK pathway such as PKC (He et al., 2009) may function as the switch controlling CAR phosphorylation and its disassociation from the retaining protein complex.

Collectively, these studies implicate rather contradictory outcomes when connecting *CYP2B* transactivation and CAR nuclear translocation to AMPK activation. Some of the disputes however, can be explained at least in part by the diverse physiological properties of different species or cell systems used in these studies, such as immortalized cell lines vs. primary hepatocytes, human cells vs. rodent cells, and *in vivo* vs. *in vitro*. In addition, energy status and nutritional environment of the cells can also influence phenobarbital regulation of the *CYP2B* gene (Yoshida et al., 1996; Rencurel et al., 2006).

### Transcriptional regulation of CAR

Although the biological function of CAR relies predominantly on chemical-mediated activation/deactivation through direct or indirect mechanisms, the expression level of CAR in response to endogenous signals or xenobiotic chemicals may also influence the downstream regulation of its target genes. It is well known that dramatic interindividual differences exist in the expression of hepatic *CYP2B6*, the prototypical target gene of hCAR (Wang and Tompkins, 2008). Nevertheless, the molecular mechanism(s) underlying this large variability remains elusive. In comparison of a panel of 12 individual human liver samples, Chang et al. revealed that substantial interindividual differences of hCAR expression in these samples were significantly and positively correlated with that of *CYP2B6*, indicating the abundance of this transcription factor may contribute to the varied expression of the *CYP2B6* gene in human liver (Chang et al., 2003). Other studies highlighted that different from cognate CAR activation, expression of CAR can be induced by a number of xenobiotics including the glucocorticoid receptor agonist (dexamethasone) (Pascucci et al., 2000) and peroxisome proliferator-activated receptor (PPAR)- $\alpha$  agonists (WY14643 and ciprofibrate) (Saito et al., 2010). *In silico* analysis of the human CAR 5'-regulatory region led to the identification of a putative glucocorticoid responsive element located between -4477 and -4410 base pair (Pascucci et al., 2003), a functional PPAR- $\alpha$  responsive element around -4400 base pair, as well as a conserved hepatocyte nuclear factor 4  $\alpha$  (HNF4 $\alpha$ ) binding site from upstream of the transcriptional start site (Ding et al., 2006).

Previous animal studies have demonstrated that fasting and caloric restriction increase the expression and activity of CAR which in turn coordinates an adaptive response by slowing down the energy expenditure. CAR knockout animals were unable to couple the metabolic adjustment and lost more weight (Maglich et al., 2004; Qatanani et al., 2005). Given that fasting typically increases the plasma level of free fatty acids that are natural ligands of PPAR $\alpha$ , and elevated interaction between PGC-1 $\alpha$  and HNF4 $\alpha$  is a hallmark of fasting adaptation, functionally establishing CAR as a target gene of PPAR $\alpha$  and HNF4 $\alpha$  provides a novel mechanistic model for CAR in energy homeostasis (Ding



**Figure 3. Signaling control of CAR activation.** Chemicals illustrating activation or deactivation of a signaling pathway are denoted in blue and red, respectively.

et al., 2006). Additional evidence indicated that stress-activated protein kinase and ERK signaling pathways are also associated with altered expression of CAR under serum-starvation stress (Osabe et al., 2009). Most recently, our own data have unexpectedly revealed that an insulin-like growth factor-1 receptor (IGF-1R) inhibitor (BMS-665351) significantly induced the expression of CYP3A4 in human primary hepatocytes without activation of either CAR or PXR, instead it selectively induced the expression of CAR (Li et al., 2012). Intriguingly, BMS-665351 did not activate either glucocorticoid receptor or PPAR $\alpha$  at concentrations that induced the expression of CYP3A4 and CAR, implying additional, yet unknown mechanisms may be involved in the transcriptional regulation of CAR. Collectively, in comparison with the heightened focus on the activation and deactivation of CAR, much less is known regarding how the expression of CAR itself is controlled under the challenge of both endogenous and xenobiotic chemicals. Clearly, transcriptional regulation of the regulator would represent another layer of CAR biology.

**CONCLUDING REMARKS**

It is evident now that CAR has evolved into a sensor of both xenobiotic and endobiotic chemicals by governing the

transcription of genes associated with drug metabolism and transport, energy homeostasis, and cell proliferation. Our understanding of the role of CAR in gene regulation as well as the mechanisms of its activation has increased remarkably during the past 15 years. As summarized in this review, an astonishing number of cellular factors and foreign compounds intertwine in the regulation of CAR biological functions. Although CAR shares several common features with its sister receptor PXR, where they overlap in a number of target genes and xenobiotic activators, the mechanisms of CAR activation have been proven to be relatively unique. To date, mounting evidence demonstrates that CAR can be activated through both classical ligand binding and ligand-independent mechanisms, with indirect activation appearing to be predominant. Seminal works by Negishi and colleagues have shown that the phosphorylation status of CAR is pivotal for its cellular localization and activation, which could be influenced by many protein kinase signals (Mutoh et al., 2009). In the meantime, the cellular expression of CAR itself appears to be affected by certain signaling molecules. Together, these data indicate that CAR may represent a cell signaling-regulated nuclear receptor rather than a typical ligand-dependent nuclear receptor (Fig. 3). Given that CAR can be activated both directly and indirectly, it is essential to keep in mind that the ligand binding and kinase signaling may interconnect to

achieve the optimal activation of this receptor. Undoubtedly, better understanding of the signaling control of CAR activation will eventually benefit the prediction of metabolism-based drug exposure as well as the development of CAR modulators as potential drug candidates.

## ACKNOWLEDGEMENTS

The authors are grateful to members of the Wang laboratory for discussions and comments on the paper, in particular, to Caitlin Lynch, Brandy Garzel, and William Hedrich for their critical reading of the manuscript. We apologize to the scientists who made contributions to the field, but have not been cited due to space limitations. This work was supported by NIH grants DK061652 and GM107058.

## ABBREVIATIONS

AICAR, 5-aminoimidazole-4-carboxamide-1- $\beta$ -D-ribofuranoside; AMPK, AMP activated protein kinase; cAMP, cyclic adenosine monophosphate; CAR, constitutive androstane receptor; CITCO, 6-(4-chlorophenyl)imidazo[2,1-b][1,3]thiazole-5-carbaldehydeO-(3,4-dichlorobenzyl)oxime; EGF, epidermal growth factor; EGFR, EGF receptor; EGR1, early growth response 1; ERK, extracellular signal regulated kinase; GRIP1, glucocorticoid receptor-interacting protein 1; HNF4 $\alpha$ , hepatocyte-enriched nuclear receptor 4 alpha; HSP90, heat shock protein 90; MAPK, mitogen-activated protein kinase; OA, okadaic acid; PBREM, phenobarbital-responsive enhancer module; PGC-1 $\alpha$ , peroxisome proliferator-activated receptor gamma coactivator-1 alpha; PKA, protein kinase A; PP2A, protein phosphatases 2A; PXR, pregnane X receptor; RACK1, receptor for activated C kinase 1; RXR, retinoid X receptor; SRC1, steroid receptor co-activator 1; TCPOBOP, 1,4-bis[2-(3,5-dichloropyridyloxy)]benzene; UGT, uridine diphosphate glucuronosyltransferase; XREM, xenobiotic-responsive enhancer module.

## COMPLIANCE WITH ETHICS GUIDELINES

Hui Yang and Hongbing Wang declare that they have no conflict of interest.

## OPEN ACCESS

This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

## REFERENCES

- Bauer D, Wolfram N, Kahl GF, Hirsch-Ernst KI (2004) Transcriptional regulation of CYP2B1 induction in primary rat hepatocyte cultures: repression by epidermal growth factor is mediated via a distal enhancer region. *Mol Pharmacol* 65:172–180
- Buckley DB, Klaassen CD (2009) Induction of mouse UDP-glucuronosyltransferase mRNA expression in liver and intestine by activators of aryl-hydrocarbon receptor, constitutive androstane receptor, pregnane X receptor, peroxisome proliferator-activated receptor alpha, and nuclear factor erythroid 2-related factor 2. *Drug Metab Dispos* 37:847–856
- Burk O, Arnold KA, Geick A, Tegude H, Eichelbaum M (2005a) A role for constitutive androstane receptor in the regulation of human intestinal MDR1 expression. *Biol Chem* 386:503–513
- Burk O, Arnold KA, Nussler AK, Schaeffeler E, Efimova E, Avery BA, Avery MA, Fromm MF, Eichelbaum M (2005b) Antimalarial artemisinin drugs induce cytochrome P450 and MDR1 expression by activation of xenosensors pregnane X receptor and constitutive androstane receptor. *Mol Pharmacol* 67:1954–1965
- Cervený L, Svecová L, Anzenbacherová E, Vrzal R, Staud F, Dvorak Z, Ulrichová J, Anzenbacher P, Pavek P (2007) Valproic acid induces CYP3A4 and MDR1 gene expression by activation of constitutive androstane receptor and pregnane X receptor pathways. *Drug Metab Dispos* 35:1032–1041
- Chakraborty S, Kanakasabai S, Bright JJ (2011) Constitutive androstane receptor agonist CITCO inhibits growth and expansion of brain tumour stem cells. *Br J Cancer* 104:448–459
- Chang TK, Bandiera SM, Chen J (2003) Constitutive androstane receptor and pregnane X receptor gene expression in human liver: interindividual variability and correlation with CYP2B6 mRNA levels. *Drug Metab Dispos* 31:7–10
- Cherrington NJ, Hartley DP, Li N, Johnson DR, Klaassen CD (2002) Organ distribution of multidrug resistance proteins 1, 2, and 3 (Mrp1, 2, and 3) mRNA and hepatic induction of Mrp3 by constitutive androstane receptor activators in rats. *J Pharmacol Exp Ther* 300:97–104
- Cherrington NJ, Slitt AL, Maher JM, Zhang XX, Zhang J, Huang W, Wan YJ, Moore DD, Klaassen CD (2003) Induction of multidrug resistance protein 3 (mrp3) in vivo is independent of constitutive androstane receptor. *Drug Metab Dispos* 31:1315–1319
- Columbano A, Ledda-Columbano GM, Pibiri M, Cossu C, Mengozzi M, Moore DD, Huang W, Tian J, Locker J (2005) Gadd45beta is induced through a CAR-dependent, TNF-independent pathway in murine liver hyperplasia. *Hepatology* 42:1118–1126
- Di Fiore PP, Segatto O, Taylor WG, Aaronson SA, Pierce JH (1990) EGF receptor and erbB-2 tyrosine kinase domains confer cell specificity for mitogenic signaling. *Science* 248:79–83
- Ding X, Lichti K, Kim I, Gonzalez FJ, Staudinger JL (2006) Regulation of constitutive androstane receptor and its target genes by fasting, cAMP, hepatocyte nuclear factor alpha, and the coactivator peroxisome proliferator-activated receptor gamma coactivator-1 alpha. *J Biol Chem* 281:26540–26551
- Dong B, Saha PK, Huang W, Chen W, Abu-Elheiga LA, Wakil SJ, Stevens RD, Ilkayeva O, Newgard CB, Chan L et al (2009) Activation of nuclear receptor CAR ameliorates diabetes and fatty liver disease. *Proc Natl Acad Sci USA* 106:18831–18836
- Du J, Chen Q, Takemori H, Xu H (2008) SIK2 can be activated by deprivation of nutrition and it inhibits expression of lipogenic genes in adipocytes. *Obesity (Silver Spring)* 16:531–538
- Faucette SR, Sueyoshi T, Smith CM, Negishi M, LeCluyse EL, Wang H (2006) Differential regulation of hepatic CYP2B6 and CYP3A4 genes by constitutive androstane receptor but not pregnane X receptor. *J Pharmacol Exp Ther* 317:1200–1209
- Ferguson SS, LeCluyse EL, Negishi M, Goldstein JA (2002) Regulation of human CYP2C9 by the constitutive androstane receptor: discovery of a new distal binding site. *Mol Pharmacol* 62:737–746

- Ferguson SS, Chen Y, LeCluyse EL, Negishi M, Goldstein JA (2005) Human CYP2C8 is transcriptionally regulated by the nuclear receptors constitutive androstane receptor, pregnane X receptor, glucocorticoid receptor, and hepatic nuclear factor 4 alpha. *Mol Pharmacol* 68:747–757
- Forman BM, Tzamelis I, Choi HS, Chen J, Simha D, Seol W, Evans RM, Moore DD (1998) Androstane metabolites bind to and deactivate the nuclear receptor CAR-beta. *Nature* 395:612–615
- Gao J, He J, Zhai Y, Wada T, Xie W (2009) The constitutive androstane receptor is an anti-obesity nuclear receptor that improves insulin sensitivity. *J Biol Chem* 284:25984–25992
- Gerbal-Chaloin S, Daujat M, Pascucci JM, Pichard-Garcia L, Vilarem MJ, Maurel P (2002) Transcriptional regulation of CYP2C9 gene. Role of glucocorticoid receptor and constitutive androstane receptor. *J Biol Chem* 277:209–217
- Giguere V (1999) Orphan nuclear receptors: from gene to function. *Endocr Rev* 20:689–725
- Goodwin B, Hodgson E, D'Costa DJ, Robertson GR, Liddle C (2002) Transcriptional regulation of the human CYP3A4 gene by the constitutive androstane receptor. *Mol Pharmacol* 62:359–365
- Handschin C, Meyer UA (2003) Induction of drug metabolism: the role of nuclear receptors. *Pharmacol Rev* 55:649–673
- Hardie DG, Ross FA, Hawley SA (2012) AMPK: a nutrient and energy sensor that maintains energy homeostasis. *Nat Rev Mol Cell Bio* 13:251–262
- He L, Sabet A, Djedjos S, Miller R, Sun X, Hussain MA, Radovick S, Wondisford FE (2009) Metformin and insulin suppress hepatic gluconeogenesis through phosphorylation of CREB binding protein. *Cell* 137:635–646
- Honkakoski P, Zelko I, Sueyoshi T, Negishi M (1998) The nuclear orphan receptor CAR-retinoid X receptor heterodimer activates the phenobarbital-responsive enhancer module of the CYP2B gene. *Mol Cell Biol* 18:5652–5658
- Huang W, Zhang J, Washington M, Liu J, Parant JM, Lozano G, Moore DD (2005) Xenobiotic stress induces hepatomegaly and liver tumors via the nuclear receptor constitutive androstane receptor. *Mol Endocrinol* 19:1646–1653
- Inoki K, Kim J, Guan KL (2012) AMPK and mTOR in cellular energy homeostasis and drug targets. *Annu Rev Pharmacol* 52:381–400
- Joannard F, Rissel M, Gilot D, Anderson A, Orfila-Lefevre L, Guillouzo A, Atfi A, Lagadic-Gossmann D (2006) Role for mitogen-activated protein kinases in phenobarbital-induced expression of cytochrome P4502B in primary cultures of rat hepatocytes. *Toxicol Lett* 161:61–72
- Johansson I, Ingelman-Sundberg M (2010) Genetic polymorphism and toxicology—with emphasis on cytochrome p450. *Toxicol Sci* 120:1–13
- Kachaylo EM, Yarushkin AA, Pustyl'nyak VO (2012) Constitutive androstane receptor activation by 2,4,6-triphenyldioxane-1,3 suppresses the expression of the gluconeogenic genes. *Eur J Pharmacol* 679:139–143
- Kamino H, Moore R, Negishi M (2011a) Role of a novel CAR-induced gene, TUBA8, in hepatocellular carcinoma cell lines. *Cancer Genet* 204:382–391
- Kamino H, Yamazaki Y, Saito K, Takizawa D, Kakizaki S, Moore R, Negishi M (2011b) Nuclear receptor CAR-regulated expression of the FAM84A gene during the development of mouse liver tumors. *Int J Oncol* 38:1511–1520
- Kanno Y, Inoue Y, Inouye Y (2010) 5-aminoimidazole-4-carboxamide-1-beta-ribofuranoside (AICAR) prevents nuclear translocation of constitutive androstane receptor by AMP-activated protein kinase (AMPK) independent manner. *J Toxicol Sci* 35:571–576
- Kawamoto T, Sueyoshi T, Zelko I, Moore R, Washburn K, Negishi M (1999) Phenobarbital-responsive nuclear translocation of the receptor CAR in induction of the CYP2B gene. *Mol Cell Biol* 19:6318–6322
- Kietzmann T, Hirsch-Ernst KI, Kahl GF, Jungermann K (1999) Mimicry in primary rat hepatocyte cultures of the in vivo perivenous induction by phenobarbital of cytochrome P-450 2B1 mRNA: role of epidermal growth factor and perivenous oxygen tension. *Mol Pharmacol* 56:46–53
- Kobayashi K, Sueyoshi T, Inoue K, Moore R, Negishi M (2003) Cytoplasmic accumulation of the nuclear receptor CAR by a tetratricopeptide repeat protein in HepG2 cells. *Mol Pharmacol* 64:1069–1075
- Koike C, Moore R, Negishi M (2007) Extracellular signal-regulated kinase is an endogenous signal retaining the nuclear constitutive active/androstane receptor (CAR) in the cytoplasm of mouse primary hepatocytes. *Mol Pharmacol* 71:1217–1221
- Li H, Wang H (2010) Activation of xenobiotic receptors: driving into the nucleus. *Expert Opin Drug Metab Toxicol* 6:409–426
- Li L, Chen T, Stanton JD, Sueyoshi T, Negishi M, Wang H (2008) The peripheral benzodiazepine receptor ligand 1-(2-chlorophenyl-methylpropyl)-3-isoquinoline-carboxamide is a novel antagonist of human constitutive androstane receptor. *Mol Pharmacol* 74:443–453
- Li H, Chen T, Cottrell J, Wang H (2009) Nuclear translocation of adenoviral-enhanced yellow fluorescent protein-tagged-human constitutive androstane receptor (hCAR): a novel tool for screening hCAR activators in human primary hepatocytes. *Drug Metab Dispos* 37:1098–1106
- Li L, Sinz MW, Zimmermann K, Wang H (2012) An insulin-like growth factor 1 receptor inhibitor induces CYP3A4 expression through a pregnane X receptor-independent, noncanonical constitutive androstane receptor-related mechanism. *J Pharmacol Exp Ther* 340:688–697
- Maglich JM, Stoltz CM, Goodwin B, Hawkins-Brown D, Moore JT, Kliewer SA (2002) Nuclear pregnane x receptor and constitutive androstane receptor regulate overlapping but distinct sets of genes involved in xenobiotic detoxification. *Mol Pharmacol* 62:638–646
- Maglich JM, Parks DJ, Moore LB, Collins JL, Goodwin B, Billin AN, Stoltz CA, Kliewer SA, Lambert MH, Willson TM et al (2003) Identification of a novel human constitutive androstane receptor (CAR) agonist and its use in the identification of CAR target genes. *J Biol Chem* 278:17277–17283
- Maglich JM, Watson J, McMillen PJ, Goodwin B, Willson TM, Moore JT (2004) The nuclear receptor CAR is a regulator of thyroid hormone metabolism during caloric restriction. *J Biol Chem* 279:19832–19838
- Makinen J, Frank C, Jyrkkarinne J, Gynther J, Carlberg C, Honkakoski P (2002) Modulation of mouse and human phenobarbital-responsive enhancer module by nuclear receptors. *Mol Pharmacol* 62:366–378
- Masuyama H, Hiramatsu Y (2012) Treatment with a constitutive androstane receptor ligand ameliorates the signs of preeclampsia

- in high-fat diet-induced obese pregnant mice. *Mol Cell Endocrinol* 348:120–127
- Molnar F, Kublbeck J, Jyrkkarinne J, Prantner V, Honkakoski P (2013) An update on the constitutive androstane receptor (CAR). *Drug Metabol Drug Interact* 28:79–93
- Moore LB, Parks DJ, Jones SA, Bledsoe RK, Consler TG, Stimmel JB, Goodwin B, Liddle C, Blanchard SG, Willson TM et al (2000) Orphan nuclear receptors constitutive androstane receptor and pregnane X receptor share xenobiotic and steroid ligands. *J Biol Chem* 275:15122–15127
- Mutoh S, Osabe M, Inoue K, Moore R, Pedersen L, Perera L, Reboloso Y, Sueyoshi T, Negishi M (2009) Dephosphorylation of threonine 38 is required for nuclear translocation and activation of human xenobiotic receptor CAR (NR1I3). *J Biol Chem* 284:34785–34792
- Mutoh S, Sobhany M, Moore R, Perera L, Pedersen L, Sueyoshi T, Negishi M (2013) Phenobarbital indirectly activates the constitutive active androstane receptor (CAR) by inhibition of epidermal growth factor receptor signaling. *Sci Signal* 6:ra31
- Nakajima Y, Takagi H, Kakizaki S, Horiguchi N, Sato K, Sunaga N, Mori M (2012) Gefitinib and gemcitabine coordinately inhibited the proliferation of cholangiocarcinoma cells. *Anticancer Res* 32:5251–5262
- Numazawa S, Shindo S, Maruyama K, Chibana F, Kawahara Y, Ashino T, Tanaka S, Yoshida T (2005) Impaired nuclear translocation of CAR in hepatic preneoplastic lesions: association with an attenuated CYP2B induction by phenobarbital. *FEBS Lett* 579:3560–3564
- Osabe M, Negishi M (2011) Active ERK1/2 protein interacts with the phosphorylated nuclear constitutive active/androstane receptor (CAR; NR1I3), repressing dephosphorylation and sequestering CAR in the cytoplasm. *J Biol Chem* 286:35763–35769
- Osabe M, Sugatani J, Fukuyama T, Ikushiro S, Ikari A, Miwa M (2008) Expression of hepatic UDP-glucuronosyltransferase 1A1 and 1A6 correlated with increased expression of the nuclear constitutive androstane receptor and peroxisome proliferator-activated receptor alpha in male rats fed a high-fat and high-sucrose diet. *Drug Metab Dispos* 36:294–302
- Osabe M, Sugatani J, Takemura A, Kurosawa M, Yamazaki Y, Ikari A, Miwa M (2009) Up-regulation of CAR expression through Elk-1 in HepG2 and SW480 cells by serum starvation stress. *FEBS Lett* 583:885–889
- Pascussi JM, Gerbal-Chaloin S, Fabre JM, Maurel P, Vilarem MJ (2000) Dexamethasone enhances constitutive androstane receptor expression in human hepatocytes: consequences on cytochrome P450 gene regulation. *Mol Pharmacol* 58:1441–1450
- Pascussi JM, Busson-Le Coniat M, Maurel P, Vilarem MJ (2003) Transcriptional analysis of the orphan nuclear receptor constitutive androstane receptor (NR1I3) gene promoter: identification of a distal glucocorticoid response element. *Mol Endocrinol* 17:42–55
- Plant N (2007) The human cytochrome P450 sub-family: transcriptional regulation, inter-individual variation and interaction networks. *Biochim Biophys Acta* 1770:478–488
- Qatanani M, Moore DD (2005) CAR, the continuously advancing receptor, in drug metabolism and disease. *Curr Drug Metab* 6:329–339
- Qatanani M, Zhang J, Moore DD (2005) Role of the constitutive androstane receptor in xenobiotic-induced thyroid hormone metabolism. *Endocrinology* 146:995–1002
- Rencurel F, Stenhouse A, Hawley SA, Friedberg T, Hardie DG, Sutherland C, Wolf CR (2005) AMP-activated protein kinase mediates phenobarbital induction of CYP2B gene expression in hepatocytes and a newly derived human hepatoma cell line. *J Biol Chem* 280:4367–4373
- Rencurel F, Foretz M, Kaufmann MR, Stroka D, Looser R, Leclerc I, da Silva Xavier G, Rutter GA, Viollet B, Meyer UA (2006) Stimulation of AMP-activated protein kinase is essential for the induction of drug metabolizing enzymes by phenobarbital in human and mouse liver. *Mol Pharmacol* 70:1925–1934
- Ross J, Plummer SM, Rode A, Scheer N, Bower CC, Vogel O, Henderson CJ, Wolf CR, Elcombe CR (2010) Human constitutive androstane receptor (CAR) and pregnane X receptor (PXR) support the hypertrophic but not the hyperplastic response to the murine nongenotoxic hepatocarcinogens phenobarbital and chlordanes in vivo. *Toxicol Sci* 116:452–466
- Roth A, Looser R, Kaufmann M, Blattler SM, Rencurel F, Huang W, Moore DD, Meyer UA (2008) Regulatory cross-talk between drug metabolism and lipid homeostasis: constitutive androstane receptor and pregnane X receptor increase Insig-1 expression. *Mol Pharmacol* 73:1282–1289
- Saito K, Kobayashi K, Mizuno Y, Fukuchi Y, Furihata T, Chiba K (2010) Peroxisome proliferator-activated receptor alpha (PPAR-alpha) agonists induce constitutive androstane receptor (CAR) and cytochrome P450 2B in rat primary hepatocytes. *Drug Metab Pharmacokinet* 25:108–111
- Saito K, Moore R, Negishi M (2013) p38 mitogen-activated protein kinase regulates nuclear receptor CAR that activates the CYP2B6 gene. *Drug Metab Dispos* 41:1170–1173
- Shan L, Vincent J, Brunzelle JS, Dussault I, Lin M, Ianculescu I, Sherman MA, Forman BM, Fernandez EJ (2004) Structure of the murine constitutive androstane receptor complexed to androstenol: a molecular basis for inverse agonism. *Mol Cell* 16:907–917
- Shin DM, Zhang H, Saba NF, Chen AY, Nannapaneni S, Amin AR, Muller S, Lewis M, Sica G, Kono S et al (2013) Chemoprevention of head and neck cancer by simultaneous blocking of epidermal growth factor receptor and cyclooxygenase-2 signaling pathways: preclinical and clinical studies. *Clin Cancer Res* 19:1244–1256
- Shindo S, Numazawa S, Yoshida T (2007) A physiological role of AMP-activated protein kinase in phenobarbital-mediated constitutive androstane receptor activation and CYP2B induction. *Biochem J* 401:735–741
- Sidhu JS, Omiecinski CJ (1995) cAMP-associated inhibition of phenobarbital-inducible cytochrome P450 gene expression in primary rat hepatocyte cultures. *J Biol Chem* 270:12762–12773
- Sidhu JS, Omiecinski CJ (1997) An okadaic acid-sensitive pathway involved in the phenobarbital-mediated induction of CYP2B gene expression in primary rat hepatocyte cultures. *J Pharmacol Exp Ther* 282:1122–1129
- Stanley LA, Horsburgh BC, Ross J, Scheer N, Wolf CR (2006) PXR and CAR: nuclear receptors which play a pivotal role in drug disposition and chemical toxicity. *Drug Metab Rev* 38:515–597
- Sueyoshi T, Moore R, Sugatani J, Matsumura Y, Negishi M (2008) PPP1R16A, the membrane subunit of protein phosphatase 1beta, signals nuclear translocation of the nuclear receptor constitutive active/androstane receptor. *Mol Pharmacol* 73:1113–1121

- Sugatani J, Nishitani S, Yamakawa K, Yoshinari K, Sueyoshi T, Negishi M, Miwa M (2005) Transcriptional regulation of human UGT1A1 gene expression: activated glucocorticoid receptor enhances constitutive androstane receptor/pregnane X receptor-mediated UDP-glucuronosyltransferase 1A1 regulation with glucocorticoid receptor-interacting protein 1. *Mol Pharmacol* 67:845–855
- Takizawa D, Kakizaki S, Horiguchi N, Yamazaki Y, Tojima H, Mori M (2011) Constitutive active/androstane receptor promotes hepatocarcinogenesis in a mouse model of non-alcoholic steatohepatitis. *Carcinogenesis* 32:576–583
- Thasler WE, Dayoub R, Muhlbauer M, Hellerbrand C, Singer T, Grabe A, Jauch KW, Schlitt HJ, Weiss TS (2006) Repression of cytochrome P450 activity in human hepatocytes in vitro by a novel hepatotrophic factor, augments liver regeneration. *J Pharmacol Exp Ther* 316:822–829
- Tolson AH, Wang H (2010) Regulation of drug-metabolizing enzymes by xenobiotic receptors: PXR and CAR. *Adv Drug Deliv Rev* 62:1238–1249
- Tzamei I, Moore DD (2001) Role reversal: new insights from new ligands for the xenobiotic receptor CAR. *Trends Endocrinol Metab* 12:7–10
- Tzamei I, Pissios P, Schuetz EG, Moore DD (2000) The xenobiotic compound 1,4-bis[2-(3,5-dichloropyridyloxy)]benzene is an agonist ligand for the nuclear receptor CAR. *Mol Cell Biol* 20:2951–2958
- Wang H, Tompkins LM (2008) CYP2B6: new insights into a historically overlooked cytochrome P450 isozyme. *Curr Drug Metab* 9:598–610
- Wang H, Faucette S, Sueyoshi T, Moore R, Ferguson S, Negishi M, LeCluyse EL (2003) A novel distal enhancer module regulated by pregnane X receptor/constitutive androstane receptor is essential for the maximal induction of CYP2B6 gene expression. *J Biol Chem* 278:14146–14152
- Wang D, Li L, Yang H, Ferguson SS, Baer MR, Gartenhaus RB, Wang H (2013) The constitutive androstane receptor is a novel therapeutic target facilitating cyclophosphamide-based treatment of hematopoietic malignancies. *Blood* 121:329–338
- Wortham M, Czerwinski M, He L, Parkinson A, Wan YJ (2007) Expression of constitutive androstane receptor, hepatic nuclear factor 4 alpha, and P450 oxidoreductase genes determines interindividual variability in basal expression and activity of a broad scope of xenobiotic metabolism genes in the human liver. *Drug Metab Dispos* 35:1700–1710
- Xia J, Kemper B (2007) Subcellular trafficking signals of constitutive androstane receptor: evidence for a nuclear export signal in the DNA-binding domain. *Drug Metab Dispos* 35:1489–1494
- Xu RX, Lambert MH, Wisely BB, Warren EN, Weinert EE, Waitt GM, Williams JD, Collins JL, Moore LB, Willson TM et al (2004) A structural basis for constitutive activity in the human CAR/RXRalpha heterodimer. *Mol Cell* 16:919–928
- Yamamoto Y, Moore R, Goldsworthy TL, Negishi M, Maronpot RR (2004) The orphan nuclear receptor constitutive active/androstane receptor is essential for liver tumor promotion by phenobarbital in mice. *Cancer Res* 64:7197–7200
- Yanagiba Y, Ito Y, Kamijima M, Gonzalez FJ, Nakajima T (2009) Octachlorostyrene induces cytochrome P450, UDP-glucuronosyltransferase, and sulfotransferase via the aryl hydrocarbon receptor and constitutive androstane receptor. *Toxicol Sci* 111:19–26
- Yang H, Garzel B, Heyward S, Moeller T, Shapiro P, Wang H (2013) Metformin represses drug-induced expression of CYP2B6 by modulating the constitutive androstane receptor signaling. *Mol Pharmacol* 85(2):249–260
- Yoshida Y, Kimura N, Oda H, Kakinuma A (1996) Insulin suppresses the induction of CYP2B1 and CYP2B2 gene expression by phenobarbital in adult rat cultured hepatocytes. *Biochem Biophys Res Commun* 229:182–188
- Yoshinari K, Kobayashi K, Moore R, Kawamoto T, Negishi M (2003) Identification of the nuclear receptor CAR:HSP90 complex in mouse liver and recruitment of protein phosphatase 2A in response to phenobarbital. *FEBS Lett* 548:17–20
- Yoshinari K, Yoda N, Toriyabe T, Yamazoe Y (2010) Constitutive androstane receptor transcriptionally activates human CYP1A1 and CYP1A2 genes through a common regulatory element in the 5'-flanking region. *Biochem Pharmacol* 79:261–269
- Zelko I, Sueyoshi T, Kawamoto T, Moore R, Negishi M (2001) The peptide near the C terminus regulates receptor CAR nuclear translocation induced by xenochemicals in mouse liver. *Mol Cell Biol* 21:2838–2846
- Zhang J, Huang W, Chua SS, Wei P, Moore DD (2002) Modulation of acetaminophen-induced hepatotoxicity by the xenobiotic receptor CAR. *Science* 298:422–424