

REVIEW

New insights into the regulation of Axin function in canonical Wnt signaling pathway

Xiaomin Song, Sheng Wang, Lin Li✉

State Key Laboratory of Molecular Biology, Institute of Biochemistry and Cell Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, China

✉ Correspondence: lli@sibs.ac.cn (L. Li)

Received November 9, 2013 Accepted December 10, 2013

ABSTRACT

The Wnt signaling pathway plays crucial roles during embryonic development, whose aberration is implicated in a variety of human cancers. Axin, a key component of canonical Wnt pathway, plays dual roles in modulating Wnt signaling: on one hand, Axin scaffolds the “ β -catenin destruction complex” to promote β -catenin degradation and therefore inhibits the Wnt signal transduction; on the other hand, Axin interacts with LRP5/6 and facilitates the recruitment of GSK3 to the plasma membrane to promote LRP5/6 phosphorylation and Wnt signaling. The differential assemblies of Axin with these two distinct complexes have to be tightly controlled for appropriate transduction of the “on” or “off” Wnt signal. So far, there are multiple mechanisms revealed in the regulation of Axin activity, such as post-transcriptional modulation, homo/hetero-polymerization and auto-inhibition. These mechanisms may work cooperatively to modulate the function of Axin, thereby playing an important role in controlling the canonical Wnt signaling. In this review, we will focus on the recent progresses regarding the regulation of Axin function in canonical Wnt signaling.

KEYWORDS Wnt signaling, Axin, post-transcriptional modification, polymerization, auto-inhibition

INTRODUCTION

Wnt signaling is one of the most conserved pathways during evolution, playing essential roles in multiple biological processes like embryonic development and adult tissue homeostasis. Dysregulation of the Wnt pathway underlies the pathogenesis of a variety of human cancers. So far, two types

of Wnt signaling pathways have been characterized: canonical Wnt pathway (also known as Wnt/ β -catenin pathway) and noncanonical Wnt pathway (including planar cell polarity pathway and Wnt/calcium pathway). The Wnt/ β -catenin pathway is distinguished from the others in the way of its dependency on β -catenin, which is nevertheless not required for the two noncanonical Wnt pathways. As a central player in canonical Wnt signaling, β -catenin dictates the signaling intensity through its cellular levels and subcellular localization. In the absence of Wnt signaling, β -catenin is sequestered in the cytoplasm by the “destruction complex” composed of Axin, glycogen synthase kinase 3 (GSK3) and adenomatous polyposis coli (APC), etc. β -catenin is phosphorylated by GSK3 and subsequently subjected to degradation through the ubiquitin-proteasome pathway. Upon Wnt stimulation, low-density lipoprotein receptor-related proteins 5/6 (LRP5/6) is co-clustered with Frizzled (Fz) and Dishevelled (Dvl) and recruits the active Axin and GSK3 complex; these proteins thus form a high-molecular-weight assemblage close to the plasma membrane that is referred to as “LRP signalosome” (Bilic et al., 2007). Following the recruitment of Axin to LRP5/6, the “destruction complex” is disassembled; and β -catenin is released and translocates into the nucleus, where it forms a transcriptional complex with LEF-1/TCF and activates the transcription of Wnt-responsive genes. Recently, Li et al., based on their findings, proposed an alternative model that Wnt stimulation, rather than cause the disassembly of “destruction complex”, suppresses β -catenin degradation and leads to saturation of the complex by the accumulated β -catenin, thus sparing the newly synthesized β -catenin, which accumulates in a free cytosolic form and consequently enters the nucleus to bind LEF-1/TCF (Li et al., 2012).

Vertebrates have two Axin isoforms: Axin1 (also referred to as Axin) is constitutively expressed, while Axin2 (also called Conductin) is subjected to the regulation of Wnt

signaling (Jho et al., 2002; Leung et al., 2002; Yan et al., 2001). Albeit different in their expression patterns, Axin and Axin2 appear to be functionally, at least partly, interchangeable in canonical Wnt pathway (Chia and Costantini, 2005). Considering its essential role in both “destruction complex” and “LRP signalosome”, the activity of Axin has to be tightly controlled. Post-translational modification is the most common mechanism for regulating protein activity. Several ways of modifications, such as phosphorylation, SUMOylation and ubiquitination, have been reported to play a role in regulating Axin stability, sub-cellular localization or association with other proteins (Callow et al., 2011; Fei et al., 2013; Kim et al., 2008; Kim and Jho, 2010; Zhang et al., 2011). Polymerization of Axin also regulates its activity in a way that currently isn't clearly defined. Axin is considered as a protein that is largely unstructured except the N-terminal RGS domain responsible for binding APC and the C-terminal DIX domain for mediating oligomerization. Besides Axin, Dvl and Ccd1 also contain a DIX domain, which is responsible for their homo- and hetero- polymerization (Liu et al., 2011; Schwarz-Romond et al., 2007a; Shiomi et al., 2003). It is reported that DIX-mediated heterotypic interaction between Axin and Dvl may disrupt Axin's association with the “destruction complex” and facilitate its recruitment to the “LRP signalosome” (Mendoza-Topaz et al., 2011; Schwarz-Romond et al., 2007b). Recently, two independent researches carried out by Sung-Eun Kim et al. and our group revealed a new mechanism of “auto-inhibition” behind Axin function, pointing out that conformational changes of Axin may play an important role in regulating its activity (Kim et al., 2013b; Wang et al., 2013). Notably, these mechanisms may be relevant and work cooperatively to fine-tune and regulate the function of Axin.

UBIQUITINATION/SUMOYLATION OF AXIN

Ubiquitination is an important regulatory process for protein, playing diverse roles in modulating protein function. Ubiquitin is covalently attached to the targeted molecules through the sequential activities of three enzymes: ubiquitin-activating enzyme (E1) for ATP-dependent activation of ubiquitin (Ub), ubiquitin-conjugating enzyme (E2) for delivery of Ub, and ubiquitin ligase (E3) for ligation of Ub to the substrates. In this process, the E3 ligases are responsible for substrate selection and the specificity of Ub-linkage. There are seven lysines (K6, K11, K27, K29, K33, K48 and K63) in Ub molecule. Generally, proteins marked by K48-linked poly-Ub chains are destined to proteasome-dependent degradation, while other types of Ub chains could regulate a variety of cellular processes by both proteolytic and non-proteolytic functions (Kulathu and Komander, 2012). Multiple key components in canonical Wnt pathway are regulated by ubiquitination, such as β -catenin, APC and Dvl (Chan et al., 2006; Choi et al., 2004; Ding et al., 2013; Hu et al., 2010; Liu et al.,

1999; Liu et al., 2001; Tran et al., 2013; Tran and Polakis, 2012; Wei et al., 2012; Winston et al., 1999). Despite limited reports so far on ubiquitination of Axin, it's increasingly clear that Axin is also subjected to ubiquitin modification. Previously, Stegmeier lab revealed that Axin stability is regulated by Tankyrase and Tankyrase-mediated poly-ADP-ribose modification (PARsylation) of Axin is linked to Axin polyubiquitylation and subsequent degradation by the proteasome (Huang et al., 2009). Later, RNF146 was uncovered to be the E3 ligase for mediating Tankyrase-dependent degradation of Axin, thus playing a positive role in Wnt signaling (Callow et al., 2011; Zhang et al., 2011). Ubiquitin-specific protease 34 (USP34) is also reported to associate with Axin and control its levels, whereby modulating Wnt signaling (Lui et al., 2011). Smurf2 (Smad ubiquitination regulatory factor 2), a C2-WW-HECT type ubiquitin ligase, is suggested to be another E3 ligase for Axin, which also plays a role in regulating its stability (Kim and Jho, 2010). Interestingly, we recently uncovered that Smurf1, an isoform of Smurf2, also targets Axin, but for a non-proteolytic purpose. Our data suggested that Smurf1 ubiquitinates Axin through Lys (K) 29-linked poly-ubiquitin chains, which disrupts its interaction with LRP5/6 rather than leads to its degradation (Fei et al., 2013). Similar to Ubiquitination, SUMO modification of proteins also plays various roles in regulating protein function. However, researches regarding Axin SUMOylation are, so far, very limited. It was reported earlier that SUMOylation of Axin may regulate its function in noncanonical Wnt pathway (Rui et al., 2002). Recently, Kim et al. suggested that SUMOylation may protect Axin from ubiquitination, thus playing a role in regulating Axin stability in canonical Wnt pathway (Kim et al., 2008).

PHOSPHORYLATION/DEPHOSPHORYLATION OF AXIN

Phosphorylation/Dephosphorylation is the most common post-translational modification for regulating protein function. Multiple components of Wnt pathway, such as LRP5/6, Dvl, β -catenin, TCF, are subject to reversible phosphorylation by kinases and phosphatases networks. Axin, as a crucial component of canonical Wnt pathway, is also under the regulation of phosphorylation/dephosphorylation. Without Wnt stimulation, Axin is phosphorylated, which enhances its binding affinity with β -catenin, leading to stabilization of Axin (Jho et al., 1999; Yamamoto et al., 1999); upon Wnt stimulation, Axin is dephosphorylated, leading to a less effective binding with β -catenin and consequently Axin degradation (Strovel et al., 2000; Willert et al., 1999). GSK3 is most likely the key kinase that phosphorylates Axin and facilitates its function in “destruction complex”. Besides GSK3, CKI was also observed to phosphorylate Axin both *in vitro* and *in vivo* (Gao et al., 2002). On the other hand, it is found that PP2A and PP2C may be the two phosphatases targeting Axin




Figure 1. The C-terminal DIX domain of Axin mediates its homo/hetero- polymerization. Schematic illustration of the domain organization for Axin, Dvl and Ccd1. These three proteins all contain a DIX domain, which mediates their homo- and hetero-interaction. The self-assembly of Axin-DIX or heterotypic interaction with Dvl-DIX may regulate Axin function in canonical Wnt signaling. (B) The three-dimensional structure of Axin DIX domain (PDB code: 1wsp) showed that it forms filaments in the crystal through head-to-tail self-interaction.

(Strovel et al., 2000; Willert et al., 1999). Recently, it was reported that phosphorylation of Axin by CKI may also improve its association with GSK3 and leads to a more active “destruction complex”, while protein phosphatase 1 (PP1) interacts with, and dephosphorylates Axin, reversing the effect conferred by CKI and hence contributing to the activation of Wnt signaling (Luo et al., 2007). Later, Jiang et al. reported that Dab2 could block the interaction between Axin and PP1, thus inhibiting Axin dephosphorylation and ultimately leading to inhibition of the Wnt signaling (Jiang et al., 2009). Previously, phosphorylation of Axin by Cyclin A/CDK2 was reported to increase its association with β -catenin (Kim et al., 2004). Recently, Axin was found to be phosphorylated by CDK5, and this phosphorylation facilitates its interaction with GSK3, which plays an essential role for axon development (Fang et al., 2011).

HOMO- AND HETERO-POLYMERIZATION OF AXIN

It is not unusual that proteins modulate their functions through altering their oligomeric states or forming hetero-oligomer with other proteins. Self- or hetero-assembly of proteins also plays important roles in regulating Wnt signaling. For example, aggregation of LRP5/6 is essential for Wnt signaling activation, which may also require the oligomerization of Dvl (Metcalf et al., 2010). Recently, we identified Caprin-2 as a new component of canonical Wnt signaling through facilitating LRP5/6 phosphorylation (Ding et al., 2008), and we further found that the oligomerization of Caprin-2 is required for its function in Wnt signaling (unpublished data). Axin contains a DIX domain of 80 amino

acids located at its C-terminus for mediating homo- or hetero-interaction, which seems to be essential for its function (Choi et al., 2010; Fiedler et al., 2011; Sakanaka and Williams, 1999; Yokoyama et al., 2012). The three-dimensional structure of Axin DIX domain showed that it forms filaments in the crystal through head-to-tail self-interaction (Schwarz-Romond et al., 2007a; Shibata et al., 2007) (Fig. 1B). Besides Axin, the homologous DIX domain is also present at the N-terminus of Dvl and the C-terminus of Ccd1 (also called DIXdc1) (Fig. 1A). It is argued that Dvl may release Axin from the “destruction complex” and deliver Axin to the “LRP signalosome” through DIX-mediated heterotypic interactions between Dvl and Axin (Schwarz-Romond et al., 2007b). Fiedler et al.’s recent work also indicated that Dvl may behave as a dominant-negative modulator of Axin to regulate its function via heterotypic interactions between their DIX domains (Choi et al., 2010; Fiedler et al., 2011). Ccd1 was initially identified as a positive regulator in Wnt signaling (Shiomi et al., 2003). Ccd1-DIX interacts with Dvl-DIX directly, converting latent polymeric Dvl to a biologically active oligomer(s) (Liu et al., 2011; Shiomi et al., 2003). However, the interaction between Ccd1 and Axin does not seem to be dominated by their DIX domains and other domains may also play an important role for their interaction (Wong et al., 2004).

AUTO-INHIBITION OF AXIN

Conformational change is an important way for proteins to regulate their functions. In many cases, when a protein adopts an “auto-inhibited” conformation, its active site for




Figure 2. The model of Axin autoinhibition. Based on the findings of Kim et al. and us, we proposed a possible model for Axin “autoinhibition”: without Wnt stimulation, the N-terminus of Axin interacts with its C-terminus, forming “type I” closed conformation, which blocks its binding with LRP5/6 but spares the binding site for β -catenin; upon Wnt stimulation, this type I auto-inhibited state of Axin is released for binding to LRP5/6 and the latter was subsequently phosphorylated, which inhibits the activity of GSK3 and thus moves Axin towards to the dephosphorylated state conferred by PP1. Dephosphorylated Axin forms a “type II” closed conformation through an intramolecular interaction between the C-terminus and the β -catenin binding region, making it unable to bind both of LRP5/6 and β -catenin.

substrates, or binding sites for other partners may be blocked and thus this state is also referred to as an “inactive” state; while, in an “active” state, the autoinhibitory conformation is released, making the active or binding sites available for interaction with substrates or regulatory partners. Normally, this reversible process of conformational change is induced and regulated by post-translational modifications or interactions with other proteins. Previous studies have suggested that Axin N-terminus (including the RGS domain and the linker region between the RGS domain and the GSK binding domain) plays an inhibitory role in Axin’s binding with its partners (Chen et al., 2009; Mao et al., 2001). However, the mechanism underlying such inhibition remains elusive. In our recent work, we revealed that the N-terminus of Axin is able to associate with its C-terminus, forming a “closed” conformation, which could be disrupted effectively by a small molecular compound, named as HLY78. HLY78 is an activator of Wnt/ β -catenin signaling pathway identified by our group through high-throughput

screening, which acts in a Wnt ligand-dependent manner (Wang et al., 2013). Our findings suggested that depending on distinct partners and post-translational modifications, Axin may undergo conformational change during its differential assemblies with “destruction complex” or “Wnt-LRP5/6 signalosome”. We further found that HLY78 targets Axin-DIX and triggers the conformational switch of Axin from a “closed” auto-inhibitory state to an “open” active state, leading to an enhanced association of Axin with LRP6 and subsequent activation of LRP6. Meanwhile, an independent work carried out by another group also proposed an “auto-inhibition” model of Axin (Kim et al., 2013b). In their model, phosphorylation by GSK3 makes Axin adopt an “open” conformation for binding to β -catenin or phospho-LRP6, while dephosphorylation by PP1 leads to a “closed” conformation (auto-inhibition) of Axin, making Axin incompetent for interacting with either β -catenin or phospho-LRP6. Notably, according to their findings, the C-terminus of Axin forms a “closed” conformation through an intramolecular interaction

with the middle β -catenin binding region rather than the N-terminal region as we found. In our view, these two models are not in conflict with each other and may function alternatively or collaboratively in different cellular contexts. A possible model that accommodates both sets of findings is proposed in Fig. 2: without Wnt stimulation, the N-terminus of Axin interacts with its C-terminus, forming “type I” closed conformation, which blocks its binding with LRP5/6 but spares the one with β -catenin; upon Wnt stimulation, this type I auto-inhibited state of Axin is released for binding to LRP5/6 and the latter was subsequently phosphorylated, which inhibits the activity of GSK3 and thus moves Axin towards to the dephosphorylated state conferred by PP1. Dephosphorylated Axin forms a “type II” closed conformation through an intramolecular interaction between the C-terminus and the β -catenin binding region, making it unable to bind both of LRP5/6 and β -catenin.

MICRORNAS MEDIATED REGULATION OF AXIN

MicroRNA (miRNA) is a small noncoding RNA molecule normally containing 19–25 nucleotides in length, which regulates gene expression both at transcriptional and post-transcriptional level. So far, thousands of transcripts have been reported to subject to microRNA regulation, playing extensive and important roles in eukaryotic development and physiology. Moreover, increasing evidences have implicated the role of miRNA deregulation in human disease including cancer. Recently, there are several researches addressing potential roles of miRNAs in Axin regulation. In a screen for miRNAs affecting Wnt pathway, Silver et al. found that ectopic miR-315 directly inhibits Axin as well as Notum (Silver et al., 2007), thus acting as an activator of Wnt signaling. Later, another miRNA—hsa-miR-34a—reportedly targets both the 5'-UTR and 3'-UTR of *Axin2*, whereby repressing *Axin2* expression (Lee et al., 2009). Recently, Egea et al. revealed that *lethal-7 (let-7)* also targets and represses the translation of *Axin2*. *lethal-7 (let-7)* is one of the first two known miRNAs (the other one is *lin-4*) (Reinhart et al., 2000), which is highly conserved across species and implicated in the regulation of a variety of important genes, such as *RAS* and *HMGA2* (Johnson et al., 2005; Mayr et al., 2007). They found miRNA let-7f promotes β -catenin activity in hMSCs by repressing the translation of *Axin2* (Egea et al., 2012). One the other hand, in a study lately published, Kim et al. reported that upregulation of miR-205 in KB oral cancer cells could trigger cell apoptosis. Further mechanistic study identified *Axin2* as the direct target of miR-205 (Lee et al., 2013). They showed that over-expressed miR-205 in KB oral cancer cells suppresses *Axin2* expression through an interaction with its own binding site at *Axin2* 3'-UTR (64–92). However, it is unclear how reduced *Axin2* expression conferred by miR-205, rather than facilitates tumorigenesis as it usually does, contributes to the suppression of KB oral cancer cells.

EPIGENETIC MODULATIONS OF AXIN IN CANCER

In addition to the post-transcriptional regulations mentioned above in affecting Axin levels, epigenetic modulations could also affect the expression of Axin and the change thereof may relate to tumor development. It is well accepted that genetic changes in components of Wnt signaling are closely related with human cancer. One leading example is colorectal cancer: alterations in Wnt-related genes, mainly APC, have been observed in 90% of colorectal cancer (Miyaki et al., 1994; Najdi et al., 2011). In addition to genetic mutations, epigenetic modification is another common cause for altered gene expression, among which promoter methylation and histone modification are the most frequently observed. Axin is also frequently found mutated in cancer, contributing to the over-activation of Wnt signaling in these cancers, while studies regarding epigenetic alterations of Axin and its relationship to cancer remain relatively scant. As a negative regulator of Wnt signaling, tumor-specific promoter methylation or histone deacetylation will cause decreased Axin expression, which may prompt over-activation of Wnt signaling in cancer. Previously, epigenetic silencing of *Axin2* was reported in colorectal carcinoma with microsatellite instability (MSI+ CRC), and forced expression of *Axin2* results in rapid cell death in an MSI+ CRC cell line, indicating a role of epigenetic silencing of *Axin2* in carcinogenesis of MSI+ CRC (Koinuma et al., 2006). Lately, CpG islands of *Axin2* were found methylated in human neuroendocrine tumors (NETs) and *Axin2* expression is correspondingly downregulated, which may contributes to the pathogenesis and growth of NETs (Kim et al., 2013a). A recent work carried out by Yang et al., also indicated that hypermethylated *Axin2* gene may significantly correlate with the progression of lung cancer (Yang et al., 2013). On the other hand, Chen G. et al. reported that overexpression of Menin, a tumor suppressor protein mutated in patients with multiple endocrine neoplasia type 1, is associated with increased H3K4 trimethylation of the *Axin2* gene promoter; while, inhibition of Menin expression by siRNA abrogates H3K4 trimethylation and gene expression of *Axin2*, indicating that Menin functions, at least partly, through affecting histone trimethylation of *Axin2* promoter (Chen et al., 2008). Recently, Histone deacetylase (HDAC) 1 and 2 are implicated in the regulation of *Axin2* expression in non-small cell lung cancer (NSCLC) (Han et al., 2012). X-radiation and siRNA could inhibit expression of HDAC1 and HDAC2 and weaken its inhibitory effect on *Axin2*, which upregulates *Axin2* expression and induces apoptosis of lung cancer cells (Han et al., 2012).

CONCLUDING REMARKS AND PERSPECTIVES

Besides those post-transcriptional modulations mentioned above, another modification—methylation—is also observed for Axin (Cha et al., 2011; Huang et al., 2009). A recent work carried out by Cha et al. revealed that PRMT1 directly

interacts with and methylates Axin, thus enhancing Axin stability (Cha et al., 2011; Huang et al., 2009). So far, our knowledge about post-transcriptional modifications of Axin is still very limited. More kinases, phosphatases, E3 ubiquitin ligases and probably other types of post-transcriptional regulations are expected to be identified in the near future to fit into the network that modulates Axin activity in canonical Wnt signaling. In light of current findings, Axin seems to be a flexible and dynamic molecule, which could undergo structural rearrangements for interacting with distinct partners. The three-dimensional structure of the full-length Axin will provide invaluable insights into the details regarding Axin auto-inhibition. On the other hand, increasing evidence indicates a role of Axin in nucleus (Nikuseva Martic et al., 2010; Pecina-Slaus et al., 2011; Schmitz et al., 2013; Willert and Jones, 2006). However, the precise role of nuclear Axin in Wnt signaling remains largely unknown, which warrants further investigation to facilitate our understanding of the function of nuclear Axin as well as its linkage to the cytoplasmic one. As an attractive therapeutic target, much effort has been made to identify chemical compounds modulating Axin activity. So far, most studies focus on small-molecules in improving Axin stability and thus attenuating Wnt signaling, among which inhibitors against Tankyrase are the best explored (Bao et al., 2012; Huang et al., 2009; James et al., 2012; Lu et al., 2009; Shultz et al., 2012). Tankyrase-mediated PARsylation of Axin stimulates Axin degradation via the ubiquitin-proteasome pathway (Huang et al., 2009). Thus, by inhibiting Tankyrase activity, Axin is spared from the consequent ubiquitination and degradation, which may antagonize Wnt signaling and inhibit tumor growth. Wnt/ β -catenin signaling normally needs to be blocked for therapeutic purpose in cancer; however, it also plays an important role in hematopoietic stem cell (HSC) formation and self-renewal, and is required for HSC recovery after injury and in transplantation. Hence, our recent identification of HLY78 as a novel Wnt activator, may provide new chances for the circumstances where activation of Wnt signaling is desired, such as for the patients with bone marrow failure or undergoing post-HSC transplantation (Wang et al., 2013).

ACKNOWLEDGEMENTS

L.L and X.S would like to thank the National Basic Research Program (973 Program) (Nos. 2010CB912100 to L.L, and 2011CD966300 to X.S) and the National Natural Science Foundation of China (Grant Nos. 31230044, 30930052 and 91213304 to L.L, and 31100532 to X.S). X.S. gratefully acknowledges the support of SA-SIBS scholarship program.

ABBREVIATIONS

APC, adenomatous polyposis coli; GSK3, glycogen synthase kinase 3; HDAC, histone deacetylase; HSC, hematopoietic stem cell; NETs, neuroendocrine tumors; NSCLC, non-small cell lung cancer; Ub, ubiquitin.

COMPLIANCE WITH ETHICS GUIDELINES

Xiaomin Song, Sheng Wang and Lin Li declare that they have no conflict of interest.

This article does not contain any studies with human or animal subjects performed by the any of the authors.

OPEN ACCESS

This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

REFERENCES

- Bao R, Christova T, Song S, Angers S, Yan X, Attisano L (2012) Inhibition of tankyrases induces Axin stabilization and blocks Wnt signalling in breast cancer cells. *PLoS ONE* 7:e48670
- Bilic J, Huang YL, Davidson G, Zimmermann T, Cruciat CM, Bienz M, Niehrs C (2007) Wnt induces LRP6 signalosomes and promotes dishevelled-dependent LRP6 phosphorylation. *Science* 316:1619–1622
- Callow MG, Tran H, Phu L, Lau T, Lee J, Sandoval WN, Liu PS, Bheddah S, Tao J, Lill JR et al (2011) Ubiquitin ligase RNF146 regulates tankyrase and Axin to promote Wnt signaling. *PLoS ONE* 6:e22595
- Cha B, Kim W, Kim YK, Hwang BN, Park SY, Yoon JW, Park WS, Cho JW, Bedford MT, Jho EH (2011) Methylation by protein arginine methyltransferase 1 increases stability of Axin, a negative regulator of Wnt signaling. *Oncogene* 30:2379–2389
- Chan DW, Chan CY, Yam JW, Ching YP, Ng IO (2006) Prickle-1 negatively regulates Wnt/ β -catenin pathway by promoting Dishevelled ubiquitination/degradation in liver cancer. *Gastroenterology* 131:1218–1227
- Chen G, AJ, Wang M, Farley S, Lee LY, Lee LC, Sawicki MP (2008) Menin promotes the Wnt signaling pathway in pancreatic endocrine cells. *Mol Cancer Res* 6:1894–1907
- Chen T, Li M, Ding Y, Zhang LS, Xi Y, Pan WJ, Tao DL, Wang JY, Li L (2009) Identification of zinc-finger BED domain-containing 3 (Zbed3) as a novel Axin-interacting protein that activates Wnt/ β -catenin signaling. *J Biol Chem* 284:6683–6689
- Chia IV, Costantini F (2005) Mouse axin and axin2/conductin proteins are functionally equivalent in vivo. *Mol Cell Biol* 25:4371–4376
- Choi J, Park SY, Costantini F, Jho EH, Joo CK (2004) Adenomatous polyposis coli is down-regulated by the ubiquitin-proteasome pathway in a process facilitated by Axin. *J Biol Chem* 279:49188–49198
- Choi SH, Choi KM, Ahn HJ (2010) Coexpression and protein-protein complexing of DIX domains of human Dvl1 and Axin1 protein. *BMB Rep* 43:609–613
- Ding Y, Xi Y, Chen T, Wang JY, Tao DL, Wu ZL, Li YP, Li C, Zeng R, Li L (2008) Caprin-2 enhances canonical Wnt signaling through regulating LRP5/6 phosphorylation. *J Cell Biol* 182:865–872
- Ding Y, Zhang Y, Xu C, Tao QH, Chen YG (2013) HECT domain-containing E3 ubiquitin ligase NEDD4L negatively regulates Wnt signaling by targeting dishevelled for proteasomal degradation. *J Biol Chem* 288:8289–8298

- Egea V, Zahler S, Rieth N, Neth P, Popp T, Kehe K, Jochum M, Ries C (2012) Tissue inhibitor of metalloproteinase-1 (TIMP-1) regulates mesenchymal stem cells through let-7f microRNA and Wnt/beta-catenin signaling. *Proc Natl Acad Sci USA* 109:E309–E316
- Fang WQ, Ip JP, Li R, Ng YP, Lin SC, Chen Y, Fu AK, Ip NY (2011) Cdk5-mediated phosphorylation of Axin directs axon formation during cerebral cortex development. *J Neurosci* 31:13613–13624
- Fei C, Li Z, Li C, Chen Y, Chen Z, He X, Mao L, Wang X, Zeng R, Li L (2013) Smurf1-mediated Lys29-linked non-proteolytic poly-ubiquitination of Axin negatively regulates Wnt/beta-catenin signaling. *Mol Cell Biol* 33:4095–4105
- Fiedler M, Mendoza-Topaz C, Rutherford TJ, Mieszczynek J, Bienz M (2011) Dishevelled interacts with the DIX domain polymerization interface of Axin to interfere with its function in down-regulating beta-catenin. *Proc Natl Acad Sci USA* 108:1937–1942
- Gao ZH, Seeling JM, Hill V, Yochum A, Virshup DM (2002) Casein kinase I phosphorylates and destabilizes the beta-catenin degradation complex. *Proc Natl Acad Sci USA* 99:1182–1187
- Han Y, Zhang Y, Yang LH, Mi XY, Dai SD, Li QC, Xu HT, Yu JH, Li G, Zhao J et al (2012) X-radiation inhibits histone deacetylase 1 and 2, upregulates Axin expression and induces apoptosis in non-small cell lung cancer. *Radiat Oncol* 7:183
- Hu T, Li C, Cao Z, Van Raay TJ, Smith JG, Willert K, Solnica-Krezel L, Coffey RJ (2010) Myristoylated Naked2 antagonizes Wnt-beta-catenin activity by degrading Dishevelled-1 at the plasma membrane. *J Biol Chem* 285:13561–13568
- Huang SM, Mishina YM, Liu S, Cheung A, Stegmeier F, Michaud GA, Charlat O, Wiellette E, Zhang Y, Wiessner S et al (2009) Tankyrase inhibition stabilizes axin and antagonizes Wnt signaling. *Nature* 461:614–620
- James RG, Davidson KC, Bosch KA, Biechele TL, Robin NC, Taylor RJ, Major MB, Camp ND, Fowler K, Martins TJ et al (2012) WIKI4, a novel inhibitor of tankyrase and Wnt/ss-catenin signaling. *PLoS ONE* 7:e50457
- Jho E, Lomvardas S, Costantini F (1999) A GSK3beta phosphorylation site in axin modulates interaction with beta-catenin and Tcf-mediated gene expression. *Biochem Biophys Res Commun* 266:28–35
- Jho EH, Zhang T, Domon C, Joo CK, Freund JN, Costantini F (2002) Wnt/beta-catenin/Tcf signaling induces the transcription of Axin2, a negative regulator of the signaling pathway. *Mol Cell Biol* 22:1172–1183
- Jiang Y, Luo W, Howe PH (2009) Dab2 stabilizes Axin and attenuates Wnt/beta-catenin signaling by preventing protein phosphatase 1 (PP1)-Axin interactions. *Oncogene* 28:2999–3007
- Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, Labourier E, Reinert KL, Brown D, Slack FJ (2005) RAS is regulated by the let-7 microRNA family. *Cell* 120:635–647
- Kim S, Jho EH (2010) The protein stability of Axin, a negative regulator of Wnt signaling, is regulated by Smad ubiquitination regulatory factor 2 (Smurf2). *J Biol Chem* 285:36420–36426
- Kim SI, Park CS, Lee MS, Kwon MS, Jho EH, Song WK (2004) Cyclin-dependent kinase 2 regulates the interaction of Axin with beta-catenin. *Biochem Biophys Res Commun* 317:478–483
- Kim MJ, Chia IV, Costantini F (2008) SUMOylation target sites at the C terminus protect Axin from ubiquitination and confer protein stability. *FASEB J* 22:3785–3794
- Kim JT, Li J, Jang ER, Gulhati P, Rychahou PG, Napier DL, Wang C, Weiss HL, Lee EY, Anthony L et al (2013a) Deregulation of Wnt/beta-catenin signaling through genetic or epigenetic alterations in human neuroendocrine tumors. *Carcinogenesis* 34:953–961
- Kim SE, Huang H, Zhao M, Zhang X, Zhang A, Semonov MV, MacDonald BT, Zhang X, Garcia Abreu J, Peng L et al (2013b) Wnt stabilization of beta-catenin reveals principles for morphogen receptor-scaffold assemblies. *Science* 340:867–870
- Koinuma K, Yamashita Y, Liu W, Hatanaka H, Kurashina K, Wada T, Takada S, Kaneda R, Choi YL, Fujiwara SI et al (2006) Epigenetic silencing of AXIN2 in colorectal carcinoma with microsatellite instability. *Oncogene* 25:139–146
- Kulathu Y, Komander D (2012) Atypical ubiquitylation - the unexplored world of polyubiquitin beyond Lys48 and Lys63 linkages. *Nat Rev Mol Cell Biol* 13:508–523
- Lee I, Ajay SS, Yook JI, Kim HS, Hong SH, Kim NH, Dhanasekaran SM, Chinnaiyan AM, Athey BD (2009) New class of microRNA targets containing simultaneous 5'-UTR and 3'-UTR interaction sites. *Genome Res* 19:1175–1183
- Lee MA, Kim WK, Park HJ, Kang SS, Lee SK (2013) Anti-proliferative activity of hydnoicarpin, a natural lignan, is associated with the suppression of Wnt/beta-catenin signaling pathway in colon cancer cells. *Bioorg Med Chem Lett* 23:5511–5514
- Leung JY, Kolligs FT, Wu R, Zhai Y, Kuick R, Hanash S, Cho KR, Fearon ER (2002) Activation of AXIN2 expression by beta-catenin-T cell factor. A feedback repressor pathway regulating Wnt signaling. *J Biol Chem* 277:21657–21665
- Li VS, Ng SS, Boersema PJ, Low TY, Karthaus WR, Gerlach JP, Mohammed S, Heck AJ, Maurice MM, Mahmoudi T et al (2012) Wnt signaling through inhibition of beta-catenin degradation in an intact Axin1 complex. *Cell* 149:1245–1256
- Liu C, Kato Y, Zhang Z, Do VM, Yankner BA, He X (1999) beta-Trcp couples beta-catenin phosphorylation-degradation and regulates Xenopus axis formation. *Proc Natl Acad Sci USA* 96:6273–6278
- Liu J, Stevens J, Rote CA, Yost HJ, Hu Y, Neufeld KL, White RL, Matsunami N (2001) Siah-1 mediates a novel beta-catenin degradation pathway linking p53 to the adenomatous polyposis coli protein. *Mol Cell* 7:927–936
- Liu YT, Dan QJ, Wang J, Feng Y, Chen L, Liang J, Li Q, Lin SC, Wang ZX, Wu JW (2011) Molecular basis of Wnt activation via the DIX domain protein Ccd1. *J Biol Chem* 286:8597–8608
- Lu J, Ma Z, Hsieh JC, Fan CW, Chen B, Longgood JC, Williams NS, Amatruda JF, Lum L, Chen C (2009) Structure-activity relationship studies of small-molecule inhibitors of Wnt response. *Bioorg Med Chem Lett* 19:3825–3827
- Lui TT, Lacroix C, Ahmed SM, Goldenberg SJ, Leach CA, Daulat AM, Angers S (2011) The ubiquitin-specific protease USP34 regulates axin stability and Wnt/beta-catenin signaling. *Mol Cell Biol* 31:2053–2065
- Luo W, Peterson A, Garcia BA, Coombs G, Kofahl B, Heinrich R, Shabanowitz J, Hunt DF, Yost HJ, Virshup DM (2007) Protein phosphatase 1 regulates assembly and function of the beta-catenin degradation complex. *EMBO J* 26:1511–1521
- Mao J, Wang J, Liu B, Pan W, Farr GH 3rd, Flynn C, Yuan H, Takada S, Kimelman D, Li L et al (2001) Low-density lipoprotein receptor-

- related protein-5 binds to Axin and regulates the canonical Wnt signaling pathway. *Mol Cell* 7:801–809
- Mayr C, Hemann MT, Bartel DP (2007) Disrupting the pairing between let-7 and Hmga2 enhances oncogenic transformation. *Science* 315:1576–1579
- Mendoza-Topaz C, Mieszczanek J, Bienz M (2011) The Adenomatous polyposis coli tumour suppressor is essential for Axin complex assembly and function and opposes Axin's interaction with Dishevelled. *Open Biol* 1:110013
- Metcalfe C, Mendoza-Topaz C, Mieszczanek J, Bienz M (2010) Stability elements in the LRP6 cytoplasmic tail confer efficient signalling upon DIX-dependent polymerization. *J Cell Sci* 123:1588–1599
- Miyaki M, Konishi M, Kikuchi-Yanoshita R, Enomoto M, Igari T, Tanaka K, Muraoka M, Takahashi H, Amada Y, Fukayama M et al (1994) Characteristics of somatic mutation of the adenomatous polyposis coli gene in colorectal tumors. *Cancer Res* 54:3011–3020
- Najdi R, Holcombe RF, Waterman ML (2011) Wnt signaling and colon carcinogenesis: beyond APC. *J Carcinog* 10:5
- Nikuseva Martic T, Pecina-Slaus N, Kusec V, Kokotovic T, Musinovic H, Tomas D, Zeljko M (2010) Changes of AXIN-1 and beta-catenin in neuroepithelial brain tumors. *Pathol Oncol Res* 16:75–79
- Pecina-Slaus N, Martic TN, Kokotovic T, Kusec V, Tomas D, Hrascan R (2011) AXIN-1 protein expression and localization in glioblastoma. *Collegium Antropologicum* 35(Suppl 1):101–106
- Reinhart BJ, Slack FJ, Basson M, Pasquinelli AE, Bettinger JC, Rougvie AE, Horvitz HR, Ruvkun G (2000) The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature* 403:901–906
- Rui HL, Fan E, Zhou HM, Xu Z, Zhang Y, Lin SC (2002) SUMO-1 modification of the C-terminal KVEKVD of Axin is required for JNK activation but has no effect on Wnt signaling. *J Biol Chem* 277:42981–42986
- Sakanaka C, Williams LT (1999) Functional domains of axin. Importance of the C terminus as an oligomerization domain. *J Biol Chem* 274:14090–14093
- Schmitz Y, Rateitschak K, Wolkenhauer O (2013) Analysing the impact of nucleo-cytoplasmic shuttling of beta-catenin and its antagonists APC, Axin and GSK3 on Wnt/beta-catenin signalling. *Cell Signal* 25:2210–2221
- Schwarz-Romond T, Fiedler M, Shibata N, Butler PJ, Kikuchi A, Higuchi Y, Bienz M (2007a) The DIX domain of Dishevelled confers Wnt signaling by dynamic polymerization. *Nat Struct Mol Biol* 14:484–492
- Schwarz-Romond T, Metcalfe C, Bienz M (2007b) Dynamic recruitment of axin by Dishevelled protein assemblies. *J Cell Sci* 120:2402–2412
- Shibata N, Tomimoto Y, Hanamura T, Yamamoto R, Ueda M, Ueda Y, Mizuno N, Ogata H, Komori H, Shomura Y et al (2007) Crystallization and preliminary X-ray crystallographic studies of the axin DIX domain. *Acta Crystallogr Sect F Struct Biol Cryst Commun* 63:529–531
- Shiomi K, Uchida H, Keino-Masu K, Masu M (2003) Ccd1, a novel protein with a DIX domain, is a positive regulator in the Wnt signaling during zebrafish neural patterning. *Curr Biol* 13:73–77
- Shultz MD, Kirby CA, Stams T, Chin DN, Blank J, Charlat O, Cheng H, Cheung A, Cong F, Feng Y et al (2012) [1,2,4]triazol-3-ylsulfanylmethyl-3-phenyl-[1,2,4]oxadiazoles: antagonists of the Wnt pathway that inhibit tankyrases 1 and 2 via novel adenosine pocket binding. *J Med Chem* 55:1127–1136
- Silver SJ, Hagen JW, Okamura K, Perrimon N, Lai EC (2007) Functional screening identifies miR-315 as a potent activator of Wingless signaling. *Proc Natl Acad Sci USA* 104:18151–18156
- Strovel ET, Wu D, Sussman DJ (2000) Protein phosphatase 2C α dephosphorylates axin and activates LEF-1-dependent transcription. *J Biol Chem* 275:2399–2403
- Tran H, Polakis P (2012) Reversible modification of adenomatous polyposis coli (APC) with K63-linked polyubiquitin regulates the assembly and activity of the beta-catenin destruction complex. *J Biol Chem* 287:28552–28563
- Tran H, Bustos D, Yeh R, Rubinfeld B, Lam C, Shriver S, Zilberleyb I, Lee MW, Phu L, Sarkar AA et al (2013) HectD1 E3 ligase modifies adenomatous polyposis coli (APC) with polyubiquitin to promote the APC-axin interaction. *J Biol Chem* 288:3753–3767
- Wang S, Yin J, Chen D, Nie F, Song X, Fei C, Miao H, Jing C, Ma W, Wang L et al (2013) Small-molecule modulation of Wnt signaling via modulating the Axin-LRP5/6 interaction. *Nat Chem Biol* 9:579–585
- Wei W, Li M, Wang J, Nie F, Li L (2012) The E3 ubiquitin ligase ITCH negatively regulates canonical Wnt signaling by targeting dishevelled protein. *Mol Cell Biol* 32:3903–3912
- Willert K, Jones KA (2006) Wnt signaling: is the party in the nucleus? *Genes Dev* 20:1394–1404
- Willert K, Shibamoto S, Nusse R (1999) Wnt-induced dephosphorylation of axin releases beta-catenin from the axin complex. *Genes Dev* 13:1768–1773
- Winston JT, Strack P, Beer-Romero P, Chu CY, Elledge SJ, Harper JW (1999) The SCF β -TRCP-ubiquitin ligase complex associates specifically with phosphorylated destruction motifs in I κ B α and beta-catenin and stimulates I κ B α ubiquitination in vitro. *Genes Dev* 13:270–283
- Wong CK, Luo W, Deng Y, Zou H, Ye Z, Lin SC (2004) The DIX domain protein coiled-coil-DIX1 inhibits c-Jun N-terminal kinase activation by Axin and dishevelled through distinct mechanisms. *J Biol Chem* 279:39366–39373
- Yamamoto H, Kishida S, Kishida M, Ikeda S, Takada S, Kikuchi A (1999) Phosphorylation of axin, a Wnt signal negative regulator, by glycogen synthase kinase-3 β regulates its stability. *J Biol Chem* 274:10681–10684
- Yan D, Wiesmann M, Rohan M, Chan V, Jefferson AB, Guo L, Sakamoto D, Caohien RH, Fuller JH, Reinhard C et al (2001) Elevated expression of axin2 and hnkcd mRNA provides evidence that Wnt/beta-catenin signaling is activated in human colon tumors. *Proc Natl Acad Sci USA* 98:14973–14978
- Yang LH, Xu HT, Li QC, Jiang GY, Zhang XP, Zhao HY, Xu K, Wang EH (2013) Abnormal hypermethylation and clinicopathological significance of Axin gene in lung cancer. *Tumour Biol* 34:749–757
- Yokoyama N, Markova NG, Wang HY, Malbon CC (2012) Assembly of Dishevelled 3-based supermolecular complexes via phosphorylation and Axin. *J Mol Signal* 7:8
- Zhang Y, Liu S, Mickanin C, Feng Y, Charlat O, Michaud GA, Schirle M, Shi X, Hild M, Bauer A et al (2011) RNF146 is a poly(ADP-ribose)-directed E3 ligase that regulates axin degradation and Wnt signalling. *Nat Cell Biol* 13:623–629