

A segment-wise dynamic programming algorithm for BSDEs

Christian Bender, Steffen Meyer

Department of Mathematics, Saarland University, Campus E 2 4, D-66123 Saarbrücken, Germany

Email: bender@math.uni-saarland.de, meyersteffen03@gmail.com

Abstract We introduce and analyze a family of linear least-squares Monte Carlo schemes for backward SDEs, which interpolate between the one-step dynamic programming scheme of Lemor, Warin, and Gobet (Bernoulli, 2006) and the multi-step dynamic programming scheme of Gobet and Turkedjiev (Mathematics of Computation, 2016). Our algorithm approximates conditional expectations over segments of the time grid. We discuss the optimal choice of the segment length depending on the ‘smoothness’ of the problem and show that, in typical situations, the complexity can be reduced compared to the state-of-the-art multi-step dynamic programming scheme.

Keywords Backward stochastic differential equations, Empirical regression, Dynamic programming, Monte Carlo methods

2020 Mathematics Subject Classification 65C30; 60H10; 62G08

1. Introduction

In this paper, we propose a family of regression-based algorithms for decoupled forward backward stochastic differential equations (FBSDEs) of the form

$$\begin{aligned} dX_t &= b(t, X_t)dt + \sigma(t, X_t)dW_t, & X_0 &= x_0, \\ -dY_t &= f(t, X_t, Y_t, Z_t)dt - Z_t dW_t, & Y_T &= \xi(X_T) \end{aligned} \quad (1)$$

driven by a (multidimensional) Brownian motion W . Recall that the solution is a triplet (X, Y, Z) of adapted processes, wherein X and Y describe the dynamics of the forward and backward SDE (BSDE), respectively, and Z steers the backward SDE into the terminal condition without making use of future information. Among the numerous applications of BSDEs we mention (nonlinear) derivative pricing problems, drift control problems, and stochastic representation formulas of Feynman-Kac type for semilinear parabolic partial differential equations, see, e.g., [7, 12, 22]. Loosely speaking, in these applications, the Y -part of the solution corresponds to the price process of the option, the value of the control problem, or the solution of the PDE, respectively, while the Z -part is required to represent the hedging portfolio, an optimal control, or the derivative of the PDE solution.

Motivated by these and other applications, the literature on numerical schemes for BSDEs has tremendously increased over the last two decades, and we refer to [3] for a comprehensive survey.

Let us emphasize that the practical choice of a numerical scheme for the FBSDE (1) crucially depends on the dimension D of the state space of the diffusion process X . If X takes values in a low dimensional space, the connection to PDEs via the four-step scheme [19] can be exploited and the application of fast PDE solvers leads to highly efficient schemes for the FBSDE, see, e.g., [5, 20, 21]. For moderate dimensions (up to about $D = 10$), simulation based regression methods are among the most popular methods [9, 11, 18], while the recent multilevel Picard approach can tackle very high-dimensional problems [6, 14].

In the present paper, we contribute to the theory of simulation-based regression schemes. More precisely, we design a family of schemes which interpolate between the one-step-dynamic programming approach (ODP) of [18] and the multi-step dynamic programming approach (MDP) of [11]. Both dynamic programming schemes rely on a backward recursion for the time discretization with nested conditional expectations which are approximated by empirical least-squares regression. The key difference is that the ODP computes the numerical solution of the BSDE at a time grid point t_i by regressing a nonlinear function of the numerical solution at the previous time grid point t_{i+1} only, while the MDP applies the numerical solutions at all grid points t_{i+1}, t_{i+2}, \dots up to terminal time T . As argued in [11] (see also [1], for related considerations), the error propagation over time for the approximation of the Y -part of the solution can be significantly reduced in the MDP making it superior to the ODP in typical situations. A closer look at both schemes reveals, however, that the approximation of the Y -part is the dominating cost in the ODP, while the cost for approximating the Z -part dominates the complexity of the MDP. In order to balance the cost for approximating Y -part and Z -part, we suggest a family of schemes, which iterates between a single ODP step and an MDP step on time segments containing $\lceil N^\alpha \rceil$ grid points—here N is the total number of grid points and $\alpha \in [0, 1]$. We call this approach ‘segmentwise dynamic programming’ (SDP) and analyze it in detail in this paper. In particular, we discuss how to optimize the choice of the segment length parameter α in dependence on the state space dimension D and on the smoothness of the PDE representation to the BSDE. It turns out that, in generic situations, the optimal choice of α is in the open interval $(0, 1)$, demonstrating that neither the ODP ($\alpha = 0$) nor the MDP ($\alpha = 1$) are optimal. For the theoretical analysis, we assume that the forward diffusion X can be perfectly simulated on the discrete time grids. One can, however, apply a higher order scheme for X on the coarser grid to implement the SDP scheme, when this assumption is not satisfied.

The paper is structured as follows: In Section 2 we introduce the setting and motivate the SDP algorithm, which is spelled out in detail in Section 3. The theoretical main results on convergence of the SDP scheme are presented in Section 4 along with a complexity analysis. The convergence behavior of the SDP scheme is illustrated and compared to the MDP scheme by a numerical experiment in Section 5. Finally, the detailed error analysis (combining ideas from [1, 11, 18]) is carried out in Section 6. Some proofs, which follow standard arguments, are provided in the online supplement.

2. Setup and motivation

Let $(\Omega, \mathcal{F}, \mathbb{F}, P)$ be a filtered probability where the filtration is the augmentation of the one generated by a \mathcal{D} -dimensional Brownian motion W . We consider a decoupled forward backward stochastic differential equation of the form

$$\begin{aligned} dX_t &= b(t, X_t)dt + \sigma(t, X_t)dW_t, & X_0 &= x_0, \\ -dY_t &= f(t, X_t, Y_t, Z_t)dt - Z_t dW_t, & Y_T &= \xi(X_T) \end{aligned}$$

with deterministic initial value $x_0 \in \mathbb{R}^D$ and the terminal time $T > 0$ under the following standing assumptions.

Assumptions 2.1

(A_ξ) The function $\xi : \mathbb{R}^D \rightarrow \mathbb{R}$ is bounded by some constant C_ξ .

(A_L) The functions $b : [0, T] \times \mathbb{R}^D \rightarrow \mathbb{R}^D$, $\sigma : [0, T] \times \mathbb{R}^D \rightarrow \mathbb{R}^{D \times D}$ and $f : [0, T] \times \mathbb{R}^D \times \mathbb{R} \times \mathbb{R}^D \rightarrow \mathbb{R}$ are measurable, $\frac{1}{2}$ -Hölder-continuous in the first variable and Lipschitz-continuous in the other variables, i.e., there exist constants L_X and L_f such that

$$\begin{aligned} |b(t, x) - b(t', x')| + |\sigma(t, x) - \sigma(t', x')| &\leq L_X \left(|t - t'|^{\frac{1}{2}} + |x - x'| \right), \\ |f(t, x, y, z) - f(t', x', y', z')| &\leq L_f \left(|t - t'|^{\frac{1}{2}} + |x - x'| + |y - y'| + |z - z'| \right) \end{aligned}$$

for all $x, x' \in \mathbb{R}^D$, $t, t' \in [0, T]$, $y, y' \in \mathbb{R}$, $z, z' \in \mathbb{R}^D$.

(A_f) The function f is uniformly bounded by a constant C_f , i.e.,

$$|f(t, x, y, z)| \leq C_f$$

for all $t \in [0, T]$, $x \in \mathbb{R}^D$, $y \in \mathbb{R}$, $z \in \mathbb{R}^D$.

Assumption (A_L) is standard for FBSDEs and yields important characteristics of the processes X , Y and Z . For once, the assumption on b and σ ensures the existence of a unique strong solution X of the SDE and that this solution satisfies $E[\sup_{t \in [0, T]} |X_t|^2] < \infty$ (see e.g. [16]). Then, paired with the assumptions on f and ξ the solution of the BSDE can be expressed by deterministic functions of the SDE solution X , i.e., there exist deterministic functions $\bar{y} : [0, T] \times \mathbb{R}^D \rightarrow \mathbb{R}$ and $\bar{z} : [0, T] \times \mathbb{R}^D \rightarrow \mathbb{R}^D$ such that $\bar{y}(t, X_t) = Y_t$ and $\bar{z}(t, X_t) = Z_t$ (see e.g. [7]). The boundedness conditions on f and ξ are posed for convenience only and could be relaxed with minor changes in the error analysis.

We now define the equidistant time grid

$$\pi := \{t_i = i\Delta; i = 0, \dots, N\}$$

with step width $\Delta = T/N$ for a fixed $N \in \mathbb{N}$. We denote the increments of the Brownian motion W on this time grid with ΔW_i , i.e., $\Delta W_i := W_{t_i} - W_{t_{i-1}}$, and define the functions

$$\begin{aligned} \bar{q}_i^N(x) &:= E[Y_{t_{i+1}} | X_{t_i} = x], \\ \bar{z}_i^N(x) &:= E\left[\frac{\Delta W_{i+1}}{\Delta} Y_{t_{i+1}} \middle| X_{t_i} = x\right] \end{aligned}$$

for all $i \in \{0, \dots, N-1\}$. Then, under the standing assumptions, it holds that

$$\lim_{N \rightarrow \infty} \left(\max_{i=0, \dots, N-1} E[|\bar{q}_i^N(X_{t_i}) - Y_{t_i}|^2] + \sum_{i=0}^{N-1} E\left[\int_{t_i}^{t_{i+1}} |\bar{z}_i^N(X_{t_i}) - Z_s|^2 ds\right] \right) = 0,$$

where the rate of convergence depends on the regularity of Z , see e.g. [23]. The functions \bar{q}_i^N and \bar{z}_i^N can, therefore, be interpreted as time-discretized versions of the BSDE solution (Y, Z) . To obtain an implementable approximation scheme for \bar{q}_i^N and \bar{z}_i^N , fix any function

$$\tau : \{0, \dots, N-2\} \rightarrow \{1, \dots, N-1\}$$

satisfying $\tau(i) \geq i+1$ for all $i \in \{0, \dots, N-2\}$. Then the tower property of the conditional expectation and the Markov property of X yield for any $i \in \{0, \dots, N-2\}$

$$\begin{aligned}
\bar{q}_i^N(x) &= E[Y_{t_{i+1}} | X_{t_i} = x] \\
&= E \left[Y_{t_{\tau(i)+1}} + \int_{t_{i+1}}^{t_{\tau(i)+1}} f(t, X_t, Y_t, Z_t) dt - \int_{t_{i+1}}^{t_{\tau(i)+1}} Z_t dW_t \middle| X_{t_i} = x \right] \\
&= E \left[\bar{q}_{\tau(i)}^N(X_{\tau(i)}) + \sum_{j=i+1}^{\tau(i)} \int_{t_j}^{t_{j+1}} f(t, X_t, Y_t, Z_t) dt \middle| X_{t_i} = x \right] \\
&\approx E \left[\bar{q}_{\tau(i)}^N(X_{\tau(i)}) + \sum_{j=i+1}^{\tau(i)} f(t_j, X_{t_j}, \bar{q}_j^N(X_{t_j}), \bar{z}_j^N(X_{t_j})) \Delta \middle| X_{t_i} = x \right],
\end{aligned}$$

and similarly

$$\begin{aligned}
\bar{z}_i^N(x) &= E \left[\frac{\Delta W_{i+1}}{\Delta} Y_{t_{i+1}} \middle| X_{t_i} = x \right] \\
&= E \left[\frac{\Delta W_{i+1}}{\Delta} \left(Y_{t_{\tau(i)+1}} + \int_{t_{i+1}}^{t_{\tau(i)+1}} f(t, X_t, Y_t, Z_t) dt - \int_{t_{i+1}}^{t_{\tau(i)+1}} Z_t dW_t \right) \middle| X_{t_i} = x \right] \\
&= E \left[\frac{\Delta W_{i+1}}{\Delta} \left(\bar{q}_{\tau(i)}^N(X_{\tau(i)}) + \sum_{j=i+1}^{\tau(i)} \int_{t_j}^{t_{j+1}} f(t, X_t, Y_t, Z_t) dt \right) \middle| X_{t_i} = x \right] \\
&\approx E \left[\frac{\Delta W_{i+1}}{\Delta} \left(\bar{q}_{\tau(i)}^N(X_{\tau(i)}) + \sum_{j=i+1}^{\tau(i)} f(t_j, X_{t_j}, \bar{q}_j^N(X_{t_j}), \bar{z}_j^N(X_{t_j})) \Delta \right) \middle| X_{t_i} = x \right].
\end{aligned}$$

This motivates the time discretization scheme

$$\begin{aligned}
Q_{N-1}^N &:= E [\xi(X_{t_N}) | \mathcal{F}_{t_{N-1}}], \\
Z_{N-1}^N &:= E \left[\frac{\Delta W_N}{\Delta} \xi(X_{t_N}) \middle| \mathcal{F}_{t_{N-1}} \right], \\
Q_i^N &:= E \left[Q_{\tau(i)}^N + \sum_{j=i+1}^{\tau(i)} f(t_j, X_{t_j}, Q_j^N, Z_j^N) \Delta \middle| \mathcal{F}_{t_i} \right], \quad i = N-2, \dots, 0, \\
Z_i^N &:= E \left[\frac{\Delta W_{i+1}}{\Delta} \left(Q_{\tau(i)}^N + \sum_{j=i+1}^{\tau(i)} f(t_j, X_{t_j}, Q_j^N, Z_j^N) \Delta \right) \middle| \mathcal{F}_{t_i} \right], \quad i = N-2, \dots, 0.
\end{aligned} \tag{2}$$

By the tower property of the conditional expectation, this definition of Q_i^N and Z_i^N does not depend on the choice of τ and is nothing but a reformulation of the backward Euler scheme by [2, 23]. However, the conditional expectations cannot be calculated in closed form in general. Hence, when attempting to solve the BSDE, one has to replace the conditional expectations with some approximation operator resulting in different schemes depending on the choice of τ . As our results will show, the choice of τ then influences both, the computational costs as well as the convergence properties.

The most natural choices for τ would be for once setting $\tau(i) = i+1$ or $\tau(i) = N-1$ for all $i \in \{0, \dots, N-2\}$. The first results in the classical one-step scheme of [18] (to which we further refer to as ODP), the latter in the multi-step forward scheme (MDP for short) by [11]. To understand the idea of the segment-wise dynamic programming algorithm that will be introduced in the next section, it is worth reviewing these two schemes and comparing the resulting algorithms.

Both algorithms work recursively backward in time by constructing estimates of the functions

\bar{q}_i^N and \bar{z}_i^N through approximating the conditional expectations in the corresponding time discretization scheme via empirical (i.e., simulation-based) orthogonal projections on finite-dimensional function spaces, where the components Q_j^N, Z_j^N with $j > i$ on the right-hand side of the discretization scheme are replaced by the approximations found in the previous steps of the recursion. As a result of the different schemes, the approximation of the ODP algorithm depends at each time point t_i only on the approximations at the time t_{i+1} while in the MDP scheme, the approximation at each step t_i depends on all the previously constructed ones, i.e., those at the time points from t_{i+1} up to t_{N-1} . Since the approximations of \bar{q}_i^N and \bar{z}_i^N have to be evaluated, one has to simulate in each step of the MDP algorithm segments of the form $(X_{t_i}, X_{t_{i+1}}, \dots, X_{t_N})$ while it suffices in the ODP algorithm to simulate values of X at just the current and the following time point. This obviously leads to higher simulation costs in the MDP scheme. However, since the algorithms recursively reuse the obtained approximations of \bar{q}_i^N and \bar{z}_i^N an error propagation between the time steps occurs. The error analysis in [11] reveals that, from this perspective, the MDP scheme features improved convergence properties compared to the ODP.

The idea of the segment-wise approach, which will be introduced in the next chapter, is to interpolate between the two cases of the ODP and the MDP scheme in order to balance these two aspects, the computational costs and convergence properties.

3. Segment-wise dynamic programming algorithm

In this section, we present the segment-wise dynamic programming algorithm (SDP, for short) in detail. We first specify a family of functions τ_α to obtain the time discretization scheme that interpolates between the ones from the ODP scheme and the MDP scheme via (2). Then the algorithm is described in detail based on this discretization scheme.

For any $\alpha \in [0, 1]$, consider the time grid

$$\bar{\pi}_\alpha := \{(\Delta n \lceil N^\alpha \rceil) \wedge (T - \Delta); n \in \mathbb{N}\}$$

with step width $\Delta \lceil N^\alpha \rceil$ (up to a possibly smaller size in the last step), that consists of $\lceil N^{1-\alpha} \rceil$ time points at most. Based on these time grids define the functions

$$\tau_\alpha : \{0, 1, \dots, N-2\} \rightarrow \{1, \dots, N-1\}$$

as

$$\tau_\alpha(i) := \min\{j > i : j\Delta \in \bar{\pi}_\alpha\}.$$

For a fixed α , the choice $\tau = \tau_\alpha$ in (2) then defines a discretization scheme where the time grid π is separated in segments consisting of $\lceil N^\alpha \rceil$ points by the coarser time grid $\bar{\pi}_\alpha$. The resulting discretization scheme corresponds to an MDP scheme on each of these segments paired with a single step of an ODP scheme between consecutive segments connecting them. Moreover choosing $\alpha = 0$ or $\alpha = 1$ results in the classical ODP or MDP scheme respectively.

Now for a fixed $\alpha \in [0, 1]$ the SDP algorithm works as follows.

Algorithm 3.1

- Choose basis functions

$$\begin{aligned} p_{q,i}^k &: \mathbb{R}^D \rightarrow \mathbb{R}, \quad k = 1, \dots, K_{q,i}, \\ p_{z,i}^k &: \mathbb{R}^D \rightarrow \mathbb{R}^D, \quad k = 1, \dots, K_{z,i}, \end{aligned}$$

for each $i \in \{0, \dots, N-1\}$ such that

$$\sum_{i=0}^{N-1} \sum_{k=1}^{K_{q,i}} E [|p_{q,i}^k(X_{t_i})|^2] + \sum_{i=0}^{N-1} \sum_{k=1}^{K_{z,i}} E [|p_{z,i}^k(X_{t_i})|^2] < \infty.$$

Here the number of basis functions $K_{q,i}, K_{z,i} \in \mathbb{N}$ may depend on the time point t_i . We denote the function spaces spanned by these basis functions with $\mathcal{K}_{q,i}$ and $\mathcal{K}_{z,i}$ respectively, i.e.,

$$\begin{aligned} \mathcal{K}_{q,i} &:= \text{span} \left(p_{q,i}^1, \dots, p_{q,i}^{K_{q,i}} \right), \\ \mathcal{K}_{z,i} &:= \text{span} \left(p_{z,i}^1, \dots, p_{z,i}^{K_{z,i}} \right). \end{aligned}$$

The algorithm will approximate \bar{q}_i^N by empirical orthogonal projections on the subspaces $\mathcal{K}_{q,i}$ and \bar{z}_i^N by projections on $\mathcal{K}_{z,i}$.

- Initialize the algorithm by setting

$$\Xi_{N-1}^{N,M}(x_N) := \xi(x_N)$$

for all $x_N \in \mathbb{R}^D$. Then, assuming $\Xi_i^{N,M}$ is already constructed, perform the following backward recursion for $i = N-1, N-2, \dots, 0$:

- 1*) If $i = N-1$: Choose $M_{N-1} \in \mathbb{N}$, then simulate M_{N-1} independent copies

$$\left(X_{t_{N-1}}^{[N-1,m,N]}, X_{t_N}^{[N-1,m,N]}, \Delta W_N^{[N-1,m,N]} \right)_{m=1, \dots, M_{N-1}}$$

of the segment $(X_{t_{N-1}}, X_{t_N}, \Delta W_N)$ and set

$$X^{[N-1,m,N]} := X_{t_N}^{[N-1,m,N]}.$$

- 1) If $i < N-1$: Choose a $M_i \in \mathbb{N}$, then simulate M_i independent copies

$$\left(X_{t_i}^{[i,m,N]}, \dots, X_{t_{\tau_\alpha(i)}}^{[i,m,N]}, \Delta W_{i+1}^{[i,m,N]} \right)_{m=1, \dots, M_i}$$

of the segment $(X_{t_i}, \dots, X_{t_{\tau_\alpha(i)}}, \Delta W_{i+1})$ and set

$$X^{[i,m,N]} := \left(X_{t_{i+1}}^{[i,m,N]}, \dots, X_{t_{\tau_\alpha(i)}}^{[i,m,N]} \right).$$

- 2) Find solutions to the linear least-squares regression problems

$$\varphi_i^{q^{N,M}} = \underset{\psi \in \mathcal{K}_{q,i}}{\text{argmin}} \left(\frac{1}{M} \sum_{m=1}^{M_i} \left| \psi \left(X_{t_i}^{[i,m,N]} \right) - \Xi_i^{N,M} \left(X_{t_i}^{[i,m,N]} \right) \right|^2 \right)$$

and

$$\varphi_i^{z^{N,M}} = \underset{\psi \in \mathcal{K}_{z,i}}{\text{argmin}} \left(\frac{1}{M} \sum_{m=1}^{M_i} \left| \psi \left(X_{t_i}^{[i,m,N]} \right) - \frac{\Delta W_{i+1}^{[i,m,N]}}{\Delta} \Xi_i^{N,M} \left(X_{t_i}^{[i,m,N]} \right) \right|^2 \right).$$

- 3) Define approximations $q_i^{N,M}$ and $z_i^{N,M}$ of the functions \bar{q}_i^N and \bar{z}_i^N via

$$q_i^{N,M} := \mathcal{T}_{C_{q,i}} \circ \varphi_i^{q^{N,M}}, \quad z_i^{N,M} := \mathcal{T}_{C_{z,i}} \circ \varphi_i^{z^{N,M}},$$

where $C_{q,i} := C_\xi + (T - t_{i+1})C_f$ and $C_{z,i} := \frac{C_{q,i}}{\Delta}$ are positive constants and \mathcal{T}_c is the truncation function defined as

$$\mathcal{T}_c(x) := \text{sign}(x) \min\{|x|, c\}$$

for any constant $c > 0$ (acting componentwise on $\varphi^{z^{N,M}}$).

4) If $i \geq 1$, set

$$\Xi_{i-1}^{N,M}(x_i, \dots, x_{\tau_\alpha(i-1)}) := q_{\tau_\alpha(i-1)}^{N,M}(x_{\tau_\alpha(i-1)}) + \sum_{j=i}^{\tau_\alpha(i-1)} \Delta f(t_j, x_j, q_j^{N,M}(x_j), z_j^{N,M}(x_j))$$

as preparation for the next iteration.

The solutions to the empirical least squares problems can be computed numerically using a singular value decomposition. Hence, the algorithm is fully implementable as long as the segments $(X_{t_i}, \dots, X_{t_{\tau_\alpha(i)}})$ can be simulated. Then, in the typical situation, for example $X = W$, the average costs for the simulation of one segment $(X_{t_i}, \dots, X_{t_{\tau_\alpha(i)}})$ are of order $\mathcal{O}(N^\alpha)$. For $\alpha < 1$ this leads to smaller computation costs through simulations as the MDP scheme, where the average costs for simulating one set $(X_{t_i}, \dots, X_{t_N})$ are of order $\mathcal{O}(N)$. When X can not be sampled perfectly, it would, in principle, be possible to replace X with some approximation scheme with minor changes in the error analysis. The problem is, however, to approximate X in a way that sustains the gain in computation costs compared to the MDP scheme, which would not be the case in the simplest approach when approximating X with a naive Euler scheme starting at the time 0. In theory, one could approximate X_{t_i} with some high-order approximation scheme [17] and use an Euler scheme inside the segment $(X_{t_i}, \dots, X_{t_{\tau_\alpha(i)}})$ and preserve at least some gain in computation costs. However, we restrict the theoretical analysis to the assumption that the values of X can be sampled directly on the time grid π .

4. Convergence rates and analysis of the complexity

In this section, we state error bounds for the quadratic error of the SDP algorithm. We then analyze how to optimally calibrate the parameters of the algorithm to the smoothness of the problem. The results show that the optimal choice of the segment length parameter α is always in the open interval $(0, 1)$, and hence the SDP algorithm presented in Section 3 is advantageous when compared to the MDP and ODP schemes.

Our first bound for the mean-squared error of the SDP scheme is given in the following theorem:

Theorem 4.1 *Under the standing assumptions,*

$$\begin{aligned} & \max_{0 \leq i \leq N-1} E \left[|q_i^{N,M}(X_{t_i}) - \bar{q}_i^N(X_{t_i})|^2 \right] + \sum_{i=0}^{N-1} \Delta E \left[|z_i^{N,M}(X_{t_i}) - \bar{z}_i^N(X_{t_i})|^2 \right] \\ & \leq c \max_{i \in \mathcal{J}} \left(N^{1-\alpha} \inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\psi(X_{t_i}) - \bar{q}_i^N(X_{t_i})|^2 \right] + N^{2-2\alpha} \frac{K_{q,i}}{M_i} + N^{2-2\alpha} \frac{K_{q,i} \log(M_i)}{M_i} \right) \\ & \quad + c \max_{0 \leq i \leq N-1} \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\psi(X_{t_i}) - \bar{q}_i^N(X_{t_i})|^2 \right] + \inf_{\psi \in \mathcal{K}_{z,i}} E \left[|\psi(X_{t_i}) - \bar{z}_i^N(X_{t_i})|^2 \right] \right) \\ & \quad + \frac{K_{q,i}}{M_i} + N \frac{K_{z,i}}{M_i} + \frac{K_{q,i} \log(M_i)}{M_i} + N \frac{K_{z,i} \log(M_i)}{M_i} \Big) + cN\mathcal{R}^N, \end{aligned}$$

where $\mathcal{J} := \{i : t_i \in \bar{\pi}_\alpha\}$, c is a positive constant not depending on N and

$$\mathcal{R}^N := \sum_{i=1}^{N-1} E \left[\left(\int_{t_i}^{t_{i+1}} E \left[f(s, X_s, Y_s, Z_s) - f(t_i, X_{t_i}, \bar{q}_i^N(X_{t_i}), \bar{z}_i^N(X_{t_i})) \mid X_{t_{i-1}} \right] ds \right)^2 \right].$$

As usual, we can think of this bound as a composition of terms due to three different error sources. The first source of error is due to the projection on the finite-dimensional subspaces. This

projection error can be controlled by the choice of basis functions. The term \mathcal{R}^N only depends on the continuous-time BSDE solution and the finer time grid (through the discretized functions \bar{q}_i^N and \bar{z}_i^N), but not on the approximation obtained by the algorithm. It can be interpreted as part of the time discretization error. The remaining terms are statistical error terms depending on the sample size. Those can be controlled by increasing the number of simulations in dependence of the number of basis functions.

The bound shows the influence of the parameter α , as the projection and statistical error terms regarding \bar{q}^N appear once at all time steps and once on the time steps of the coarser time grid $\bar{\pi}$ with different factors that are decreasing in α . Although larger values for α appear to be favorable from this perspective, increasing the parameter α also results in higher computational costs as paths then have to be simulated on larger segments. Hence, when optimizing α a trade-off between convergence properties and computational costs has to be taken into account.

For $\alpha = 1$ the terms in the first bracket are dominated by the remaining terms, and we essentially reproduce the error analysis of the MDP scheme in [11] with some minor differences: In particular, the projection error in Theorem 4.1 is formulated in terms of the true continuous time solution (Y, Z) of the BSDE via the functions \bar{q}_i^N and \bar{z}_i^N while it is stated in terms of the backward Euler discretization scheme for BSDEs in [11]. When choosing $\alpha = 0$, one ends up with an error analysis for the ODP scheme with independent re-simulation at every time point. This differs from the error analysis in [18], as they only apply one cloud of simulations, which is re-used at any time point, resulting in an additional (and dominating) interdependency error. Hence, the bound obtained with Theorem 4.1 for $\alpha = 0$ also allows for a comparison of the ODP and MDP algorithms in a unified setting.

Typical bounds for the term \mathcal{R}^N in dependence of N are of order N^{-2} or N^{-3} , depending on regularity assumptions as illustrated by the following theorems. These theorems can be derived from Theorem 4.1 by applying standard techniques to bound the expected quadratic difference between the functions \bar{q}^N and \bar{z}^N and their continuous time counterparts \bar{y} and \bar{z} depending on the regularity of the BSDE, see [8, 23]. Detailed proofs are provided in an online supplement¹.

Theorem 4.2 *Under the standing assumptions, suppose that \bar{z} is $\frac{1}{2}$ -Hölder continuous in t and Lipschitz continuous in x . Then,*

$$\begin{aligned} & \max_{0 \leq i \leq N-1} E \left[|q_i^{N,M}(X_{t_i}) - \bar{y}(t_i, X_{t_i})|^2 \right] + \sum_{i=0}^{N-1} E \left[\int_{t_i}^{t_{i+1}} |z_i^{N,M}(X_{t_i}) - \bar{z}(s, X_s)|^2 ds \right] \\ & \leq c \max_{i \in \mathcal{J}} \left(N^{1-\alpha} \inf_{\psi \in \mathcal{K}_{q,i}} E [|\psi(X_{t_i}) - \bar{y}(t_i, X_{t_i})|^2] + N^{2-2\alpha} \frac{K_{q,i}}{M_i} + N^{2-2\alpha} \frac{K_{q,i} \log(M_i)}{M_i} \right) \\ & \quad + c \max_{0 \leq i \leq N-1} \left(\inf_{\psi \in \mathcal{K}_{q,i}} E [|\psi(X_{t_i}) - \bar{y}(t_i, X_{t_i})|^2] + \inf_{\psi \in \mathcal{K}_{z,i}} E [|\psi(X_{t_i}) - \bar{z}(t_i, X_{t_i})|^2] \right) \\ & \quad + \frac{K_{q,i}}{M_i} + N \frac{K_{z,i}}{M_i} + \frac{K_{q,i} \log(M_i)}{M_i} + N \frac{K_{z,i} \log(M_i)}{M_i} \Big) + cN^{-1}, \end{aligned}$$

where $\mathcal{J} := \{i : t_i \in \bar{\pi}_\alpha\}$ and c is a positive constant not depending on N .

Theorem 4.2 provides, in the context of Theorem 4.1, the standard error bound of order $N^{-1/2}$ for the time discretization within the backward Euler scheme [2, 23]. Note that the projection errors are measured with respect to the continuous time PDE representation of the

¹ Available at: <https://www.uni-saarland.de/fakultaet-mi/stochastik/prof-dr-christian-bender/publications.html>

BSDE (1). Hence the choice of the basis functions can be adapted to known regularity results for the corresponding semilinear parabolic Cauchy problem, see, e.g., [4].

Under additional regularity assumptions, it is known from [8] that a time discretization error of order N^{-1} can be achieved. In the context of the SDP, the following result can be derived in this respect.

Theorem 4.3 *Additionally to the standing assumptions, suppose that $X = W$, and that f and \bar{y} are twice, resp. $s + 1$ times continuously differentiable ($s \geq 2$) with bounded derivatives. Then,*

$$\begin{aligned} & \max_{0 \leq i \leq N-1} E \left[|q_i^{N,M}(W_{t_i}) - \bar{y}(t_i, W_{t_i})|^2 \right] + \sum_{i=0}^{N-1} \Delta E \left[|z_i^{N,M}(W_{t_i}) - \bar{z}(t_i, W_{t_i})|^2 \right] \\ & \leq c \max_{i \in \mathcal{J}} \left(N^{1-\alpha} \inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\psi(W_{t_i}) - \bar{q}_i^N(W_{t_i})|^2 \right] + N^{2-2\alpha} \frac{K_{q,i}}{M_i} + N^{2-2\alpha} \frac{K_{q,i} \log(M_i)}{M_i} \right) \\ & \quad + c \max_{0 \leq i \leq N-1} \left(\inf_{\psi \in \mathcal{K}_{z,i}} E \left[|\psi(W_{t_i}) - \bar{q}_i^N(W_{t_i})|^2 \right] + \inf_{\psi \in \mathcal{K}_{z,i}} E \left[|\psi(W_{t_i}) - \bar{z}_i^N(W_{t_i})|^2 \right] \right. \\ & \quad \left. + \frac{K_{q,i}}{M_i} + N \frac{K_{z,i}}{M_i} + \frac{K_{q,i} \log(M_i)}{M_i} + N \frac{K_{z,i} \log(M_i)}{M_i} \right) + cN^{-2}, \end{aligned}$$

where $\mathcal{J} := \{i : t_i \in \bar{\pi}_\alpha\}$ and c is a positive constant not depending on N . Furthermore, \bar{q}^N and \bar{z}^N are bounded and $s + 1$ times, respectively, s times continuously differentiable with bounded derivatives.

Before we prove the theorems presented above in Section 6, we analyze how to optimally choose α and derive the resulting complexity of the algorithm. To this end, we first calibrate the algorithm in dependence of α to achieve a squared error of the order $N^{-2\theta}$ with $\theta \in \{1/2, 1\}$ with local polynomials as basis functions, cp. [11, 18].

For simplicity we use the same approximation space \mathcal{K}_z for the approximation of \bar{z}^N in each time step. Due to the additional projection error on the coarser time grid $\bar{\pi}_\alpha$ we distinguish between the time points inside and outside $\bar{\pi}_\alpha$ for the approximation of \bar{q}^N . We assume that the same approximation space $\bar{\mathcal{K}}_q$ is used for all time points $t_i \in \bar{\pi}_\alpha$ while we use a possibly different approximation space \mathcal{K}_q at the other time points. Analogously we assume that we use $M_i = M$ simulations at each time point t_i outside the coarser time grid $\bar{\pi}$ and $M_i = \bar{M}$ simulations otherwise. As already mentioned, we assume that one segment $(X_i, \dots, X_{\tau_\alpha(i)})$ of a path of X can be simulated at the cost of N^α . Under these assumptions, assuming we can evaluate the driver and the basis functions at cost 1, the performed simulations and evaluations during the algorithm lead to costs of order

$$NN^\alpha M + N^{1-\alpha} N^\alpha \bar{M}.$$

As basis functions we take local polynomials on cubes which we choose disjoint such that their union contains the set $\{x \in \mathbb{R}^D : |x| < C_b\}$ for a constant $C_b > 0$. We suppose that the edge length of the cubes is $\bar{\delta}_q$ for the approximation of \bar{q}^N on the time grid $\bar{\pi}_\alpha$, δ_q for the approximation of \bar{q}^N at all other time points and δ_z for the approximation of \bar{z}^N . Assuming that \bar{y} and \bar{z} are $s + 1$ times, respectively, s times continuously differentiable with bounded derivatives we set the degree of the polynomials as s for the approximation of \bar{q}^N and $s - 1$ for \bar{z}^N . We denote the set of polynomials of degree less than or equal to l by P_l . Then, the projection error in the context of Theorem 4.2 can be estimated by a Taylor expansion on each cube:

$$\begin{aligned}
& \inf_{\psi \in \mathcal{K}_q} E [|\psi(X_{t_i}) - \bar{y}(t_i, X_{t_i})|^2] \\
& \leq E \left[|\bar{y}(t_i, X_{t_i})|^2 \mathbb{1}_{|X_{t_i}| > C_b} \right] + \sum_{H \in \mathcal{H}_q} \inf_{\psi \in P_s} E [|\psi(X_{t_i}) - \bar{y}(t_i, X_{t_i})|^2 \mathbb{1}_{X_{t_i} \in H}] \\
& \leq \|\bar{y}(t_i, \cdot)\|_\infty^2 P(|X_{t_i}| > C_b) + c \|\bar{y}(t_i, \cdot)\|_\infty^{(s+1)} (\delta_q^{s+1})^2,
\end{aligned} \tag{3}$$

where \mathcal{H}_q denotes the collection of cubes applied for the approximation of \bar{q}^N at all time points outside $\bar{\pi}$. Under the assumption that $\sup_{0 \leq i \leq N} E[e^{\varpi |X_{t_i}|}] < \infty$ for some $\varpi > 0$, it follows by the Markov inequality that the choice $C_b = 2\theta\varpi^{-1} \log(N+1)$ ensures that the first term in (3) is of order $N^{-2\theta}$. The same bound holds for the second term when choosing the edge length of the hypercubes as $\delta_q = cN^{-\frac{\theta}{s+1}}$. Therefore it suffices to choose K_q of the order $N^{D\frac{\theta}{s+1}} \log^D(N+1)$ to ensure that

$$\inf_{\psi \in \mathcal{K}_q} E[|\psi(X_i) - \bar{y}(t_i, X_{t_i})|^2] \in O(N^{-2\theta}).$$

Following the same argumentation, we set

$$\bar{K}_q = cN^{D\frac{\theta + \frac{1-\alpha}{2}}{s+1}} \log^D(N+1), \quad K_z = cN^{D\frac{\theta}{s}} \log^D(N+1)$$

for a positive constant c to ensure

$$N^{1-\alpha} \inf_{\psi \in \bar{\mathcal{K}}_q} E [|\psi(X_i) - \bar{y}(t_i, X_{t_i})|^2] \in O(N^{-2\theta}), \quad \inf_{\psi \in \mathcal{K}_z} E [|\psi(X_i) - \bar{z}(t_i, X_{t_i})|^2] \in O(N^{-2\theta}),$$

where the change from $s+1$ to s in the number of basis functions in the approximation of \bar{z}^N occurs due to the lower smoothness of \bar{z}^N and \bar{K}_q and K_z denote the number of basis functions of the space $\bar{\mathcal{K}}_q$ and \mathcal{K}_z respectively. The same argument applies in the context of Theorem 4.3, replacing \bar{y} and \bar{z} by their discrete counterparts \bar{q}^N and \bar{z}^N .

Given the size of the approximation spaces we hence have to choose M of the order

$$N^{2\theta} \max\{K_q, NK_z\} = N^{1+2\theta} K_z$$

and \bar{M} of the order

$$N^{2\theta} \max\{N^{2-2\alpha} \bar{K}_q, NK_z\}$$

in order to bound the statistical error terms asymptotically by a multiple of $N^{-2\theta}$ (assuming that the driver f is not independent of Z and ignoring log-factors from now on). Then, in dependence of α , the computation costs of the algorithm grows as

$$C = \max\{N^{1+\alpha} M, N\bar{M}\} =: \max\{C_\pi, C_{\bar{\pi}}\}.$$

Here C_π is increasing in α and $C_{\bar{\pi}}$ is decreasing in α . The optimal choice of α is, thus, obtained by equating both terms, leading to

$$\alpha_{opt} = \frac{\frac{1}{2} - \frac{\theta}{s} + \frac{s+1}{D}}{\frac{1}{2} + \frac{3(s+1)}{D}}.$$

It always lies in the open interval $(0, 1)$, provided $\theta = \frac{1}{2}$, $s \geq 1$ or $\theta = 1$, $s \geq 2$. The resulting computational costs are of the order

$$C = NN^{\alpha_{opt}} M = N^{2+2\theta+D\frac{\theta}{s} + \frac{\frac{1}{2} - \frac{\theta}{s} + \frac{s+1}{D}}{\frac{1}{2} + \frac{3(s+1)}{D}}} = N^{3+2\theta+D\frac{\theta}{s} - \frac{\frac{\theta}{s} + \frac{2(s+1)}{D}}{\frac{1}{2} + \frac{3(s+1)}{D}}}.$$

In comparison, choosing $\alpha = 1$ (the MDP case), the computational costs are of the order

$$\mathcal{C} = cN^{3+2\theta+D\frac{\theta}{s}},$$

as already shown in [11]. Hence, the suggested SDP algorithm with the optimal choice of the segment length reduces the complexity by a factor of the order

$$N^{1-\alpha_{opt}} = N^{\frac{\theta/s+2(s+1)/D}{1/2+3(s+1)/D}}$$

depending on the smoothness s of the problems and the dimension D of the state space of X .

Remark 4.4 *Whenever some arbitrary function basis (increasing in the number N of time discretization steps) has been fixed and one wishes to control the squared simulation error by $N^{-2\theta}$ (up to log-factors), the previous reasoning can be adapted in a straightforward way to tune the segment length parameter α in order to minimize the algorithm's complexity.*

Remark 4.5 *Due to memory restrictions, it may be prohibitive to run the SDP algorithm with high order monomials as local basis functions in practice, in particular when the state space is high-dimensional. We therefore now fix a (small) degree s and apply local monomials up to degree s for the approximation of the Y -part and the Z -part of the solution (assuming that the problem is smooth enough, i.e., \bar{y} and \bar{z} are at least $s+1$ times continuously differentiable with bounded derivatives). A straightforward modification of the previous reasoning shows that (ignoring log-factors again) a number M^* of independent sample paths (segments) of the order*

$$N^{1+2\theta+D\frac{\theta}{s+1}}$$

must be simulated at each grid point of the fine grid, to achieve a squared error of the order $N^{-2\theta}$ for the Z -part. This number M^ corresponds to the memory requirement for the Z -part (the argumentation in [10] for the MDP scheme carries over to SDP without changes). Note that M^* is independent of the segment length parameter α . Choosing the number of independent sample paths at the grid points of the coarse grid of the same order as M^* and making sure that the squared error for the Y -part is also of the order $N^{-2\theta}$ leads to the condition that*

$$1 + 2\theta + D\frac{\theta}{s+1} = 2\theta + 2 - 2\alpha + D\frac{\theta + \frac{1-\alpha}{2}}{s+1},$$

i.e.,

$$\alpha_{mem} := \alpha = 1 - \left(2 + \frac{D}{2(s+1)}\right)^{-1}.$$

With this choice of α_{mem} , the SDP has the same memory requirement as the MDP and both schemes feature the same guaranteed rate of convergence of the order $N^{-\theta}$. The cost of the SDP scheme is, then, still smaller by a factor of $N^{1-\alpha_{mem}}$ compared to the MDP scheme. Noting that $\alpha_{mem} \in (1/2, 1)$ and that it approaches 1, if (for fixed s) the dimension D tends to infinity, we observe that applying the SDP is also beneficial (compared to the MDP) when the prime objective is to minimize the memory requirement, but the gain is more moderate compared to minimizing the complexity as objective. We stress that stratification proved useful for reducing the memory requirement in the context of the MDP scheme, see [10], and we expect that an analogous stratified variant of the SDP scheme could be analyzed similarly.

5. Numerical example

In this section, we compare the SDP and MDP approach in a numerical test case in order to illustrate our theoretical results.

For this purpose, we define for each $x \in \mathbb{R}^D$ and all $t \in [0, 0.2]$ the function

$$\varphi(t, x) := \exp\left(-\sum_{d=1}^D |x^{(d)} - t|^{0.3}\right) \prod_{d=1}^D (x^{(d)} - t)^2$$

and for $d = 1, \dots, D$

$$\phi_d(t, x) := \exp\left(-\sum_{e=1}^D |x^{(e)} - t|^{0.3}\right) (x^{(d)} - t) \left(2 - 0.3|x^{(d)} - t|^{0.3}\right) \prod_{e=1, e \neq d}^D (x^{(e)} - t)^2.$$

We then consider the BSDE driven by a Brownian motion $X = W$ with terminal time $T = 0.2$, the terminal condition

$$\xi(W_T) = \varphi(T, W_T)$$

and driver

$$f(t, x, y, z) := \min\{|z|, c\} - |\nabla\varphi(t, x)| - \left(\partial_t + \frac{1}{2}\Delta\right)\varphi(t, x).$$

for $c := \sup_{(t,x) \in [0,T] \times \mathbb{R}^D} |\nabla\varphi(t, x)|$, noting that φ is bounded and twice continuously differentiable with bounded derivatives. It can be easily checked by Itô's formula that the analytic solution to this BSDE is given by

$$Y_t = \varphi(t, W_t), \quad Z_t^{(d)} = \frac{\partial}{\partial x^{(d)}}\varphi(t, W_t) = \phi_d(t, W_t), \quad d = 1, \dots, D,$$

and that the standing assumptions are satisfied.

For a comparison between the MDP algorithm and the SDP algorithm, we calibrate the function basis and the number of sample paths as functions of the number of time steps N in line with the complexity analysis of the previous section with $\theta = 1/2$ and $s = 1$. We increase the number of time steps N and re-run each algorithm 40 times in dimension $D = 2$. Below we report the mean-squared errors and the average run times of both algorithms across the 40 repetitions. More precisely, we proceed in the following way.

Calibration:

We use piecewise linear functions for the approximation of \bar{q}_i^N and piecewise constant functions for the one of \bar{z}_i^N in both algorithms. At each time, we set the outer bound of the hypercubes as a multiple of the standard deviation of the Brownian motion at that time. More precisely, we choose $C_{b,i} := (2 \log(N) + 2)\sqrt{t_i}$ at the time t_i . As edge length of the cubes we choose $\delta_z = \sqrt{T}/N^{\frac{1}{2}}$ and $\delta_q = \sqrt{T}/N^{\frac{1}{4}}$ for the MDP algorithm leading to $K_{z,i} \in \mathcal{O}(\lceil R_i/\delta_z \rceil^D) = \mathcal{O}(N \log^2(N))$ and $K_{q,i} \in \mathcal{O}(\lceil R_i/\delta_q \rceil^D) = \mathcal{O}(N^{1/2} \log^2(N))$ at time t_i . We re-simulate the sample paths for the approximation of each conditional expectation and use $M_{q,i} = 10N K_{q,i} \in \mathcal{O}(N^{3/2} \log^2(N))$ simulations for Q at time t_i and $M_{z,i} = 5N^2 K_{z,i} \in \mathcal{O}(N^3 \log^2(N))$ simulations for Z .

We may choose δ_z , $K_{z,i}$ and $M_{z,i}$ for the SDP algorithm in the same way as in the MDP algorithm as well as δ_q and $K_{q,i}$ and $M_{q,i}$, if $t_i \notin \bar{\pi}$. The optimal value of the segment length parameter is $\alpha_{opt} = \frac{2}{7}$. For the time points in $\bar{\pi}$ we choose $\bar{\delta}_q = 1.5\sqrt{T}/N^{-\frac{1}{2} + \frac{\alpha}{4}}$ leading to $\bar{K}_{q,i} \in \mathcal{O}(N^{1-1/7} \log^2(N))$. As number of simulations for the approximation of Q at these time points we choose $\bar{M}_{q,i} = N^{3-2\alpha} \bar{K}_{q,i} \in \mathcal{O}(N^{3+2/7} \log^2(N))$.

Measuring the errors:

We measure the mean-squared error (MSE) of both algorithms for the approximation of \bar{q}_i^N

and \bar{z}_i^N separately and consider the average error over all the time steps. Additionally, we report the approximation error of Y at time zero, which is a relevant quantity in many applications such as nonlinear option pricing. Precisely, we consider the average over the 40 repetitions of the three indicators

$$\begin{aligned} \mathcal{C}_{y,av} &:= \frac{1}{N} \sum_{i=1}^{N-1} \sum_{H \in \mathcal{H}_{q,i}} |q_i^{M,N}(\Theta_H) - \bar{y}(t_i, \Theta_H)|^2 P(X_{t_i} \in H), \\ \mathcal{C}_{y,0} &:= |q_0^{M,N}(0) - \bar{y}(0, 0)|^2, \\ \mathcal{C}_{z,av} &:= \frac{1}{N} \sum_{i=0}^{N-1} \sum_{H \in \mathcal{H}_{z,i}} |z_i^{M,N}(\Theta_H) - \bar{z}(t_i, \Theta_H)|^2 P(X_{t_i} \in H), \end{aligned}$$

where Θ_H is the center of the cube H , and $\mathcal{H}_{q,i}$ and $\mathcal{H}_{z,i}$ are the sets of cubes used at time t_i for the approximation of \bar{q}_i^N and \bar{z}_i^N , respectively. We refer to the arithmetic mean of $\mathcal{C}_{y,av}$ and $\mathcal{C}_{z,av}$ over the 40 repetitions as average mean-squared error in time and analogously to the arithmetic mean of $\mathcal{C}_{y,0}$ over the 40 repetitions as mean-squared error at time 0.

Numerical results:

Figure 1 depicts \log_{10} - \log_{10} -plots of the mean-squared errors (as introduced above) and the average run times of both algorithms as the number N of time discretization points increases




Figure 1 Numerical results in dimension $D = 2$

from $N = 6$ to $N = 30$. In view of the results in Theorem 4.2 and the calibration of the algorithms we expect that the mean-squared errors decrease at the order N^{-1} . The numerical values of the three error indicators in Figures 1a–1c (marked by blue ‘x’) for the MDP scheme are largely in accordance with this expected convergence behavior. The numerical results for the SDP scheme (red circles) feature a convergence rate of N^{-1} for the average mean-squared error of the Z -component (Figure 1(b)) but a faster convergence rate of about N^{-2} for the mean-squared errors of the Y -components (Figures 1(a) and 1(c)). A possible explanation of this excellent empirical convergence rate are potential variance benefits when running empirical regressions on shorter time segments, see Remark 5.1. Figures 1(a) and 1(c) also illustrate a higher variance of the Y -approximations of the MDP scheme across the repetitions compared to the SDP scheme. Finally, the average run time over the repetitions are plotted against the number of time discretization points in Figure 1d. As before, the blue ‘x’-markers and the red circles display the numerical results for the MDP algorithm and the SDP algorithm, respectively. The lines correspond to the expected run time behavior of N^5 for the MDP scheme and $N^{4+2/7}$ for the SDP scheme derived from the complexity analysis at the end of Section 4. Taking into account that log-factors, which have been neglected throughout the complexity analysis, play a role for small values of N in both algorithms, Figure 1d confirms the theoretical run time benefits of the SDP scheme in the practical implementation.

Note that this example is only supposed to serve as a proof of concept of the improved complexity of the SDP scheme compared to the MDP scheme. Parallelization and variance reduction techniques can be incorporated analogously to the MDP case and are of prime importance when implementing regression on local polynomials for higher dimensional problems (of around $D = 10$), see, e.g., [10].

Remark 5.1 *For a possible explanation of the faster convergence of the Y -part in the SDP scheme, let us have a closer look at the simulation error: When estimating a regression function $m(x) = E[\Xi|X = x]$ via (truncated) linear least-squares regression, the squared statistical error can be bounded by*

$$\text{const.} \max\{\sigma^2, L\} \frac{K \log(M)}{M},$$

where σ^2 is an upper bound for the conditional variance $\text{Var}(\Xi|X)$, L is a bound for the regressand Ξ , K is the number of basis functions, and M is the sample size; see, e.g., Theorem 11.3 in [13]. The contribution involving the bound L of Ξ arises from changing from the empirical L^2 -norm to the L^2 -norm w.r.t. the law of X (cp., e.g., Lemma 6.5 below for a related result) and may be considered as a ‘worst case’ estimate. In each segment of the SDP scheme, one moves forward by a time step of the order $N^{-(1-\alpha)}$ at most. Hence, under sufficient regularity conditions, σ^2 is of the order $N^{-(1-\alpha)}$ for the Y -part. Neglecting the worst case contribution, one, thus, may hope for a faster convergence of the simulation error in some numerical test cases than the guaranteed rate stated in Theorem 4.1.

6. Error analysis

In this section, we present a complete and detailed error analysis of the SDP algorithm. In the first subsection, we introduce additional notation used in the proofs and present some key tools for the error analysis. The following subsections are then dedicated to the derivation of error bounds for the approximation of \bar{q}^N and \bar{z}^N respectively. We will establish a sort of recursion

formula for both parts, allowing us to bound the quadratic error at the time t_i by the one at time t_{i+1} plus an additional driver-dependent term. This illustrates the error propagation between the time steps. We will then derive global bounds for the quadratic error for both approximations before we analyze the driver-dependent terms appearing in both recursions further in Subsection 6.5. Finally, in the last subsection, we combine the obtained bounds to derive the result presented in Theorem 4.1.

6.1 Preliminaries and key tools

In this section, we discuss some ramifications for the error analysis by introducing additional objects and presenting key tools for the analysis.

First, recall the definition of the functions \bar{q}_i^N and \bar{z}_i^N . By setting

$$\begin{aligned} \Xi_{N-1}^N(\underline{x}_{N-1}) &:= \xi(x_N), \\ \Xi_i^N(\underline{x}_i) &:= \bar{q}_{\tau_\alpha(i)}^N(x_{\tau_\alpha(i)}) + \int_{t_{i+1}}^{t_{\tau_\alpha(i)+1}} f(s, x_s, \bar{y}(s, x_s), \bar{z}(s, x_s)) ds, \quad i \in \{0, \dots, N-2\}, \end{aligned}$$

for any $\underline{x}_i := (x_s)_{t_i \leq s \leq T} \in (\mathbb{R}^D)^{[t_i, T]}$, it holds

$$\bar{q}_i^N(x) = E [\Xi_i^N((X_s)_{t_i \leq s \leq T}) | X_{t_i} = x]$$

and

$$\bar{z}_i^N(x) = E \left[\frac{\Delta W_{i+1}}{\Delta} \Xi_i^N((X_s)_{t_i \leq s \leq T}) \middle| X_{t_i} = x \right].$$

Note that Ξ_i^N differs from $\Xi_i^{N,M}$ in two ways: in the functions $\Xi_i^{N,M}$, the true solution (Y, Z) of the BSDE is replaced by approximations of the algorithm and the integral is discretized. In order to make use of properties of the least squares projection, we need the analogs of the least squares solutions $\varphi^{q^{N,M}}$ and $\varphi^{z^{N,M}}$ based on the functions Ξ^N . Since those depend on the whole path of X rather than just the values on the time grid π additional fictitious simulations are required for the theoretical error analysis which motivates the following definition:

Definition 6.1 For $i \in \{0, \dots, N-1\}$, let $\mathcal{S}_i := \{\Delta W_{i+1}^{[i,m]}, X^{[i,m]} : m = 1, \dots, M\}$ be a cloud of independent random variables defined on the probability space $(\Omega^M, \mathcal{F}^M, P^M)$ with $X^{[i,m]} = (X_s^{[i,m]})_{t_i \leq s \leq T}$ such that $X^{[i,m]}$ is distributed like a segment of the SDE solution X . Furthermore, we assume that these simulations match the ones used in the SDP algorithm on the time grid π , i.e., $\Delta W_{i+1}^{[i,m]} = \Delta W_{i+1}^{[i,m,N]}$ and $X_{t_j}^{[i,m]} = X_{t_j}^{[i,m,N]}$ for all $i \in \{0, \dots, N-1\}$, $j \geq i$ and $m \in \{1, \dots, M_i\}$. Then, for every $\omega \in \Omega^M$, let $\nu_i^M(\omega, \cdot)$ be the measure on $((\mathbb{R}^D)^{[t_i, T]} \times \mathbb{R}^D, \mathcal{B}((\mathbb{R}^D)^{[t_i, T]} \times \mathbb{R}^D))$ defined by

$$\nu_i^M(\omega, B) := \frac{1}{M_i} \sum_{m=1}^{M_i} \delta_{(\Delta W_{i+1}^{[i,m]}(\omega), X^{[i,m]}(\omega))}(B),$$

where $\delta_c(\cdot)$ is the Dirac-measure on c .

The following calculations are done on the probability space $(\Omega^M, \mathcal{F}^M, P^M)$, where we suppose that there exists a D -dimensional Brownian motion W on $(\Omega^M, \mathcal{F}^M, P^M)$, which is independent of all simulations and hence also a copy X of the SDE solution independent of the simulations.

Given these additional random variables, we denote with $\varphi_i^{\bar{q}^N}$ and $\varphi_i^{\bar{z}^N}$ the solutions of the least-squares problems

$$\varphi_i^{\bar{q}^N} := \operatorname{argmin}_{\psi \in \mathcal{K}_{q,i}} \left(\frac{1}{M} \sum_{m=1}^M \left| \psi \left(X_{t_i}^{[i,m]} \right) - \Xi_i^N \left(X^{[i,m]} \right) \right|^2 \right)$$

and

$$\varphi_i^{\bar{z}^N} := \operatorname{argmin}_{\psi \in \mathcal{K}_{z,i}} \left(\frac{1}{M} \sum_{m=1}^{M_i} \left| \psi \left(X_{t_i}^{[i,m]} \right) - \frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \Xi_i^N \left(X^{[i,m]} \right) \right|^2 \right).$$

Remark 6.2 Recall the functions $\varphi_i^{q^{N,M}}$ and $\varphi_i^{z^{N,M}}$ from the SDP algorithm. Since it holds by definition of the ghost sample that $X_\pi^{[i,m]} := (X_{t_j}^{[i,m]})_{t_i < t_j \in \pi} = X^{[i,m,N]}$, we have

$$\varphi_i^{q^{N,M}} = \operatorname{argmin}_{\psi \in \mathcal{K}_{q,i}} \left(\frac{1}{M} \sum_{m=1}^M \left| \psi \left(X_{t_i}^{[i,m]} \right) - \Xi_i^{N,M} \left(X_\pi^{[i,m]} \right) \right|^2 \right)$$

and

$$\varphi_i^{z^{N,M}} = \operatorname{argmin}_{\psi \in \mathcal{K}_{z,i}} \left(\frac{1}{M} \sum_{m=1}^{M_i} \left| \psi \left(X_{t_i}^{[i,m]} \right) - \frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \Xi_i^{N,M} \left(X_\pi^{[i,m]} \right) \right|^2 \right),$$

i.e., for any fixed outcomes of the simulations $X^{[i,m]}$, both pairs of functions $\varphi_i^{q^{N,M}}$ and $\varphi_i^{z^{N,M}}$, and $\varphi_i^{\bar{q}^N}$ and $\varphi_i^{\bar{z}^N}$ solve a least squares problem with respect to the same measure $\nu_i^M(\omega, \cdot)$.

This allows us to utilize the following lemma, which is a key tool in the error analysis. It matches essentially Proposition 4.12 in [11] where the domain of the function Ξ is generalized in order to cover our setting. The proof presented in [11] still holds for this setting.

Lemma 6.3 For each $\omega \in \Omega^M$, let $(A, \mathcal{A}, \nu(\omega, \cdot))$ be a measurable space with

$$\nu(\omega, \cdot) = \frac{1}{M} \sum_{m=1}^M \delta_{\chi^{[m]}(\omega)}$$

for i.i.d random variables $\chi^{[1]} \dots, \chi^{[M]} : \Omega^M \rightarrow A$. Furthermore, let \mathcal{K} be a linear function space spanned by \mathbb{R}^l -valued basis functions $\{p^k(\cdot), 1 \leq k \leq K\}$ with $\sum_{k=1}^K E[|p^k(\chi^{[m]})|^2] < \infty$ for all m . For any $\mathcal{F}^M \otimes \mathcal{A}$ -measurable, \mathbb{R}^l -valued random variable Ξ with $\Xi(\omega, \cdot) \in L^2(\mathcal{A}, \nu(\omega, \cdot))$ for P^M -a.e. ω , set

$$\varphi(\omega, \cdot) := \operatorname{arginf}_{\psi \in \mathcal{K}} \frac{1}{M} \sum_{m=1}^M \left| \psi \left(\chi^{[m]}(\omega) \right) - \Xi \left(\omega, \chi^{[m]}(\omega) \right) \right|^2.$$

Then:

- (i) The mapping $\Xi \mapsto \varphi$ is linear.
- (ii) It holds

$$\|\varphi\|_{L^2(\mathcal{A}, \nu(\omega, \cdot))} \leq \|\Xi\|_{L^2(\mathcal{A}, \nu(\omega, \cdot))},$$

where we denote with $\|\cdot\|_{L^2(\mathcal{A}, \nu(\omega, \cdot))}$ the L^2 -norm with respect to the measure $\nu(\omega, \cdot)$.

(iii) Suppose \mathcal{G} is a sub- σ -field of \mathcal{F}^M such that $(p^k(\chi^{[1]}), \dots, p^k(\chi^{[M]}))$ is \mathcal{G} -measurable for each $k = 1, \dots, K$. Then

$$E[\varphi|\mathcal{G}](\omega, \cdot) = \operatorname{argmin}_{\psi \in \mathcal{K}} \frac{1}{M} \sum_{i=1}^M \left| \psi \left(\chi^{[i]}(\omega) \right) - \Xi_{\mathcal{G}} \left(\chi^{[i]}(\omega) \right) \right|^2,$$

where $\Xi_{\mathcal{G}}(x) := E[\Xi(x)|\mathcal{G}]$.

(iv) In the situation of (iii), suppose that \mathcal{G} is given by $\sigma(g(\chi^{[m]})_{m=1,\dots,M})$ for a \mathcal{A} -measurable function

$$g : A \rightarrow \mathbb{R}'.$$

Furthermore, assume that there is a sub- σ -field \mathcal{H} independent of $\sigma((\chi^{[m]})_{m=1,\dots,M})$ such that Ξ is $\mathcal{H} \otimes \mathcal{A}$ -measurable and that the conditional variance

$$E \left[\left| \Xi(\chi^{[m]}) - E[\Xi(\chi^{[m]})|\mathcal{G} \vee \mathcal{H}] \right|^2 \middle| \mathcal{G} \vee \mathcal{H} \right]$$

is P^M -almost surely uniformly bounded by some constant σ^2 for all $m \in \{1, \dots, M\}$. Then

$$E \left[\|\varphi - E[\varphi|\mathcal{G} \vee \mathcal{H}] \|_{L^2(\mathcal{A}, \nu(\omega, \cdot))}^2 \middle| \mathcal{G} \vee \mathcal{H} \right] \leq \sigma^2 \frac{K}{M}.$$

While Lemma 6.3 and the objects defined so far help us to utilize projection properties, we still need tools to handle the dependency on the different sets of simulations used in the algorithm. For this purpose, we first consider the following norms allowing us to distinguish between the dependence on simulations or the actual law of the true SDE solution X more clearly:

Definition 6.4 Let $\varphi : \Omega^M \times \mathbb{R}^D \rightarrow \mathbb{R}^l$ be $\mathcal{F}^M \times \mathcal{B}(\mathbb{R}^D)$ -measurable. For each $i = 0, \dots, N-1$, define the random norms $\|\cdot\|_{i,\infty}$ and $\|\cdot\|_{i,M}$ via

$$\|\varphi\|_{i,\infty}^2 := \int_{\mathbb{R}^D} |\varphi(x)|^2 P_{X_{t_i}}(dx), \quad \|\varphi\|_{i,M}^2 := \frac{1}{M} \sum_{m=1}^M \left| \varphi \left(X_{t_i}^{[i,m]} \right) \right|^2,$$

where $P_{X_{t_i}}$ denotes the distribution of X_{t_i} .

Note that we are interested in the error with respect to the law of the SDE solution X , i.e., in the difference between the approximations $q_i^{N,M}$ and $z_i^{N,M}$, and the functions \bar{q}_i^N and \bar{z}_i^N respectively in the $\|\cdot\|_{i,\infty}$ -norm. The following lemma allows us to lead this difference back to the one in the $\|\cdot\|_{i,M}$ -norm that appears naturally in the error analysis due to the use of simulations. It is a straightforward adaptation of Proposition 4.10 in [11] where the analogous result is shown for $\epsilon = 1$ instead of $\epsilon \in (0, 1]$.

Lemma 6.5 It holds for all $i = 0, \dots, N-1$ and any $\epsilon \in (0, 1]$ that

$$\begin{aligned} E \left[\|q_i^{N,M} - \bar{q}_i^N\|_{i,\infty}^2 \right] &\leq (1 + \epsilon) E \left[\|q_i^{N,M} - \bar{q}_i^N\|_{i,M}^2 \right] + \frac{C_1 K_{q,i} \log(C_2 M_i)}{M_i \epsilon}, \\ E \left[\|z_i^{N,M} - \bar{z}_i^N\|_{i,\infty}^2 \right] &\leq (1 + \epsilon) E \left[\|z_i^{N,M} - \bar{z}_i^N\|_{i,M}^2 \right] + \frac{\mathcal{D} C_1 K_{z,i} \log(C_2 M_i)}{\Delta M_i \epsilon} \end{aligned}$$

for positive constants C_1, C_2 independent of ϵ , Δ , and M_i .

Finally, the following lemma allows us to reduce the dependency on a sampled path of X to only the value of the sample at one time point t_i . A proof can be found in Chapter 5 of [15].

Lemma 6.6 Let \mathcal{G} and \mathcal{H} be independent sub- σ -fields of \mathcal{F}^M . For $l \geq 1$ let $F : \Omega^M \times \mathbb{R}^D \rightarrow \mathbb{R}^l$ be bounded and $\mathcal{G} \times \mathcal{B}(\mathbb{R}^D)$ -measurable and $U : \Omega^M \rightarrow \mathbb{R}^D$ be \mathcal{H} -measurable. Then $E[F(U)|\mathcal{H}] = j(U)$ where $j(x) = E[F(x)]$ for all $x \in \mathbb{R}^D$.

To see how we can utilize this lemma, note that each sample $X^{[i,m]}$ satisfies

$$X_{t_i+s}^{[i,m]} = X_{t_i}^{[i,m]} + \int_{t_i}^{t_i+s} b(l, X_l^{[i,m]}) dl + \int_{t_i}^{t_i+s} \sigma(l, X_l^{[i,m]}) dW_l^{[i,m]}$$

for a Brownian motion $W^{[i,m]}$. Substituting the time variable and setting $\tilde{W}_u = W_{u+t_i}^{[i,m]} - W_{t_i}^{[i,m]}$ leads to

$$X_{t_i+s}^{[i,m]} = X_{t_i}^{[i,m]} + \int_0^s b\left(u+t_i, X_{u+t_i}^{[i,m]}\right) du + \int_0^{t-s} \sigma\left(u+t_i, X_{u+t_i}^{[i,m]}\right) d\tilde{W}_u,$$

which shows that the sample $X^{[i,m]}$ is the solution to a forward SDE starting in t_i with initial value $X_{t_i}^{[i,m]}$ with respect to the Brownian motion \tilde{W}_u . Hence, we can express the path $X^{[i,m]}$ as a deterministic function h of $X_{t_i}^{[i,m]}$ and $(\tilde{W}_u)_{t_i \leq u \leq T}$ only, i.e., we can write $X_s^{[i,m]} = h(s, X_{t_i}^{[i,m]}, (\tilde{W}_u)_{0 \leq u \leq s-t_i})$ for any $s \geq t_i$. Since the Brownian motion \tilde{W} is independent of $\sigma(X_{t_i}^{[i,m]})$, we can then use Lemma 6.6 on the function

$$\begin{aligned} F(x_{t_i}) &= \int_{t_{i+1}}^{t_{\tau(i)+1}} f\left(s, h(s, x_{t_i}, (\tilde{W}_u)_{0 \leq u \leq s-t_i}), Y_s, Z_s\right) ds \\ &\quad - \Delta \sum_{j=i+1}^{\tau(i)} f\left(t_j, h(t_j, x_{t_i}, (\tilde{W}_u)_{0 \leq u \leq t_j-t_i}), q_j^{N,M}\left(h(t_j, x_{t_i}, (\tilde{W}_u)_{0 \leq u \leq t_j-t_i})\right), \right. \\ &\quad \left. z_j^{N,M}\left(h(t_j, x_{t_i}, (\tilde{W}_u)_{0 \leq u \leq t_j-t_i})\right)\right) \end{aligned}$$

and get

$$\begin{aligned} &E\left[\Xi_i^N\left(X^{[i,m]}\right) - \Xi_i^{N,M}\left(X^{[i,m]}\right) \middle| \sigma(\mathcal{S}_j : j > i) \vee \sigma\left(X_{t_i}^{[i,m]}\right)\right] \\ &= E\left[\Xi_i^N(\underline{X}_{t_i}) - \Xi_i^{N,M}(\underline{X}_{t_i}) \middle| \sigma(\mathcal{S}_j : j > i), X_{t_i} = X_{t_i}^{[i,m]}\right], \end{aligned} \quad (4)$$

where we set $\underline{X}_{t_i} = (X_s)_{s \in [t_i, T]}$. Completely analogously, it holds

$$\begin{aligned} &E\left[\frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \left(\Xi_i^N\left(X^{[i,m]}\right) - \Xi_i^{N,M}\left(X^{[i,m]}\right)\right) \middle| \sigma(\mathcal{S}_j : j > i) \vee \sigma\left(X_{t_i}^{[i,m]}\right)\right] \\ &= E\left[\frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \left(\Xi_i^N(\underline{X}_{t_i}) - \Xi_i^{N,M}(\underline{X}_{t_i})\right) \middle| \sigma(\mathcal{S}_j : j > i), X_{t_i} = X_{t_i}^{[i,m]}\right]. \end{aligned} \quad (5)$$

Hence, Lemma 6.6 allows us to reduce the dependency on a sample path to a conditional expectation according to the actual law of X given the value of the sample path at the current time.

We close this section with some abbreviations for the notation: Throughout the rest of this chapter, we denote with $\mathcal{F}_i^M := \sigma(\mathcal{S}_k : k > i) \vee \sigma(X_{t_i}^{[i,m]} : m = 1, \dots, M_i)$ the σ -field generated by the simulations used up to the time k (backwards starting from N) for a fixed $N \in \mathbb{N}$. With $\mathcal{F}_i := \sigma((W_s)_{0 \leq s \leq t_i})$ we denote the σ -field generated by the Brownian motion which is independent of the simulations. The conditional expectations given those σ -fields we denote with $E_i^M[\cdot] = E[\cdot | \mathcal{F}_i^M]$ and $E_i[\cdot] := E[\cdot | \mathcal{F}_i]$. Additionally, we shorten the notation of the driver f by dropping clearly indicated function arguments through the notation $f(t, x, g, g') := f(t, x, g(x), g'(x))$ for any functions $g, g' : \mathbb{R}^D \rightarrow \mathbb{R}^l$ and $f(t, x, g_t, g'_t) := f(t, x, (g(t, x), g'(t, x)))$ for any functions $g, g' : [0, T] \times \mathbb{R}^D \rightarrow \mathbb{R}^l$.

After these considerations we are ready for the derivation of the error bounds.

6.2 Error of the Q approximation

In this section, we analyze the expected quadratic error of the approximation of \bar{q}^N in the form of the terms

$$E \left[\|q_i^{N,M} - \bar{q}_i^N\|_{i,M}^2 \right].$$

We first establish a bound on the error propagation between the time steps, allowing us to express the expected difference of this term by the one in the next time step. Afterwards, we derive a local error bound for the term as well as a bound for the global error by bounding the maximum of the quadratic error terms over all time points $t_i \in \pi$. The first part of the proof, in which we establish the error propagation, is inspired by the error analysis of the MDP scheme in [11].

Error propagation: First, note that

$$\begin{aligned} E_i^M \left[\Xi_i^N \left(X^{[i,m]} \right) \right] &= E_i^M \left[\bar{q}_{\tau_\alpha(i)}^N \left(X_{t_{\tau_\alpha(i)}}^{[i,m]} \right) + \int_{t_{i+1}}^{t_{\tau_\alpha(i)+1}} f \left(s, X_s^{[i,m]}, \bar{y}_s, \bar{z}_s \right) ds \right] \\ &= E \left[\bar{q}_{\tau_\alpha(i)}^N \left(X_{t_{\tau_\alpha(i)}} \right) + \int_{t_{i+1}}^{t_{\tau_\alpha(i)+1}} f \left(s, X_s, Y_s, Z_s \right) ds \middle| X_{t_i} = X_{t_i}^{[i,m]} \right] \\ &= \bar{q}_i^N \left(X_{t_i}^{[i,m]} \right). \end{aligned}$$

Hence, it holds by Lemma 6.3 (iii) that $E_i^M[\varphi_i^{\bar{q}^N}]$ is the least squares projection of \bar{q}_i^N on the subspace $\mathcal{K}_{q,i}$ with respect to the measure ν_i^M , i.e., it holds

$$E_i^M \left[\varphi_i^{\bar{q}^N} \right] = \operatorname{arginf}_{\psi \in \mathcal{K}_{q,i}} \left(\frac{1}{M} \sum_{i=1}^M \left| \psi \left(X_{t_i}^{[i,m]} \right) - \bar{q}_i^N \left(X_{t_i}^{[i,m]} \right) \right|^2 \right).$$

Therefore, by the properties of least square projections, $\bar{q}_i^N - E_i^M[\varphi_i^{\bar{q}^N}]$ is orthogonal on $E_i^M[\varphi_i^{\bar{q}^N}] - \psi_i^{q^{N,M}}$ with respect to ν_i^M . Additionally, note that, since ξ and f are bounded due to the assumptions (A_ξ) and (A_f) , it follows by a simple backward recursion that

$$|\bar{q}_i^N(x)| \leq C_{q,i} = C_\xi + (T - t_{i+1})C_f \leq C_q := C_\xi + TC_f$$

for all $x \in \mathbb{R}^D$. We conclude that $\mathcal{T}_{C_{q,i}}(\bar{q}_i^N(x)) = \bar{q}_i^N(x)$ and obtain:

$$\begin{aligned} E \left[\|\bar{q}_i^N(\cdot) - q_i^{N,M}(\cdot)\|_{i,M}^2 \right] &= E \left[\left\| \mathcal{T}_{C_{q,i}}(\bar{q}_i^N(\cdot)) - \mathcal{T}_{C_{q,i}}(\varphi_i^{q^{N,M}}(\cdot)) \right\|_{i,M}^2 \right] \\ &\leq E \left[\left\| \bar{q}_i^N(\cdot) - E_i^M[\varphi_i^{\bar{q}^N}(\cdot)] + E_i^M[\varphi_i^{\bar{q}^N}(\cdot)] - \varphi_i^{q^{N,M}}(\cdot) \right\|_{i,M}^2 \right] \\ &= E \left[\left\| \bar{q}_i^N(\cdot) - E_i^M[\varphi_i^{\bar{q}^N}(\cdot)] \right\|_{i,M}^2 \right] + E \left[\left\| E_i^M[\varphi_i^{\bar{q}^N}(\cdot)] - \varphi_i^{q^{N,M}}(\cdot) \right\|_{i,M}^2 \right] \\ &\leq E \left[\left\| \bar{q}_i^N(\cdot) - E_i^M[\varphi_i^{\bar{q}^N}(\cdot)] \right\|_{i,M}^2 \right] \\ &\quad + (1 + \kappa) E \left[\left\| E_i^M[\varphi_i^{q^{N,M}}(\cdot)] - \varphi_i^{\bar{q}^N}(\cdot) \right\|_{i,M}^2 \right] \\ &\quad + (1 + \kappa^{-1}) E \left[\left\| E_i^M[\varphi_i^{q^{N,M}}(\cdot)] - \varphi_i^{q^{N,M}}(\cdot) \right\|_{i,M}^2 \right], \end{aligned}$$

where κ is an arbitrary positive constant. We handle the terms separately:

As argued before, it holds

$$E_i^M \left[\varphi_i^{\bar{q}^N}(\cdot) \right] = \operatorname{arginf}_{\psi \in \mathcal{K}_{q,i}} \|\bar{q}_i^N(\cdot) - \psi(\cdot)\|_{i,M}^2.$$

Hence,

$$\begin{aligned}
E \left[\left\| \bar{q}_i^N(\cdot) - E_i^M \left[\varphi_i^{\bar{q}^N}(\cdot) \right] \right\|_{i,M}^2 \right] &= E \left[\inf_{\psi \in \mathcal{K}_{q,i}} \left\| \bar{q}_i^N(\cdot) - \psi(\cdot) \right\|_{i,M}^2 \right] \\
&\leq \inf_{\psi \in \mathcal{K}_{q,i}} \frac{1}{M} \sum_{i=1}^M E \left[\left| \bar{q}_i^N(X_{t_i}^{[i,m]}) - \psi(X_{t_i}^{[i,m]}) \right|^2 \right] \\
&= \inf_{\psi \in \mathcal{K}_{q,i}} E \left[\left| \bar{q}_i^N(X_{t_i}) - \psi(X_{t_i}) \right|^2 \right].
\end{aligned}$$

This term describes the best approximation error due to the projection on the subspace $\mathcal{K}_{q,i}$ and is part of the final error representation.

For the next term, note that $\Xi_i^{N,M}$ is bounded by $C_{q,i}$ under assumptions (A_ξ) and (A_f) , since the approximations $q_i^{N,M}$ are (due to the truncation in the algorithm). Additionally, for each i , the function $\Xi_i^{N,M}$ is built using only the simulations in the sets \mathcal{S}_k for $k > i$. Hence, it follows directly by Lemma 6.3 (iv) with $\mathcal{H} = \sigma(\mathcal{S}_k, k > i)$ and $g(X^{[i,m]}) = X_{t_i}^{[i,m]}$ that

$$E \left[\left\| E_i^M \left[\varphi_i^{q_i^{N,M}}(\cdot) \right] - \varphi_i^{q_i^{N,M}}(\cdot) \right\|_{i,M}^2 \right] \leq C_{q,i}^2 \frac{K_{q,i}}{M_i}.$$

Since, as argued before, the functions \bar{q}_i^N are bounded by $C_{q,i}$ as well, we can apply a similar argument to the functions Ξ_i^N . Those are again built using only the simulations in \mathcal{S}_k for $k > i$. Setting $\xi_i^q(x) := E[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) | X_{t_i} = x, \mathcal{F}_0^M]$, we have by (4) that

$$E_i^M \left[\Xi_i^{N,M}(X^{[i,m]}) - \Xi_i^N(X^{[i,m]}) \right] = \xi_i^q(X_i^{[i,m]}).$$

Then Lemma 6.3 (i) and (iii) imply that

$$E_i^M \left[\varphi_i^{q_i^{N,M}}(\cdot) - \varphi_i^{\bar{q}_i^N}(\cdot) \right] = \operatorname{arginf}_{\psi \in \mathcal{K}_{q,i}} \left(\frac{1}{M} \sum_{i=1}^M \left| \psi(X_{t_i}^{[i,m]}) - \xi_i^q(X_{t_i}^{[i,m]}) \right|^2 \right).$$

Hence, by Lemma 6.3 (ii) it holds

$$E \left[\left\| E_i^M \left[\varphi_i^{q_i^{N,M}}(\cdot) - \varphi_i^{\bar{q}_i^N}(\cdot) \right] \right\|_{i,M}^2 \right] \leq E \left[\left\| \xi_i^q(\cdot) \right\|_{i,M}^2 \right] = E \left[\xi_i^q(X_{t_i})^2 \right].$$

Plugging in the estimates derived so far we have:

$$\begin{aligned}
E \left[\left\| \bar{q}_i^N(\cdot) - q_i^{N,M}(\cdot) \right\|_{i,M}^2 \right] &\leq \inf_{\psi \in \mathcal{K}_{q,i}} E \left[\left(\bar{q}_i^N(X_{t_i}) - \psi(X_{t_i}) \right)^2 \right] \\
&\quad + (1 + \kappa^{-1}) C_{q,i}^2 \frac{K_{q,i}}{M_i} + (1 + \kappa) E \left[\xi_i^q(X_{t_i})^2 \right]. \tag{6}
\end{aligned}$$

To further estimate $E[\xi_i^q(X_{t_i})^2]$, we have to distinguish between the time points. If the time points t_i and t_{i+1} are in the same segment defined by $\bar{\pi}_\alpha$, i.e., at time points t_i such that $t_{i+1} \notin \bar{\pi}_\alpha$, it holds $\tau_\alpha(i) = \tau_\alpha(i+1)$ by our choice of τ_α and hence:

$$\begin{aligned}
\Xi_i^{N,M}(\underline{X}_{t_i}) &= q_{\tau_\alpha(i)}^{N,M}(X_{t_{\tau_\alpha(i)}}) + \sum_{j=i+1}^{\tau_\alpha(i)} \Delta f \left(t_j, X_{t_j}, q_j^{N,M}, z_j^{N,M} \right) \\
&= \Xi_{i+1}^{N,M}(\underline{X}_{t_{i+1}}) + \Delta f \left(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M} \right)
\end{aligned}$$

and

$$\begin{aligned}
\Xi_i^N(\underline{X}_{t_i}) &= \bar{q}_{\tau_\alpha(i)}^N(X_{t_{\tau_\alpha(i)}}) + \int_{t_{i+1}}^{t_{\tau_\alpha(i)+1}} f(t, X_t, Y_t, Z_t) dt \\
&= \Xi_{i+1}^N(\underline{X}_{t_{i+1}}) + \int_{t_{i+1}}^{t_{i+2}} f(t, X_t, Y_t, Z_t) dt.
\end{aligned}$$

This allows us to estimate $E[\xi_i^q(X_{t_i})^2]$ as:

$$\begin{aligned}
& E \left[\xi_i^q(X_{t_i})^2 \right] \\
&= E \left[E \left[\bar{q}_{\tau_\alpha(i)}^N(X_{t_{\tau_\alpha(i)}}) - q_{\tau_\alpha(i)}^{N,M}(X_{t_{\tau_\alpha(i)}}) \right. \right. \\
&\quad \left. \left. + \sum_{j=i+1}^{\tau_\alpha(i)} \int_{t_j}^{t_{j+1}} f(s, X_s, Y_s, Z_s) - f(t_j, X_{t_j}, q_j^{N,M}, z_j^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
&\leq (1 + \Gamma\Delta) E \left[E \left[\Xi_{i+1}^N(\underline{X}_{t_{i+1}}) - \Xi_{i+1}^{N,M}(\underline{X}_{t_{i+1}}) \middle| \mathcal{F}_0^M, X_{t_i}, X_{t_{i+1}} \right] \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \\
&\quad + \left(1 + \frac{1}{\Gamma\Delta} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
&\leq (1 + \Gamma\Delta) E \left[(\xi_{i+1}^q(X_{t_{i+1}}))^2 \right] \\
&\quad + \left(1 + \frac{1}{\Gamma\Delta} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right].
\end{aligned}$$

Here we first used Young's inequality with some constant $\Gamma > 0$ that will be specified later and the tower property of the conditional expectation in the first inequality. Then in the second step, we used Jensen's inequality, the Markov property of X and once more the tower property of the conditional expectation. The calculation shows that the expected error term $E[\xi_i^q(X_i)^2]$ is bounded by the one in the next time step and a driver-dependent term.

At time points at the end of a segment, i.e., for time points t_i such that $t_{i+1} \in \bar{\pi}_\alpha$, the function $\Xi_i^{N,M}$ and $\Xi_{i+1}^{N,M}$ are defined on different segments and hence the first inequality in the calculations above does not hold true in this case. However, to get a similar bound, we can once more use Young's inequality and a zero addition to get

$$\begin{aligned}
& E \left[(\xi_i^q(X_{t_i}))^2 \right] \\
&= E \left[E \left[\bar{q}_{\tau_\alpha(i)}^N(X_{t_{\tau_\alpha(i)}}) - q_{\tau_\alpha(i)}^{N,M}(X_{t_{\tau_\alpha(i)}}) \right. \right. \\
&\quad \left. \left. + \int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
&\leq (1 + \Gamma\Delta) E \left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] \\
&\quad + \left(1 + \frac{1}{\Gamma\Delta} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
&\leq (1 + \Gamma\Delta)(1 + \kappa)(1 + \epsilon) E \left[(\xi_{i+1}^q(X_{t_{i+1}}))^2 \right] \\
&\quad + \left(1 + \frac{1}{\Gamma\Delta} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
&\quad + (1 + \Gamma\Delta) \left(E \left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] - (1 + \kappa)(1 + \epsilon) E \left[(\xi_{i+1}^q(X_{t_{i+1}}))^2 \right] \right)_+,
\end{aligned}$$

with positive constants κ and ϵ , which we will specify later. Again we have bounded the error term $E[\xi_i^q(X_{t_i})^2]$ by the one at the next time point and a driver-dependent term, but now with an additional error term that depends on the approximation of \bar{q}^N at the time point t_{i+1} . This additional error term could be expected since the SDP algorithm works on the segment containing t_i like the MDP scheme where the correct terminal condition of the BSDE restricted to the corresponding time segment has been replaced by the approximation $q_{\tau_\alpha(i)}^N$ at the time

point $\tau_\alpha(i) = t_{i+1}$.

Local and global bounds: Iterating this step leads to the following local and global bounds for the quadratic error of the approximation of \bar{q}^N , that are stated in terms of the expectations $E[\xi_i^q(X_i)^2]$ for later use. To obtain a corresponding bound for the terms $E[\|\bar{q}_i^N - q_i^{N,M}\|_{i,M}^2]$, one can simply follow the arguments that lead to (6) and apply the lemma afterwards.

Lemma 6.7 *For a positive constant Γ , set $\lambda_i := (1 + \Gamma\Delta)^i(1 + N^{\alpha-1})^{2|\{j \leq i: t_j \in \bar{\pi}_\alpha\}|}$ for $i \in \{0, \dots, N\}$. It then holds under the standing assumptions that*

$$\begin{aligned} E\left[(\xi_i^q(X_{t_i}))^2\right] &\leq \lambda_i E\left[(\xi_i^q(X_{t_i}))^2\right] \\ &\leq \left(1 + \frac{1}{\Delta\Gamma}\right) \sum_{j=i}^{N-2} \lambda_j E\left[E\left[\int_{t_{j+1}}^{t_{j+2}} f(s, X_s, Y_s, Z_s) - f\left(t_{j+1}, X_{t_{j+1}}, q_{j+1}^{N,M}, z_{j+1}^{N,M}\right) ds \middle| \mathcal{F}_0^M, X_{t_j}\right]^2\right] \\ &\quad + [N^{1-\alpha}] \lambda_N \sup_{j \in \mathcal{J}} \left(E\left[\|\bar{q}_j^N - q_j^{N,M}\|_{j,\infty}^2\right] - (1 + N^{\alpha-1})^2 E\left[(\xi_j^q(X_{t_j}))^2\right]\right)_+ \end{aligned}$$

for all $i \in \{0, \dots, N-2\}$ and

$$\begin{aligned} &\max_{0 \leq i \leq N-1} E\left[(\xi_i^q(X_{t_i}))^2\right] \\ &\leq \sum_{j=0}^{N-2} \left(1 + \frac{1}{\Delta\Gamma}\right) \lambda_j E\left[E\left[\int_{t_{j+1}}^{t_{j+2}} f(s, X_s, Y_s, Z_s) - f\left(t_{j+1}, X_{t_{j+1}}, q_{j+1}^{N,M}, z_{j+1}^{N,M}\right) ds \middle| \mathcal{F}_0^M, X_{t_j}\right]^2\right] \\ &\quad + [N^{1-\alpha}] \lambda_N \max_{j \in \mathcal{J}} \left((1 + N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,j}} E\left[|\bar{q}_j^N(X_{t_j}) - \psi(X_{t_j})|^2\right] + (1 + N^{1-\alpha}) \frac{C_{q,j}^2 K_{q,j}}{M_j} \right) \right. \\ &\quad \left. + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right). \end{aligned}$$

where $\mathcal{J} := \{i : t_i \in \bar{\pi}_\alpha\}$.

Proof Iterating the previous calculations where we choose $\kappa = \epsilon = N^{\alpha-1}$ yields

$$\begin{aligned} E\left[(\xi_i^q(X_{t_i}))^2\right] &\leq \lambda_i E\left[(\xi_i^q(X_{t_i}))^2\right] \\ &\leq \lambda_{i+1} E\left[(\xi_{i+1}^q(X_{t_{i+1}}))^2\right] \\ &\quad + \lambda_i \left(1 + \frac{1}{\Delta\Gamma}\right) E\left[E\left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f\left(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}\right) ds \middle| \mathcal{F}_0^M, X_{t_i}\right]^2\right] \\ &\quad + \lambda_{i+1} \left(E\left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1,\infty}^2\right] - (1 + N^{\alpha-1})^2 E\left[(\xi_{i+1}^q(X_{t_{i+1}}))^2\right]\right)_+ \mathbf{1}_{\mathcal{J}}(i+1) \\ &\leq \lambda_{N-1} E\left[(\xi_{N-1}^q(X_{t_{N-1}}))^2\right] \\ &\quad + \left(1 + \frac{1}{\Delta\Gamma}\right) \sum_{j=i}^{N-2} \lambda_j E\left[E\left[\int_{t_{j+1}}^{t_{j+2}} f(s, X_s, Y_s, Z_s) - f\left(t_{j+1}, X_{t_{j+1}}, q_{j+1}^{N,M}, z_{j+1}^{N,M}\right) ds \middle| \mathcal{F}_0^M, X_{t_j}\right]^2\right] \\ &\quad + [N^{1-\alpha}] \lambda_N \max_{j \in \mathcal{J}} \left(E\left[\|\bar{q}_j^N - q_j^{N,M}\|_{j,\infty}^2\right] - (1 + N^{\alpha-1})^2 E\left[(\xi_j^q(X_{t_j}))^2\right]\right)_+. \end{aligned}$$

Then we have by definition $E[\xi_{N-1}^q(X_{t_{N-1}})^2] = 0$ since $\Xi_{N-1}^{N,M} = \Xi_{N-1}^N$ and the recursion terminates. This already proves the first statement of Lemma 6.7. Additionally, using Lemma 6.5 with $\epsilon = N^{\alpha-1}$ and the inequality (6) for $\kappa = N^{\alpha-1}$ we have for any j :

$$\begin{aligned}
& \left(E \left[\|\bar{q}_j^N - q_j^{N,M}\|_{j,\infty}^2 \right] - (1 + N^{\alpha-1})^2 E \left[(\xi_j^q(X_{t_j}))^2 \right] \right)_+ \\
& \leq \left((1 + N^{\alpha-1}) E \left[\|\bar{q}_j^N - q_j^{N,M}\|_{j,M}^2 \right] + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right. \\
& \quad \left. - (1 + N^{\alpha-1})^2 E \left[(\xi_j^q(X_{t_j}))^2 \right] \right)_+ \\
& \leq \left((1 + N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,j}} E \left[|\bar{q}_j^N(X_{t_j}) - \psi(X_{t_j})|^2 \right] + (1 + N^{1-\alpha}) \frac{C_{q,j}^2 K_{q,j}}{M_j} \right. \right. \\
& \quad \left. \left. + (1 + N^{\alpha-1}) E \left[(\xi_j^q(X_{t_j}))^2 \right] \right) + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{\epsilon M_j} - (1 + N^{\alpha-1})^2 E \left[(\xi_j^q(X_{t_j}))^2 \right] \right)_+ \\
& \leq (1 + N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,j}} E \left[|\bar{q}_j^N(X_{t_j}) - \psi(X_{t_j})|^2 \right] + (1 + N^{1-\alpha}) \frac{C_{q,j}^2 K_{q,j}}{M_j} \right) + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j}.
\end{aligned} \tag{7}$$

The second statement of Lemma 6.7 then follows by plugging in the estimate above in the first statement and taking the maximum over all time points. \square

6.3 Error of the Z approximation

Analogously to the previous section, we now analyze the quadratic error of the approximation of \bar{z}^N via the terms $E[\|\bar{z}_i^N - z_i^{N,M}\|_{i,M}^2]$. Again, we first establish a bound on the error propagation between the time steps before deriving global bounds.

Error propagation: While the later steps require changes, we can get an analog of the inequality (6) by applying the same arguments as before. It holds that

$$E_i^M \left[\frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \Xi_i^N \left(X_{t_i}^{[i,m]} \right) \right] = \bar{z}_i^N \left(X_{t_i}^{[i,m]} \right)$$

and therefore, we have by Lemma 6.3 (iii)

$$E_i^M \left[\varphi_i^{\bar{z}^N} \right] = \operatorname{arginf}_{\psi \in \mathcal{K}_{z,i}} \left(\frac{1}{M} \sum_{i=1}^M \left| \psi \left(X_{t_i}^{[i,m]} \right) - \frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \bar{z}_i^N \left(X_{t_i}^{[i,m]} \right) \right|^2 \right). \tag{8}$$

We conclude that $\bar{z}_i^N - E_i^M[\varphi_i^{\bar{z}^N}]$ is orthogonal on $E_i^M[\varphi_i^{\bar{z}^N}] - \varphi_i^{\bar{z}^N}$ with respect to $\|\cdot\|_{i,M}$. Additionally, since $|\Xi_i^N| \leq C_{q,i}$ due to assumptions (A_ξ) and (A_f) , it holds

$$|\bar{z}_i^{N,(d)}(x)| \leq C_{z,i} = \frac{C_{q,i}}{\sqrt{\Delta}}$$

for each component $\bar{z}_i^{N,(d)}$, $d = 1, \dots, \mathcal{D}$ of \bar{z}_i^N , $x \in \mathbb{R}^{\mathcal{D}}$ and $i \in \{0, \dots, N-1\}$. With that, we obtain for an arbitrary $\kappa > 0$ that

$$\begin{aligned}
\Delta E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i,M}^2 \right] &= \Delta E \left[\|\mathcal{T}_{C_{z,i}}(\bar{z}_i^N(\cdot)) - \mathcal{T}_{C_{z,i}}(\varphi_i^{z^{N,M}}(\cdot))\|_{i,M}^2 \right] \\
&\leq \Delta E \left[\|\bar{z}_i^N(\cdot) - E_i^M[\varphi_i^{\bar{z}^N}(\cdot)] + E_i^M[\varphi_i^{\bar{z}^N}(\cdot)] - \varphi_i^{z^{N,M}}(\cdot)\|_{i,M}^2 \right] \\
&= \Delta \left(E \left[\|\bar{z}_i^N(\cdot) - E_i^M[\varphi_i^{\bar{z}^N}(\cdot)]\|_{i,M}^2 \right] + E \left[\|E_i^M[\varphi_i^{\bar{z}^N}(\cdot)] - \varphi_i^{z^{N,M}}(\cdot)\|_{i,M}^2 \right] \right) \\
&\leq \Delta \left(E \left[\|\bar{z}_i^N(\cdot) - E_i^M[\varphi_i^{\bar{z}^N}(\cdot)]\|_{i,M}^2 \right] \right. \\
&\quad \left. + (1 + \kappa^{-1}) + E \left[\|E_i^M[\varphi_i^{z^{N,M}}(\cdot)] - \varphi_i^{z^{N,M}}(\cdot)\|_{i,M}^2 \right] \right. \\
&\quad \left. + (1 + \kappa) E \left[\|E_i^M[\varphi_i^{z^{N,M}}(\cdot)] - \varphi_i^{\bar{z}^N}(\cdot)\|_{i,M}^2 \right] \right).
\end{aligned}$$

Once more we handle the appearing terms separately:

First, analogously to the corresponding term in the previous section, it follows due to equation (8) that

$$E[\|\bar{z}_i^N(\cdot) - E_i^M[\varphi_i^{\bar{z}^N}(\cdot)]\|_{i,M}^2] \leq \inf_{\psi \in \mathcal{K}_{z,i}} E[|\psi(X_{t_i}) - \bar{z}_i^N(X_{t_i})|^2],$$

which describes the best approximation error of \bar{z}^N using the basis functions and is part of the final error representation.

For the next term, note again that $\Xi_i^{N,M}$ is bounded by $C_{q,i}$ for all $i \in \{0, \dots, N-1\}$. We conclude that

$$\begin{aligned}
&E_i^M \left[\left| \frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \Xi_i^{N,M}(X^{[i,m]}) - E_i^M \left[\frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \Xi_i^{N,M}(X^{[i,m]}) \right] \right|^2 \right] \\
&\leq E_i^M \left[\left| \frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \Xi_i^{N,M}(X^{[i,m]}) \right|^2 \right] \\
&\leq \frac{\mathcal{D}C_{q,i}^2}{\Delta}.
\end{aligned}$$

Then, since $\Xi_i^{N,M}$ is built using only simulations of the clouds \mathcal{S}_k for $k > i$, it follows by Lemma 6.3 (iv) that $E[\|E_i^M[\varphi_i^{z^{N,M}}] - \varphi_i^{z^{N,M}}\|_{i,M}^2]$ is bounded by $C_{q,i}^2 \frac{\mathcal{D}K_{z,i}}{\Delta M_i}$.

For the last term, we have by (5) that

$$E_i^M \left[\frac{\Delta W_{i+1}^{[i,m]}}{\Delta} \left(\Xi_i^{N,M}(X^{[i,m]}) - \Xi_i^N(X^{[i,m]}) \right) \right] = \xi_i^z(X_{t_i}^{[i,m]})$$

with

$$\xi_i^z(x) = E \left[\frac{\Delta W_{i+1}}{\Delta} \left(\Xi_i^N(\underline{X}_{t_i}) - \Xi_i^{N,M}(\underline{X}_{t_i}) \right) \middle| X_{t_i} = x, \mathcal{F}_0^M \right].$$

Then, by Lemma 6.3 (i) and (iii), it follows that

$$E_i^M \left[\varphi_i^{z^{N,M}}(\cdot) - \varphi_i^{\bar{z}^N}(\cdot) \right] = \operatorname{arginf}_{\psi \in \mathcal{K}_{z,i}} \left(\frac{1}{M} \sum_{i=1}^M \left| \psi(X_{t_i}^{[i,m]}) - \xi_i^z(X_{t_i}^{[i,m]}) \right|^2 \right).$$

Hence, by Lemma 6.3 (ii) we have

$$E \left[\left\| E_i^M \left[\psi_i^{z_i^{N,M}} - \psi_i^{\bar{z}_i^N} \right] \right\|_{i,M}^2 \right] \leq E \left[\left\| \xi_i^z \right\|_{i,M}^2 \right] = E \left[\left(\xi_i^z(X_{t_i}) \right)^2 \right].$$

Plugging in the estimates obtained so far we have

$$\begin{aligned} \Delta E \left[\left\| \bar{z}_i^N - z_i^{N,M} \right\|_{i,M}^2 \right] &\leq \Delta \left(\inf_{\psi \in \mathcal{K}_{z,i}} E \left[\left| \psi(X_{t_i}) - \bar{z}_i^N(X_{t_i}) \right|^2 \right] \right. \\ &\quad \left. + (1 + \kappa^{-1}) C_{q,i}^2 \frac{\mathcal{D}K_{z,i}}{\Delta M_i} + (1 + \kappa) E \left[\left(\xi_i^z(X_{t_i}) \right)^2 \right] \right). \end{aligned} \quad (9)$$

Now using the tower property and a zero addition, we get

$$\begin{aligned} \Delta E \left[\left(\xi_i^z(X_{t_i}) \right)^2 \right] &= \Delta E \left[E \left[\frac{\Delta W_{i+1}}{\Delta} \left(\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \right) \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\ &= \Delta E \left[E \left[\frac{\Delta W_{i+1}}{\Delta} E \left[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \middle| \mathcal{F}_0^M, \mathcal{F}_{t_{i+1}} \right] \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\ &= \Delta E \left[E \left[\frac{\Delta W_{i+1}}{\Delta} \left(E \left[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \middle| \mathcal{F}_0^M, \mathcal{F}_{t_{i+1}} \right] \right. \right. \right. \\ &\quad \left. \left. - E \left[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \middle| \mathcal{F}_0^M, X_{t_i} \right] \right) \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\ &\leq \mathcal{D}E \left[E \left[\left(E \left[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \middle| \mathcal{F}_0^M, \mathcal{F}_{t_{i+1}} \right] \right. \right. \right. \\ &\quad \left. \left. - E \left[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \middle| \mathcal{F}_0^M, X_{t_i} \right] \right)^2 \middle| \mathcal{F}_0^M, X_{t_i} \right] \right] \\ &\leq \mathcal{D}E \left[E \left[E \left[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \middle| \mathcal{F}_0^M, \mathcal{F}_{t_{i+1}} \right]^2 \middle| \mathcal{F}_0^M, X_{t_i} \right] \right. \\ &\quad \left. - \mathcal{D}E \left[\Xi_i^{N,M}(\underline{X}_{t_i}) - \Xi_i^N(\underline{X}_{t_i}) \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right]. \end{aligned}$$

For the next step we have to distinguish between the time points again. If t_i and t_{i+1} are in the same segment, i.e., at all time points t_i such that $t_{i+1} \notin \bar{\pi}_\alpha$, it holds $\tau_\alpha(i) = \tau_\alpha(i+1)$ and we get for a $\Gamma > 0$ which will be specified later that

$$\begin{aligned} &\Delta E \left[\left(\xi_i^z(X_{t_i}) \right)^2 \right] \\ &\leq \mathcal{D}E \left[E \left[\bar{q}_{\tau_\alpha(i)}(X_{t_{\tau_\alpha(i)}}) - q_{\tau_\alpha(i)}^{N,M}(X_{t_{\tau_\alpha(i)}}) + \sum_{j=i+1}^{\tau_\alpha(i)} \int_{t_j}^{t_{j+1}} f(s, X_s, Y_s, Z_s) \right. \right. \\ &\quad \left. \left. - f(t_j, X_{t_j}, q_j^{N,M}, z_j^{N,M}) \right) ds \middle| \mathcal{F}_0^M, \mathcal{F}_{t_{i+1}} \right]^2 - \mathcal{D}E \left[E \left[\Xi_i^{N,M}(\underline{X}_i) - \Xi_i^N(\underline{X}_i) \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\ &\leq (1 + \Gamma \Delta) \mathcal{D}E \left[\left(\xi_{i+1}^q(X_{t_{i+1}}) \right)^2 \right] \\ &\quad + \mathcal{D} \left(1 + \frac{1}{\Delta \Gamma} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\ &\quad - \mathcal{D}E \left[\left(\xi_i^q(X_{t_i}) \right)^2 \right]. \end{aligned}$$

Like in the derivation of the error of the approximation of \bar{q}^N , the last inequality in the calculations above does not hold true when considering $E[\xi_i^z(X_{t_i})^2]$ at time points t_i at the end of a segment, i.e., t_i such that $t_{i+1} \in \bar{\pi}_\alpha$. However, by adding and subtracting a multiple of $E[(\xi_{i+1}^q(X_{t_{i+1}}))^2]$, we again get the similar bound

$$\begin{aligned} & \Delta E \left[(\xi_i^z(X_{t_i}))^2 \right] \\ & \leq \mathcal{D} (1 + \Gamma \Delta) (1 + \kappa) (1 + \epsilon) E \left[(\xi_{i+1}^q(X_{t_{i+1}}))^2 \right] + \mathcal{D} \left(1 + \frac{1}{\Delta \Gamma} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) \right. \right. \\ & \quad \left. \left. - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] - \mathcal{D} E \left[(\xi_i^q(X_{t_i}))^2 \right] \\ & \quad + \mathcal{D} (1 + \Gamma \Delta) \left(E \left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] - (1 + \kappa) (1 + \epsilon) E \left[(\xi_{i+1}^q(X_{t_{i+1}}))^2 \right] \right)_+ \end{aligned}$$

for these time points, with an additional error term that depends on the approximation of \bar{q}^N at the next time point t_{i+1} . As argued in the analysis of the approximation of \bar{q}^N , this term results from the single use of an ODP step in the discretization scheme that is used to connect two time segments.

Next we want to derive a global error bound for the approximation of \bar{z}^N . Since \bar{z}^N appears in the discretization scheme only as argument of the driver, we state this term as an averaged sum over the time steps rather than the maximum.

Lemma 6.8 *Let Γ be a positive constants and set $\lambda_i := (1 + \Gamma \Delta)^i ((1 + N^{\alpha-1})^{2|\{j \leq i: t_j \in \bar{\pi}_\alpha\}|})$ for $i \in \{0, \dots, N-1\}$. Then*

$$\begin{aligned} & \sum_{i=0}^{N-1} \Delta \lambda_i E \left[(\xi_i^z(X_{t_i}))^2 \right] \\ & \leq \mathcal{D} \sum_{i=0}^{N-2} \lambda_i \left(1 + \frac{1}{\Delta \Gamma} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\ & \quad + \lambda_N \mathcal{D} [N^{1-\alpha}] \max_{j \in \mathcal{J}} \left((1 + N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,j}} E \left[|\psi(X_{t_j} - \bar{q}_i^N(X_{t_j}))|^2 \right] + (1 + N^{1-\alpha}) \frac{C_{q,j}^2 K_{q,j}}{M_j} \right) \right. \\ & \quad \left. + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right). \end{aligned}$$

Proof First, note that $E[(\xi_{N-1}^z(X_{t_{N-1}}))^2] = 0$ by definition since $\Xi_{N-1}^{N,M} = \Xi_{N-1}^N$. Then, by plugging in the estimate for $\Delta E[(\xi_i^z(X_{t_i}))^2]$ from the analysis of the error propagation where we choose $\epsilon = \kappa = N^{\alpha-1}$ for all i and summing up we get

$$\begin{aligned} & \sum_{i=0}^{N-2} \Delta \lambda_i E \left[(\xi_i^z(X_{t_i}))^2 \right] \\ & \leq \mathcal{D} \sum_{i=0}^{N-2} \left(\lambda_{i+1} E \left[(\xi_{i+1}^q(X_{t_{i+1}}))^2 \right] + \lambda_i \left(1 + \frac{1}{\Delta \Gamma} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) \right. \right. \right. \\ & \quad \left. \left. - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] - \lambda_i E \left[(\xi_i^q(X_{t_i}))^2 \right] \end{aligned}$$

$$\begin{aligned}
& + \lambda_i(1 + \Gamma\Delta) \left(E \left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1,\infty}^2 \right] - (1 + N^{\alpha-1})^2 E \left[(\xi_{i+1}^q(X_{t_{i+1}}))^2 \right] \right)_+ \mathbb{1}_{\mathcal{J}}(i+1) \Big) \\
& \leq \mathcal{D} \sum_{i=0}^{N-2} \lambda_i \left(1 + \frac{1}{\Delta\Gamma} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \Big| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
& \quad + \mathcal{D} \lambda_N [N^{1-\alpha}] \sup_{j \in \mathcal{J}} \left(\inf_{\psi \in \mathcal{K}_{q,j}} E \left[|\psi(X_{t_j}) - \bar{q}_j^N(X_{t_j})|^2 \right] - (1 + N^{\alpha-1})^2 E \left[(\xi_j^q(X_{t_j}))^2 \right] \right)_+ .
\end{aligned}$$

Plugging in the estimate for $E[\|\bar{q}_i^N - q_i^{N,M}\|_{i,\infty}^2]$ from (7) derived in the proof of Lemma 6.7 finishes the proof. \square

6.4 Bounds for the driver-dependent terms

In this section, we derive a bound for the sum of the terms

$$E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \Big| \mathcal{F}^M, X_{t_i} \right]^2 \right]$$

over the time steps that appears in the bounds of Lemma 6.7 and 6.8. For this purpose, note that for any $i \in \{0, \dots, N-1\}$, it holds by Fubini's theorem

$$\begin{aligned}
& E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \Big| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
& \leq E \left[\left(\int_{t_{i+1}}^{t_{i+2}} E_i [f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N)] \right. \right. \\
& \quad \left. \left. + E [f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) \Big| \mathcal{F}_0^M, X_{t_i}] ds \right)^2 \right] \\
& \leq 2E \left[\left(\int_{t_{i+1}}^{t_{i+2}} E_i [f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N)] ds \right)^2 \right] \\
& \quad + 2\Delta^2 E \left[E [f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) \Big| \mathcal{F}_0^M, X_{t_i}]^2 \right].
\end{aligned}$$

Then, by the Lipschitz assumption on f we get

$$\begin{aligned}
& E \left[E [f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) \Big| \mathcal{F}_0^M, X_{t_i}]^2 \right] \\
& \leq 2L_f^2 E \left[E [|\bar{q}_{i+1}^N(X_{t_{i+1}}) - q_{i+1}^{N,M}(X_{t_{i+1}})| \Big| \mathcal{F}_0^M, X_{t_i}]^2 \right] \\
& \quad + E [|\bar{z}_{i+1}^N(X_{t_{i+1}}) - z_{i+1}^{N,M}(X_{t_{i+1}})| \Big| \mathcal{F}_0^M, X_{t_i}]^2 \\
& \leq 2L_f^2 \left(E [\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1,\infty}^2] + E [\|\bar{z}_{i+1}^N - z_{i+1}^{N,M}\|_{i+1,\infty}^2] \right).
\end{aligned}$$

Hence it holds

$$\begin{aligned}
& \sum_{i=0}^{N-2} \lambda_i E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
& \leq \sum_{i=0}^{N-2} \lambda_i 2E \left[\left(\int_{t_{i+1}}^{t_{i+2}} E_i [f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N)] ds \right)^2 \right] \\
& \quad + \sum_{i=0}^{N-2} \lambda_i 4\Delta^2 L_f^2 \left(E \left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] + E \left[\|\bar{z}_{i+1}^N - z_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] \right) \\
& \leq \sum_{i=0}^{N-2} \lambda_i 2E \left[\left(\int_{t_{i+1}}^{t_{i+2}} E_i [f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N)] ds \right)^2 \right] \\
& \quad + 4\Delta L_f^2 T \max_{0 \leq i \leq N-1} \lambda_i E \left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] + 4\Delta L_f^2 \sum_{i=1}^{N-1} \lambda_i \Delta E \left[\|\bar{z}_{i+1}^N - z_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] \\
& \leq 2\lambda_N \mathcal{R}^N + 4\Delta L_f^2 T \max_{0 \leq i \leq N-1} \lambda_i E \left[\|\bar{q}_{i+1}^N - q_{i+1}^{N,M}\|_{i+1, \infty}^2 \right] + 4\Delta L_f^2 \sum_{i=1}^{N-1} \lambda_i \Delta E \left[\|\bar{z}_{i+1}^N - z_{i+1}^{N,M}\|_{i+1, \infty}^2 \right]
\end{aligned}$$

with

$$\mathcal{R}^N := \sum_{i=0}^{N-2} E \left[\left(\int_{t_{i+1}}^{t_{i+2}} E_i [f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, \bar{q}_{i+1}^N, \bar{z}_{i+1}^N)] ds \right)^2 \right]$$

as defined in Theorem 4.1. The term \mathcal{R}^N does not depend on our approximation of the BSDE but only on the real solution Y, Z , the semi-continuous versions \bar{q}^N, \bar{z}^N and the solution of the forward SDE X . It can be bounded in different ways depending on the regularity of these functions, which leads to the different bounds of the total quadratic error in Theorem 4.2 and 4.3.

6.5 Final error bounds

Using the bounds derived throughout this section, we are now ready to proof Theorem 4.1.

Proof of Theorem 4.1

In the following calculations, c denotes a positive constant that does not depend on N and may change from line to line. First, we can write the quadratic error as

$$\begin{aligned}
& \max_{0 \leq i \leq N-1} E \left[|\bar{q}_i^N(X_{t_i}) - q_i^{N,M}(X_{t_i})|^2 \right] + \sum_{i=0}^{N-1} \Delta E \left[|\bar{z}^N(X_{t_i}) - z_i^{N,M}(X_{t_i})|^2 \right] \\
& \leq \max_{0 \leq i \leq N-1} \lambda_i E \left[|\bar{q}_i^N(X_{t_i}) - q_i^{N,M}(X_{t_i})|^2 \right] + \sum_{i=0}^{N-2} \Delta \lambda_i E \left[|\bar{z}^N(X_{t_i}) - z_i^{N,M}(X_{t_i})|^2 \right] \\
& = \max_{0 \leq i \leq N-1} \lambda_i E \left[\|\bar{q}_i^N - q_i^{N,M}\|_{i, \infty}^2 \right] + \sum_{i=0}^{N-1} \Delta \lambda_i E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i, \infty}^2 \right].
\end{aligned}$$

We can now estimate this term by Lemma 6.5 with $\epsilon = 1$ as

$$\begin{aligned}
& \max_{0 \leq i \leq N-1} \lambda_i E \left[\|\bar{q}_i^N - q_i^{N,M}\|_{i,\infty}^2 \right] + \sum_{i=0}^{N-1} \Delta \lambda_i E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i,\infty}^2 \right] \\
& \leq \max_{0 \leq i \leq N-1} \left(2\lambda_i E \left[\|\bar{q}_i^N - q_i^{N,M}\|_{i,M}^2 \right] + \lambda_i \frac{C_1 K_{q,i} \log(C_2 M_i)}{M_i} \right) \\
& \quad + \sum_{i=0}^{N-2} 2\Delta \lambda_i E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i,M}^2 \right] + \Delta \lambda_i \frac{C_1 K_{z,i} \log(C_2 M_i)}{\Delta M_i}.
\end{aligned}$$

By the inequalities (6) and (9) we then get with the choice $\kappa = 1$

$$\begin{aligned}
& \max_{0 \leq i \leq N-1} \lambda_i E \left[\|\bar{q}_i^N - q_i^{N,M}\|_{i,\infty}^2 \right] + \sum_{i=0}^{N-1} \Delta \lambda_i E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i,\infty}^2 \right] \\
& \leq \lambda_N \max_{0 \leq i \leq N-1} \left(4 \frac{C_{q,i} K_{q,i}}{M_i} + 2 \inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + \frac{C_1 K_{q,i} \log(C_2 M_i)}{M_i} \right) \\
& \quad + \sum_{i=0}^{N-1} \Delta \lambda_i \left(4 \frac{C_{z,i}^2 K_{z,i}}{\Delta M_i} + 2 \inf_{\psi \in \mathcal{K}_{z,i}} E \left[|\bar{z}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + \frac{C_1 K_{z,i} \log(C_2 M_i)}{\Delta M_i} \right) \\
& \quad + 4 \left(\max_{0 \leq i \leq N-1} \lambda_i E \left[(\xi_i^q(X_{t_i}))^2 \right] + \sum_{i=0}^{N-1} \Delta \lambda_i E \left[(\xi_i^z(X_{t_i}))^2 \right] \right).
\end{aligned}$$

Now the bounds in Lemma 6.7 and Lemma 6.8 yield

$$\begin{aligned}
& \max_{0 \leq i \leq N-1} \lambda_i E \left[(\xi_i^q(X_{t_i}))^2 \right] + \sum_{i=0}^{N-1} \Delta \lambda_i E \left[(\xi_i^z(X_{t_i}))^2 \right] \\
& \leq \sum_{i=0}^{N-2} \left(1 + \frac{1}{\Delta \Gamma} \right) \lambda_i E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
& \quad + [N^{1-\alpha}] \lambda_N \max_{j \in \mathcal{J}} \left((1 + N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + (1 + N^{1-\alpha}) \frac{C_{q,i}^2 K_{q,j}}{M_j} \right) \right. \\
& \quad \left. + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right) + \mathcal{D} \sum_{i=0}^{N-2} \lambda_i \left(1 + \frac{1}{\Delta \Gamma} \right) E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) \right. \right. \\
& \quad \left. \left. - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] + \lambda_N \mathcal{D} [N^{1-\alpha}] \max_{j \in \mathcal{J}} \left((1 + N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) \right. \right. \right. \\
& \quad \left. \left. - \psi(X_{t_i})|^2 \right] + (1 + N^{1-\alpha}) \frac{C_{q,i}^2 K_{q,j}}{M_j} \right) + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \left. \right) \\
& = \sum_{i=0}^{N-2} (1 + \mathcal{D}) \left(1 + \frac{1}{\Delta \Gamma} \right) \lambda_i E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
& \quad + (1 + \mathcal{D}) [N^{1-\alpha}] \lambda_N \max_{j \in \mathcal{J}} \left((1 + N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + (1 + N^{1-\alpha}) \frac{C_{q,i}^2 K_{q,j}}{M_j} \right) \right. \\
& \quad \left. + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right).
\end{aligned}$$

By plugging in the bounds derived in Section 6.4, we can estimate this term further as

$$\begin{aligned}
& \max_{0 \leq i \leq N-1} \lambda_i E \left[(\xi_i^q(X_{t_i}))^2 \right] + \sum_{i=0}^{N-1} \Delta \lambda_i E \left[(\xi_i^z(X_{t_i}))^2 \right] \\
& \leq \sum_{i=0}^{N-2} (1+\mathcal{D}) \left(1 + \frac{1}{\Delta \Gamma} \right) \lambda_i E \left[E \left[\int_{t_{i+1}}^{t_{i+2}} f(s, X_s, Y_s, Z_s) - f(t_{i+1}, X_{t_{i+1}}, q_{i+1}^{N,M}, z_{i+1}^{N,M}) \, ds \middle| \mathcal{F}_0^M, X_{t_i} \right]^2 \right] \\
& \quad + (1+\mathcal{D}) \lceil N^{1-\alpha} \rceil \lambda_N \max_{j \in \mathcal{J}} \left((1+N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + (1+N^{1-\alpha}) \frac{C_{q,i}^2 K_{q,j}}{M_j} \right) \right. \\
& \quad \left. + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right) \\
& \leq \left[\left(\Delta + \frac{1}{\Gamma} \right) (1+\mathcal{D}) \right] 4(T \vee 1) L_f^2 \left(\max_{0 \leq i \leq N-1} \lambda_i E \left[\|\bar{q}_i^N - q_i^{N,M}\|_{i,\infty}^2 \right] + \sum_{i=0}^{N-2} \Delta \lambda_i E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i,\infty}^2 \right] \right) \\
& \quad + \left[\left(\Delta + \frac{1}{\Gamma} \right) (1+\mathcal{D}) \right] \Delta^{-1} \lambda_N \mathcal{R}^N + \lceil N^{1-\alpha} \rceil \lambda_N (1+\mathcal{D}) \max_{j \in \mathcal{J}} \left((1+N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) \right. \right. \right. \\
& \quad \left. \left. \left. - \psi(X_{t_i}) \right|^2 \right] + (1+N^{1-\alpha}) \frac{C_{q,j}^2 K_{q,j}}{M_j} \right) + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right).
\end{aligned}$$

Now, assuming that N and Γ are sufficiently large such that $[(\Delta + \frac{1}{\Gamma})(1+\mathcal{D})]16L^2(T \vee 1) \leq \frac{1}{2}$, we have

$$\begin{aligned}
& \max_{0 \leq i \leq N-1} E \left[\|\bar{q}_i^N - q_i^{N,M}\|_{i,\infty}^2 \right] \lambda_i + \sum_{i=0}^{N-1} \lambda_i E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i,\infty}^2 \right] \Delta \\
& \leq \frac{1}{2} \left(\max_{0 \leq i \leq N-1} E \left[\|\bar{q}_i^N - q_i^{N,M}\|_{i,\infty}^2 \right] \lambda_i + \sum_{i=0}^{N-1} \lambda_i E \left[\|\bar{z}_i^N - z_i^{N,M}\|_{i,\infty}^2 \right] \Delta \right) \\
& \quad + c \lambda_N \lceil N^{1-\alpha} \rceil \max_{j \in \mathcal{J}} \left((1+N^{\alpha-1}) \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + (1+N^{1-\alpha}) \frac{C_{q,j}^2 K_{q,j}}{M_j} \right) \right. \\
& \quad \left. + \frac{N^{1-\alpha} C_1 K_{q,j} \log(C_2 M_j)}{M_j} \right) + c \lambda_N \max_{0 \leq i \leq N-1} \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\bar{q}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + \frac{K_{q,i}}{M_i} \right. \\
& \quad \left. + \frac{K_{q,i} \log(M_i)}{M_i} + \inf_{\psi \in \mathcal{K}_{z,i}} E \left[|\bar{z}_i^N(X_{t_i}) - \psi(X_{t_i})|^2 \right] + \frac{K_{z,i}}{\Delta M_i} + \frac{K_{z,i} \log(M_i)}{\Delta M_i} \right) + c \Delta^{-1} \lambda_N \mathcal{R}^N.
\end{aligned}$$

Considering that λ_N is bounded by a constant independent of N , since

$$\begin{aligned}
\lambda_N & = \left(1 + \frac{T\Gamma}{N} \right)^N (1+N^{\alpha-1})^{2\lceil N^{1-\alpha} \rceil} \\
& \leq e^{T\Gamma N^{-1}N} e^{2\lceil N^{1-\alpha} \rceil N^{\alpha-1}} = e^{T\Gamma+4},
\end{aligned}$$

this implies

$$\begin{aligned}
& \max_{0 \leq i \leq N-1} E \left[|\bar{q}_i^N(X_{t_i}) - q_i^{N,M}(X_{t_i})|^2 \right] + \sum_{i=0}^{N-2} \Delta E \left[|\bar{z}_i^N(X_{t_i}) - z_i^{N,M}(X_{t_i})|^2 \right] \\
& \leq c \max_{i \in \mathcal{J}} \left(N^{1-\alpha} \inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\psi(X_{t_i}) - \bar{q}_i^N(X_{t_i})|^2 \right] + N^{2-2\alpha} \frac{K_{q,i}}{M_i} + N^{2-2\alpha} \frac{K_{q,i} \log(C_2 M_i)}{M_i} \right) \\
& \quad + c \max_{0 \leq i \leq N-1} \left(\inf_{\psi \in \mathcal{K}_{q,i}} E \left[|\psi(X_{t_i}) - \bar{q}_i^N(X_{t_i})|^2 \right] + \inf_{\psi \in \mathcal{K}_{z,i}} E \left[|\psi(X_{t_i}) - \bar{z}_i^N(X_{t_i})|^2 \right] \right) \\
& \quad + \frac{K_{q,i}}{M_i} + N \frac{K_{z,i}}{M_i} + \frac{K_{q,i} \log(C_2 M_i)}{M_i} + N \frac{K_{z,i} \log(C_2 M_i)}{M_i} \\
& \quad + cN\mathcal{R}^N,
\end{aligned}$$

and finishes the proof. \square

References

- [1] Bender, C. and Denk, R., [A forward scheme for backward SDEs](#), Stochastic Processes and their Applications, 2007, 117(12): 1793–1812.
- [2] Bouchard, B. and Touzi, N., [Discrete-time approximation and Monte-Carlo simulation of backward stochastic differential equations](#), Stochastic Processes and their Applications, 2004, 111(2): 175–206.
- [3] Chessari, J., Kawai, R., Shinozaki, Y. and Yamada, T., Numerical methods for backward stochastic differential equations: A survey, Probability Surveys, 2023, 20: 486–567.
- [4] Crisan, D. and Delarue, F., [Sharp derivative bounds for solutions of degenerate semi-linear partial differential equations](#), Journal of Functional Analysis, 2012, 263(10): 3024–3101.
- [5] Douglas, J., Ma, J. and Protter, P., Numerical methods for forward-backward stochastic differential equations, The Annals Applied Probability, 1996, 6(3): 940–968.
- [6] E, W., Huttenhaler, M., Jentzen, A. and Kruse, T., [Multilevel Picard iterations for solving smooth semilinear parabolic heat equations](#), Partial Differential Equations and Applications, 2021, 2: 80.
- [7] El Karoui, N., Peng, S. and Quenez, M. C., [Backward stochastic differential equations in finance](#), Mathematical Finance, 1997, 7(1): 1–71.
- [8] Gobet, E. and Labart, C., [Error expansion for the discretization of backward stochastic differential equations](#), Stochastic Processes and their Applications, 2007, 117(7): 803–829.
- [9] Gobet, E., Lemor, J.-P. and Warin, X., A regression-based Monte Carlo method to solve backward stochastic differential equations, The Annals Applied Probability, 2005, 15(3): 2172–2202.
- [10] Gobet, E., López-Salas, J. G., Turkedjiev, P. and Vázquez, C., [Stratified regression Monte-Carlo scheme for semilinear PDEs and BSDEs with large scale parallelization on GPUs](#), SIAM Journal on Scientific Computing, 2016, 38(6): C652–C677.
- [11] Gobet, E. and Turkedjiev, P., Linear regression MDP scheme for discrete backward stochastic differential equations under general conditions, Mathematics of Computation, 2016, 85(299): 1359–1391.
- [12] Guyon, J. and Henry-Labordère, P., Nonlinear Option Pricing, Chapman & Hall/CRC Financial Mathematics Series, CRC Press, Boca Raton, 2014.
- [13] Györfi, L., Kohler, M., Krzyżak, A. and Walk, H., A Distribution-Free Theory of Nonparametric Regression, Springer Series in Statistics, Springer, New York, 2002.
- [14] Huttenhaler, M., Jentzen, A., Kruse, T. and Nguyen, T. A., Overcoming the curse of dimensionality in the numerical approximation of backward stochastic differential equations, Journal of Numerical Mathematics, 2023, 31(1): 1–28.

- [15] Kallenberg, O., *Foundations of Modern Probability*, Springer, Cham, 1997.
- [16] Karatzas, I. and Shreve, S., *Brownian Motion and Stochastic Calculus*, Springer, New York, 2012.
- [17] Kloeden, P. E. and Platen, E., *Numerical Solution of Stochastic Differential Equations*, Springer, Berlin, Heidelberg, 1992.
- [18] Lemor, J.-P., Gobet, E. and Warin, X., Rate of convergence of an empirical regression method for solving generalized backward stochastic differential equations, *Bernoulli*, 2006, 12(5): 889–916.
- [19] Ma, J., Protter, P. and Yong, J. M., [Solving forward-backward stochastic differential equations explicitly—a four step scheme](#), *Probability Theory and Related Fields*, 1994, 98(3): 339–359.
- [20] Ma, J., Shen, J. and Zhao, Y., [On numerical approximations of forward-backward stochastic differential equations](#), *SIAM Journal on Numerical Analysis*, 2008, 46(5): 2636–2661.
- [21] Milstein, G. N. and Tretyakov, M. V., [Numerical algorithms for forward-backward stochastic differential equations](#), *SIAM Journal on Scientific Computing*, 2006, 28(2): 561–582.
- [22] Pardoux, E. and Peng, S., Backward stochastic differential equations and quasilinear parabolic partial differential equations, In: Rozovskii, B. L. and Sowers, R. B.(eds.), *Stochastic Partial Differential Equations and their Applications* (Charlotte, NC, 1991), *Lecture Notes in Control and Information Sciences*, Springer, Berlin, 1992, 176: 200–217.
- [23] Zhang, J., A numerical scheme for BSDEs, *The Annals Applied Probability*, 2004, 14(1): 459–488.