

REVIEW

Systems and synthetic biology approaches in understanding biological oscillators

Zhengda Li^{1,2} and Qiong Yang^{1,2,*}

¹ Department of Biophysics, University of Michigan, Ann Arbor, MI 48109, USA

² Department of Computational Medicine & Bioinformatics, University of Michigan, Ann Arbor, MI 48109, USA

* Correspondence: qiongy@umich.edu

Received December 16, 2016; Revised April 17, 2017; Accepted April 18, 2017

Background: Self-sustained oscillations are a ubiquitous and vital phenomenon in living systems. From primitive single-cellular bacteria to the most sophisticated organisms, periodicities have been observed in a broad spectrum of biological processes such as neuron firing, heart beats, cell cycles, circadian rhythms, etc. Defects in these oscillators can cause diseases from insomnia to cancer. Elucidating their fundamental mechanisms is of great significance to diseases, and yet challenging, due to the complexity and diversity of these oscillators.

Results: Approaches in quantitative systems biology and synthetic biology have been most effective by simplifying the systems to contain only the most essential regulators. Here, we will review major progress that has been made in understanding biological oscillators using these approaches. The quantitative systems biology approach allows for identification of the essential components of an oscillator in an endogenous system. The synthetic biology approach makes use of the knowledge to design the simplest, *de novo* oscillators in both live cells and cell-free systems. These synthetic oscillators are tractable to further detailed analysis and manipulations.

Conclusion: With the recent development of biological and computational tools, both approaches have made significant achievements.

Keywords: biological oscillators; synthetic oscillators; circuit design principles


Author summary: Biological oscillators drive neuron firing, cell cycles, sleep-wake patterns, and many other essential processes in living systems. Quantitative investigation of how biological oscillators are designed can help us better understand and cure diseases. Over the past decades, development of techniques in the field of systems and synthetic biology has paved the way to decipher the mechanisms behind these oscillators. Here we review a series of experimental and theoretical studies in quantitative system and synthetic biology to discuss the progress that has been made in studying biological oscillators.

INTRODUCTION

Studies over the past decades have made major progress in identifying genes, proteins, and metabolites that are involved in the rhythmic phenomena. A typical pathway of an oscillator that can be found in databases nowadays is rather complicated. For example, the cell cycle pathway in KEGG (Figure 1A) contains hundreds of molecules.

Biological oscillators are not only complicated, but also

appear to be extremely diverse in their dynamic properties and compositions. From sub-second neural spikes, to daily circadian rhythms, to annual reproduction in plants and animals, the periods of oscillators span orders of magnitudes. Molecules that drive these events are very different from oscillators to oscillators: i) cytoplasmic oscillators, in particular glycolytic oscillators, are formed by protein enzymes and small molecule metabolites that mainly interact within the cytoplasm; ii) membrane



04110_46516
© Kaenlisia Laboratories

















Oscillators	Network descriptions	Period	Oscillator type	Function	Ref.
 Cardiomyocytes calcium spike		~1 s	Membrane oscillators	Endogenous pacemaker	Periasamy <i>et al.</i> 2008 [3]; Liu <i>et al.</i> 2008 [4]
 Xenopus embryo cell cycle		10 min–24 h	Cytoplasmic oscillators	Endogenous pacemaker	Yang <i>et al.</i> 2013 [1]
 Zebrafish segmentation clock		25 min (zebrafish) 90 min (chicks) 2 h (mice)	Genetic oscillators	Endogenous pacemaker	Mara and Holley 2007 [6]
 Mammal circadian clock		24 h	Genetic oscillators	Endogenous pacemaker	Bell-Pedersen <i>et al.</i> 2005 [7]
 Action potential		0.001 s–10 s	Membrane oscillators	Signal processing and transduction	Hodgkin and Huxley 1952 [8]
 p38 oscillation		~60 min	Cytoplasmic oscillators	Signal processing and transduction	Tomida <i>et al.</i> 2015 [9]
 NF-κB spike		~100 min	Genetic oscillators	Signal processing and transduction	Zambrano <i>et al.</i> 2016 [10]
 p53 oscillation		~6 h	Genetic oscillators	Signal processing and transduction	Batchelor <i>et al.</i> 2011 [11]

Figure 1. Quantitative studies have revealed core architectures of biological oscillators that are both complicated and diverse. (A) Left: The cell cycle pathway diagram of *Xenopus laevis* adopted from Kyoto Encyclopedia of Genes and Genomes (KEGG) database. Right: A simplified version of *Xenopus* early embryonic cell cycle circuitry that contains only the most essential components [1,2]. (B) Examples of biological oscillators and their network structures. From top to bottom: calcium spikes in cardiomyocytes [3,4], embryonic cell cycles in *Xenopus* [1,2], segmentation clock in Zebrafish [5,6], and circadian clock in Mammals [7], action potential in neuron [8], p38 oscillations in HeLa cells [9], NF-κB spikes in fibroblasts [10], p53 oscillation in human cell lines [11]. Their periods span orders of magnitude.

oscillators, such as action potentials in neural and cardiac pacemaker cells, are formed by regulatory ion channels restricted to the membrane; iii) genetic oscillators are composed of a set of genes that regulate each other inside cells.

These oscillators perform several functions in general (Figure 1B). Most oscillators function as an endogenous pacemaker. Circadian clocks, for example, exist in almost all organisms on earth and help them adapt to the natural periodicity of the day-night alternations, by orchestrating their intrinsic gene expressions with a period of approximately 24 hours [7,12,13]. Heartbeat [14], respiration [15], and cell cycle [16–18], etc., are other examples in this category. The pacemaker oscillators also play a role in the spatial organization of development. Through an excitable medium, a local oscillatory signal may trigger waves that propagate over a large distance much faster than through pure diffusion [19]. Examples of trigger waves include polar regeneration in *Acetabularia* [20], cAMP waves in the aggregation and differentiation of *Dictyostelium discoideum* [21], and mitotic waves recently reconstituted in *Xenopus* cell-free extracts [22]. Alternatively, in multidimensional or metameric systems, the phase differences among cell-autonomous periodic events in cells as a function of their spatial locations, resulted in another type of wave, called phase wave. These may be exemplified by pattern formation of hypostome in *Hydra* [23] and periodic tissue morphogenesis of vertebrates [5,24]. In addition to being a pacemaker, oscillators can also generate frequency-modulated signals to regulate their downstream gene expression levels and influence cellular fate decisions [25,26] in an accurate manner through information transduction. Examples of these have been seen in frequency coding of neurons [27] and a number of signal transduction pathways that have oscillatory dynamics, such as NF-kappaB [28], p53 [11,29], p38 [9], Ca signaling [30,31]. These oscillators are flexible in tuning their frequencies.

The question whether a common mechanism is shared by all these seemingly different oscillators has stimulated much interest and its answer may allow for a fundamental understanding of such widespread phenomena. However, the high complexity and broad diversity of biological oscillators make it extremely challenging to identify the fundamental mechanisms. To address such difficulty, approaches in systems and synthetic biology have been effective in dissecting or creating an oscillatory circuit and analyzing its functions beyond the molecular level. In this review, we attempt to summarize the major progress in quantitative systems biology and synthetic biology that aim to delineate the mechanisms and understand oscillatory properties at the systems level.

QUANTITATIVE ANALYSIS OF BIOLOGICAL OSCILLATORS IN ENDOGENOUS SYSTEMS

The biological oscillators are amenable to quantitative analysis both in theory and experiments. One of the first characterized biological clocks is the action potential in a giant nerve fibre of *Loligo*. In their Nobel Prize winning studies, Alan Hodgkin and Andrew Huxley carried out a series of measurements of the nerve cell membrane ionic current under a “voltage clamp” [32–36]. Under certain conditions, the nervous system exhibits an oscillatory behavior, with electrical pulses repeatedly generated in response to a stimulus. To interpret the findings, they developed a set of ordinary differential equations (ODEs) [8] that include multiple voltage-dependent currents of ion (e.g., potassium and sodium) channels, known as the Hodgkin-Huxley model. The model marks the starting point for theoretical biophysics of action potential.

About a decade later, another oscillator is observed in the metabolism pathway glycolysis, which drives repetitive fluctuations of concentrations of metabolites intermediates. This time, the oscillations were not investigated in the membrane of a giant cell, but in a whole cell suspension [37] and a cytosolic cell-free system [38,39] of yeast cells *Saccharomyces carlsbergensis*. Although these early metabolic assays only captured damped sinusoidal oscillations, theoretical work describing six reaction equations using ODE models for phosphofructokinase and the associated glycolytic intermediates predicted the existence of self-sustained limit-cycle behaviors in glycolytic oscillations [40]. Such limit-cycle oscillations were soon observed in a later experiment [41]. Other metabolite oscillations have also been observed in cAMP synthesis [42] and peroxidase-oxidase reactions [43].

The first genetic oscillator in theory was proposed by Brian Goodwin back in 1963 [44], shortly after the model of gene regulation developed by Francois Jacob and Jacques Monod. This classical Goodwin oscillator contains only a single gene whose product represses itself after a sufficient delay, allowing for periodic gene expression to occur. Remarkably, this theoretical work predicts that a genetic circuit is able to generate the limit cycle oscillations, even before any real genetic oscillator is discovered experimentally. The experimental investigations on genetic oscillators lagged behind until the development of genetic engineering techniques in modern molecular biology and the rapidly growing popularity of luminescence and fluorescence microscopy techniques. One of the most characterized genetic oscillators is circadian clock, which has been found in all eukaryotes

and some prokaryotes such as cyanobacteria. Circadian clocks endogenously drive cell-autonomous oscillations roughly once per day, so that organisms have the ability to anticipate the time of day. Studies on circadian clocks have been performed on various organisms (*Synechococcus* [45], *Neurospora* [46], *Drosophila* [47], mouse [48] and *Arabidopsis* [49]).

These studies have successfully combined experimental measurements and mathematical modeling to enable understanding of how each of these biological oscillators functions in great quantitative details. Following these studies, the quantitative approaches have been applied to the discovery and characterization of many more oscillators. For readers interested in a thorough and detailed description of biological oscillators, the classical book and review article by Goldbeter [50,51] are recommended.

Although major progress has been made in understanding relevant components and functions of biological oscillators, challenges for further quantitative analysis of these oscillators in living systems are obvious. One major obstacle is that the core architecture of an oscillator is often embedded in a significantly more complicated network. It also tends to interfere with other pathways or couple with other oscillators [52–54]. These make it hard to isolate the central oscillator circuitry for investigation. Another obstacle is the limited capability of dissecting feedback loops and manipulating oscillatory behaviors in living systems. To address these problems, synthetic biologists create well-defined *de novo* oscillatory circuitry in a system that is amenable to detailed analysis and accurate control. In the next two sections of this review, we introduce how synthetic biology has become an increasingly important alternative approach that will complement studies in the endogenous system. We will specifically focus on recent development of synthetic oscillators in both live cells and cell-free extracts.

BUILD TO UNDERSTAND — A SYNTHETIC BIOLOGY APPROACH IN THE UNDERSTANDING OF BIOLOGICAL OSCILLATORS

As reflected by Richard Feynman’s 1988 quote “What I cannot create, I do not understand”, synthetic biologists seek to improve understanding of oscillators by making them from scratch. The extensive quantitative studies on various biological clocks have revealed that, although the molecular forms can look vastly different from oscillators to oscillators, their network structures share common features. Notably, most of them seem to reduce down to a core architecture that contains positive plus negative feedback loops. This may support the possibility of building a simple oscillator that contains only the most

essential core components. To build such an oscillator, an inevitable question is — what are the minimum requirements for generating sustained oscillations? The answer to this question may also help us identify fundamental mechanisms that are shared by all biological oscillators.

Basic requirements of making oscillators in theory

Ever since the early discoveries of biological oscillations, theorists have started to pursue the answers to such question by building simplest possible models that capture fundamental mechanisms of various oscillatory systems. Here we will mainly focus on the development of ODE models which have the longest history and are one of the most widely used models to describe oscillatory behaviors. Some other method like Boolean model is also useful when dealing with highly complicated networks with few known parameters [55]. Stochastic differential equation models have also been used to reliably explain more complicated biological clock behaviors [56,57]. Right after the discovery of metabolic oscillations, Ilya Prigogine approached three biochemical oscillators of distinct catalytic properties in physicochemical terms by arguing that all these biochemical oscillations are not different than non-equilibrium spatial and temporal self-organization as dissipative structures in chemical systems [58]. This view provides a conceptual framework that supports the later research of unifying all biological oscillators [51].

Throughout the years, efforts have been made in simplifying existing detailed models, the complexity of which may obscure the fundamental design principles underlying the core architecture of an oscillator. In contrast to the complicated models, a model with only the most essential components is more accessible to experimental validation as it requires less parameters to measure to build an accurate model. The original Hodgkin-Huxley investigations of the action potential [8] describes in a detailed manner the voltage-current relationship based on experimental observations, which requires more than twenty parameters. It was simplified in the FitzHugh-Nagumo (FHN) model [59] and later in the Morris-Lecar model [60], where two differential equations describe the system as coupled positive and negative feedback loops. Importantly, despite the simplifications, these models capture the main dynamic responses, and by adding diffusion, the FHN model generates trigger wave propagations as seen in axons [19].

Simple models have also helped postulate the fundamental requirements for designing an oscillator. Since the design of the Goodwin oscillator [44,61], the inhibitory feedback loop and a source of delay in this feedback have been postulated as two required elements for limit cycle oscillations [62]. However, a time-delayed negative feedback could not explain the noise-resistance behavior












Molecular network description	Organism	Period	Year
	<i>E. coli</i>	~150 min	Elowitz and Leibler, 2000 [66] Niederholtmeyer <i>et al.</i> 2015 [79] Potvin-Trottier <i>et al.</i> 2016 [75]
	<i>E. coli</i>	10 h–20 h	Atkinson <i>et al.</i> 2003 [67]
	<i>E. coli</i>	45 min–3 h	Fung <i>et al.</i> 2005 [69]
	<i>E. coli</i>	13 min–100 min	Stricker <i>et al.</i> 2008 [68] Mondragón-Palomino <i>et al.</i> 2011 [72] Hussain <i>et al.</i> 2014 [74] Butzin <i>et al.</i> 2016 [73]
	Mamalian cell line (CHO K1)	2 h–8 h	Tigges <i>et al.</i> 2009 [70]
	<i>E. coli</i>	~50 min	Dies <i>et al.</i> 2016 [84]
	<i>E. coli</i> population	52 min–90 min	Danino <i>et al.</i> 2010 [76]
	Mamalian cell line (CHO K1)	26 h	Tigges <i>et al.</i> 2010 [71]
	Mamalian cell line (MCF7)	~6 h	Toettcher <i>et al.</i> 2010 [78]
	<i>E. coli</i> population	100 min–250 min	Chen <i>et al.</i> 2015 [85]
	<i>E. coli</i> extract	2 h–20 h	Niederholtmeyer <i>et al.</i> 2015 [79]

Figure 2. Examples of synthetic oscillators. For each synthetic oscillator, the information such as its molecular network structure, the organism that the oscillator is built in, and the period, has been listed [66–79].

in circadian rhythms. The activator-repressor oscillator was thus proposed by Barkai and Leibler [63]. It was a two-component gene circuit that contains both an activator and a repressor. In addition to the negative feedback loop formed by the activator activating its own repressor, the activator can also auto-activate itself resulting in a self-positive feedback loop. Remarkably, it turns out later that such “design principle”, i.e., the coupled positive and negative feedback loops, is widely found in biological oscillators. In the 2008 review article, Novak and Tyson have extensively discussed the general requirements for biochemical oscillations [64]. Besides the negative feedback and time delay, sufficient non-linearity and properly balanced timescales of opposing chemical reactions are also reported to be indispensable. In addition, the positive feedback is helpful to amplify and delay the negative-feedback signal, which makes the oscillator more robust. A computational study has suggested that adding a positive feedback may increase the robustness and frequency range of the system [65]. All together, these computational studies have provided key theoretical bases for creating synthetic oscillators, and experiments have been demanded for verifications.

Pioneering work of making simple oscillators

With the development of the genetic engineering tools, creating a *de novo* designed oscillator in real biological systems becomes possible. A list of well-known synthetic oscillators has been summarized in Figure 2 and will be described in more details below.

In 2000, Elowitz and Leibler, in a landmark study in synthetic biology, constructed the first genetic oscillator in *Escherichia coli* cells, named repressilator [66]. In this oscillator, three negative transcriptional regulators, TetR, λ CI and LacI, repress each other to form a delayed negative feedback loop. A green fluorescence protein GFP under the control of TetR promoter has been used to report the oscillatory behavior of the system, confirming that a single negative feedback is sufficient for generating oscillations. However, it was not robust, with only 40% of the cells oscillatory. Based on this work, many new oscillators have been proposed aiming for better performance.

In 2003, Atkinson *et al.* constructed for the first time the activator-repressor genetic oscillator [67], reminiscent of the theoretical clock of Barkai and Leibler [63]. Turbidostat cell cultures of *E. coli* containing this oscillator achieved synchronized oscillations. However, the oscillations become damped over time, possibly due to a loss of synchrony among cells. Further development in bacteria is made by Stricker *et al.* [68]. Besides a negative feedback formed by *araC* and *lacI*, *araC* also activates itself and *lacI* inhibits itself. This time, the

oscillator was tested in microfluidic devices using single-cell fluorescence microscopy. Unlike the ensemble measurements, the ability of tracking single cells relieves the requirement of synchronization of cells. Single-cell data has shown that the period of this oscillator can be as short as 13 min and the oscillation is self-sustained and robust.

The first metabolic circuit using glycolytic flux to generate oscillations was designed in *E. coli* by Fung *et al.*, called metabolator [69]. Like all genetic oscillators at the time, this synthetic metabolic oscillator was designed in bacteria cells. The first synthetic oscillator in mammalian cells was reported by Tigges *et al.* in 2009 [70]. Using auto-regulated sense-antisense transcription control, they built a molecular network that resembles the typical activator-repressor circuit, where tetracycline-dependent transactivator (tTA) functions as an activator and pristinamycin-dependent transactivator (PIT) a repressor. Further development of this system has led to an oscillator with frequency comparable to that of a circadian clock [71].

These pioneer studies of synthetic oscillators play an important role in testing the minimal design principles postulated by theoretical studies. In a review article, Purcell *et al.* have focused on several basic *in silico* designs that have been implemented in synthetic genetic oscillators [80]. The simple synthetic oscillators have also paved the way for further more complicated designs.

Making more complicated synthetic oscillators and their applications

At present, while new designs continue to expand the list of synthetic oscillators, the focus of the research has shifted from making new circuits to improving the existing oscillators with new functions or applications.

First, more features have been added to existing oscillators. By modifying the aforementioned synthetic oscillator with coupled positive- and negative-feedback loops [68], Mondragon-Palomino *et al.* have built an oscillator that can be entrained by external periodic signals [72]. Butzin *et al.* [73] further demonstrated that such synthetic oscillators can also be entrained by aperiodic signals, similar to the entrainment of cells in our body by the noisy natural signals. Modifying the same dual-feedback oscillator [68] through a single amino acid mutation to its core repressor, Hussain *et al.* have built an oscillator with temperature compensation [74]. The temperature compensation, i.e., keeping a constant period over a range of temperatures, is an essential property of circadian clock. In addition, Potvin-Trottier *et al.* focused on noise resistance and built an improved version of repressilator that can lead to synchrony in bacteria without coupling [75].

Second, the ability to program population-level dynamics and emergent collective behavior has become increasingly interesting in synthetic biology. Using quorum sensing to couple individual genetic oscillators, Danino *et al.* was able to construct synchronized oscillations at the colony level. In microfluidics devices, they demonstrated a variety of spatiotemporal waves propagating across cellular populations [76]. This work was further developed by Prindle *et al.* by coupling 12,000 such quorum-sensing synchronized colonies, called “biopixels”, in a macroscopic array, through gas-phase redox signaling. The ability to synchronize a large scale of colonies across the entire array enabled the construction of a low-cost biosensor that can detect heavy metals like arsenic via modulation of the frequency [81]. Chen *et al.* further extended the monoclonal system to create a synthetic microbial consortium containing two distinct cell types. The “activator” cells and “repressor” cells express orthogonal cell-signaling molecules and can only oscillate when both strains were present [77]. They also showed that certain network topologies of the two-strain circuit exhibited more robust oscillations than others. A recent study has developed a more complicated synthetic microbial community through multiplexed quorum sensing circuits [82]. Although all above-mentioned studies have utilized quorum sensing as a key design to produce population-level dynamics, Marguet *et al.* were able to construct oscillations in bacterial population density that required no quorum-sensing genes or promoters. Instead, the oscillations arise through the unexpected interplay of the host cell and the density-dependent plasmid amplification that established a population-level negative feedback. This study has highlighted the importance of considering “hidden interactions” between the synthetic circuits and the pre-existing metabolic and regulatory networks in complex host cells [83].

Third, integrating synthetic components into natural biological oscillators can modulate the endogenous oscillator behaviors. By adding MTF1 in the p53 signaling pathway, Toettcher *et al.* constructed an oscillatory system that is tunable in frequency [78]. In another study, Dies *et al.* linked the cell division cycle to a dual-feedback oscillator [68] in *E. coli*, by driving the *hda* and *dnaN* genes that inhibit the initiation of chromosomal replication, under the oscillator. In this engineered system, they observed the entrainment between the synthetic oscillator and the cell cycle [84].

Finally, studies have started to introduce synthetic circuits into organisms to enable a new function or application. A recent study has reconstructed the cyanobacterial KaiABC oscillator in *E. coli*, making this endogenously non-circadian bacterium perform circadian rhythms [85]. This demonstrated that a circadian

oscillator is transplantable to a heterologous organism. Moreover, a study engineered a bacterium capable of synchronous lysis at a threshold population density [86]. Introducing the lysis strain in combination with a chemotherapy, the study also demonstrated its clinical significance.

ANALYSIS OF BIOLOGICAL OSCILLATORS USING A CELL FREE SYSTEM

While many oscillators have been created in live bacteria and cell lines, oscillators have also been reconstituted *in vitro* in well-defined cell-free systems for decades. Comparing to live cells, a cell-free system has several unique advantages. First, a cell-free system usually contains only the most essential components in a test tube, which reduces potential interferences from the complex intracellular and extracellular environment. Second, it is convenient to introduce recombinant plasmids, mRNAs, proteins, as well as small molecules and drugs into the cell-free system, to precisely tune its oscillatory reactions, without worrying about the cytotoxicity, delivery efficiency, cross-talks, etc. Because of such flexibility and specificity of introducing molecules, it is more efficient and less time-consuming to design and test a functional circuit in cell-free systems than in living systems. Remarkably, it makes dissection of the circuits much easier, allowing for obtaining the steady state response function of each dissected reaction, parameters of which are crucial to build models.

In this section, we will specifically review the extensive application of a cell-free extract system [87], as a successful example, that has made major advances in understanding the cell cycle progression [1,22,88–92]. We will then extend this discussion to include a broader spectrum of cell-free systems and oscillators.

The cell free assays have made major contributions to the initial discovery and characterization of the central mitotic regulators. Extracts prepared from eggs of *Xenopus laevis* have enabled the first purification and *in vitro* kinase activity characterization of the maturation-promoting factor (MPF), later known as the protein complex cyclin B1-Cdk1 [87,93,94]. Clam oocyte extracts have allowed for the first discovery of anaphase-promoting complex or cyclosome (APC/C) [95], functioning as a E3 ubiquitin ligase. The activation of cyclin B1-Cdk1 drives mitotic entry and activates APC/C-Cdc20, which in turn marks the cyclin B1 for degradation and deactivates Cdk1, resulting in mitotic exit, and completing a core negative feedback loop. Studies making use of cycloheximide-treated interphase *Xenopus* egg extracts have shown that when adding APC-resistant cyclin mutants into the extracts, the extracts

approach a steady state of Cdk1 activity rather than oscillating [87]. This confirms the essential role of the negative-feedback loop in mitotic oscillations.

In theory, a negative feedback alone can generate oscillations. However, additional positive feedback loops through the regulations of Wee1 and Cdc25 are evolutionarily conserved [96,97]. Compromising the positive feedback loops will suppress oscillations in *Xenopus* egg extracts [91], suggesting that these are essential for sustained embryonic cell cycle oscillations. Together, these studies have identified the core architecture of cell cycles as interlinked positive and negative feedback loops, a commonly occurring motif also found in many other biological oscillators.

In addition to discovering the core architecture, cell-free assays have been combined with theory to dissect the interconnected feedback loops and investigate the function of each leg in isolation, in greater details. Tyson & Novak [98] and Thron [99] first proposed that the Cdk1/Cdc25/Wee1 system functions as a bistable trigger for mitosis. This hypothesis was then examined in quantitative measurements of *Xenopus* cell-free systems, showing that both Wee1 and Cdc25 respond to Cdk1 in an ultrasensitive (i.e., highly non-linear) manner [90–92,100]. In a rate balance analysis [101], this mirror-image, two-loop structure composed of Cdk1, Cdc25, and Wee1 with ultrasensitive responses makes the system substantially easier to generate a bistable response than otherwise. Bistable switch is a key consequence of positive feedback that delays the negative-feedback signal and prevents the system damped into a stable steady state [64]. Indeed, this has an experimental evidence that short-circuiting positive feedback in cell-free cycling *Xenopus* extracts makes the cell cycle oscillations damped [91].

Further studies, by integrating real-time fluorescence assays into the cell-free system, have revealed the core negative feedback system to operate as a time-delayed, digital switch, with a time lag of ~15 min between Cdk1 and APC/C-Cdc20 activation and a tremendously high degree of ultrasensitivity [1,2]. A simple, analytically tractable model has been developed to show how the attributes of time delay and ultrasensitivity contribute to the generation of robust, clock-like oscillations. A mechanism centered on the multisite phosphorylation of APC has also been postulated to explain how the ultrasensitivity and time delay are generated.

All together, these studies have proved that cell-free extracts are amenable to quantitative biochemical and synthetic approaches. By reconstituting and analyzing mitotic cycles in cell-free extracts, they have identified the key mechanisms for the mitotic oscillations: negative feedback, sufficient time delay, sufficient “nonlinearity” of the reaction kinetics, positive feedback, all consistent with the general requirements for biochemical oscillations [64].

Cell cycles are not the only study of oscillatory circuit that benefits from detailed quantitative analysis enabled by a simple cell-free system. In a pioneering study, Kondo and colleagues have reconstituted a simplest possible circadian oscillator *in vitro* using only three proteins: KaiA, KaiB, and KaiC [102]. In the presence of ATP as an energy supply, these three proteins can generate self-sustained oscillations with a period of about 24 hours and that are temperature compensated. Both the 24-hour period and temperature compensation are the defining characteristics of circadian oscillations observed *in vivo*. Without transcription or translation, this cell-free system has paved the way to examining the kinetic interactions of three Kai proteins in a detailed and quantitative manner [103].

In addition to recreating natural circuits such as cell cycles and circadian clocks, *de novo* engineered artificial networks have also been increasingly constructed thanks to the recent advances in synthesizing well-defined biopolymers and replicating systems *in vitro* [104]. These synthetic networks can not only perform various functions such as polymerase chain reaction (PCR) [105], transcription/translation machinery [106–108], bistable switches [109], etc., but can also behave as oscillators, such as DNA-based predator-prey molecular ecosystems [110,111], *in vitro* transcriptional oscillators [112–116], inorganic chemical reactions such as the Belousov-Zhabotinsky (BZ) reaction [117,118]. The recent development in microfluidics techniques has significantly improved the cell-free system by simplifying the fabrication process or creating high-throughput arrays [119–122].

A broad variety of cell-free systems (e.g., extracts from archaea, protozoa, plants, insects, etc.) allowing for *in vitro* transcription and translation [123] can serve as a powerful platform for synthesizing and examining oscillatory circuits. Besides the well-established *Xenopus* egg extracts [93], extracts widely prepared from other eukaryotic systems including yeast cells [124], clam oocytes [95], *Drosophila* embryos [120,121], mammalian cell lines [125,126], have been applied to the investigation of mechanisms underlying cell cycles and metabolic cycles. Prokaryotic cell-free systems such as *E. coli* cytoplasmic extracts have been developed to provide a highly versatile cell-free platform to construct complex biological systems. These include the protein synthesis using recombinant elements (PURE) system [127], the hybrid bacteriophage — *Escherichia coli* transcription and translation (TX-TL) system [128], and more recently, the all *E. coli* TX-TL system [129]. Oscillators built in these prokaryotic cell-free systems, e.g., the *in vitro* ring oscillators, show similar properties as the transcriptional oscillatory systems built *in vivo* [79,112]. Recent development in synthetic biology also makes it possible

to produce modular oscillators using nucleic acid [130,131] or organic molecules [132]. It is also possible to generate biological trigger waves [22] in cell-free systems when allowing diffusion, similar to the well-known BZ waves.

PERSPECTIVE

Systems and synthetic biology has been a promising approach to delineate an oscillator's mechanism by building and analyzing the functions at the systems level. An ultimate goal of synthetic biology is to engineer from the bottom-up a complete customized system with desired functions and applications. To build such an oscillatory system, a set of building blocks, which correspond to the functional modules in biological systems, must be available on hand. Using theoretical and experimental approaches, a set of modules of oscillation have been found and verified, including time delayed negative feedbacks and amplified negative feedbacks [133]. However, besides central oscillatory modules, peripheral modifications are also found to be highly conserved among different biological systems [65]. Different approaches have been used to identify functions of these auxiliary topological structures. These include: testing a hypothesis in a few oscillatory structures [65,134–136]; performing evolutionary network search [137,138]; and doing an exhaustive network enumeration [139,140]. Although these studies have made progress in identifying some peripheral modifications that help promote the robustness of an oscillator, the role of many other peripheral modifications remains unclear. Further investigations, both theoretically and experimentally, are needed to understand it.

The development of statistical tools and bioinformatics methods has also been used to push forward progress in system and synthetic biology. A deeper understanding of genomics is a breakthrough in 21st century biology. With the help of functional genome predictions, synthetic biologist can now make much more complicated biological networks to facilitate the understanding of biological system [141]. Development in data science and computational methods is another significant event that helps understanding and designing of biological oscillators. The statistical method like machine learning is used to identify functional proteins in biological systems [142]. Artificial intelligence is also helping the development of smarter cell-free systems that can automatically produce better proteins in parallel [143]. With greater computational power, we can now simulate much more complicated networks in shorter time, which is immensely helpful in theoretical research, but it also means that methods in data science will be widely used to investigate the data produced by high throughput simulations [144].

There are still many opening questions in biological oscillators, the understanding of which will be of both academic and clinical interest. Findings from studying essential biological and synthetic oscillators have been applied in the development of effective drugs for clock-related diseases [145,146] and other medical application [86]. The already established quantitative approaches in systems and synthetic biology will be facilitated by new methods from other fields such as engineering and bioinformatics, which will enable greater achievements in the scientific and medical studies on biological oscillators.

ACKNOWLEDGEMENTS

This work was supported by the National Science Foundation (Early CAREER Grant #1553031) and the National Institutes of Health (MIRA #GM119688).

COMPLIANCE WITH ETHICAL GUIDELINES

The authors Zhengda Li and Qiong Yang declare that they have no conflict of interests.

This article is a review article and does not contain any studies with human or animal subjects performed by any of the authors.

REFERENCES

1. Yang, Q. and Ferrell, J. E. Jr. (2013) The Cdk1-APC/C cell cycle oscillator circuit functions as a time-delayed, ultrasensitive switch. *Nat. Cell Biol.*, 15, 519–525
2. Ferrell, J. E. Jr, Tsai, T. Y. and Yang, Q. (2011) Modeling the cell cycle: why do certain circuits oscillate? *Cell*, 144, 874–885
3. Periasamy, M., Bhupathy, P. and Babu, G. J. (2008) Regulation of sarcoplasmic reticulum Ca²⁺ ATPase pump expression and its relevance to cardiac muscle physiology and pathology. *Cardiovasc. Res.*, 77, 265–273
4. Liu, N. and Priori, S. G. (2008) Disruption of calcium homeostasis and arrhythmogenesis induced by mutations in the cardiac ryanodine receptor and calsequestrin. *Cardiovasc. Res.*, 77, 293–301
5. Bénazéraf, B. and Pourquié, O. (2013) Formation and segmentation of the vertebrate body axis. *Annu. Rev. Cell Dev. Biol.*, 29, 1–26
6. Mara, A. and Holley, S. A. (2007) Oscillators and the emergence of tissue organization during zebrafish somitogenesis. *Trends Cell Biol.*, 17, 593–599
7. Bell-Pedersen, D., Cassone, V. M., Earnest, D. J., Golden, S. S., Hardin, P. E., Thomas, T. L. and Zoran, M. J. (2005) Circadian rhythms from multiple oscillators: lessons from diverse organisms. *Nat. Rev. Genet.*, 6, 544–556
8. Hodgkin, A. L. and Huxley, A. F. (1952) A quantitative description of membrane current and its application to conduction and excitation in nerve. *J. Physiol.*, 117, 500–544
9. Tomida, T., Takekawa, M. and Saito, H. (2015) Oscillation of p38 activity controls efficient pro-inflammatory gene expression. *Nat.*

- Commun., 6, 8350
10. Zambrano, S., De Toma, I., Piffer, A., Bianchi, M. E. and Agresti, A. (2016) NF- κ B oscillations translate into functionally related patterns of gene expression. *eLife*, 5, e09100
 11. Batchelor, E., Loewer, A., Mock, C. and Lahav, G. (2011) Stimulus-dependent dynamics of p53 in single cells. *Mol. Syst. Biol.*, 7, 488
 12. Dunlap, J. C. (1999) Molecular bases for circadian clocks. *Cell*, 96, 271–290
 13. Gallego, M. and Virshup, D. M. (2007) Post-translational modifications regulate the ticking of the circadian clock. *Nat. Rev. Mol. Cell Biol.*, 8, 139–148
 14. Brown, H., DiFrancesco, D. and Noble, S. (1979) Cardiac pacemaker oscillation and its modulation by autonomic transmitters. *J. Exp. Biol.*, 81, 175–204
 15. Paydarfar, D. and Eldridge, F. L. (1987) Phase resetting and dysrhythmic responses of the respiratory oscillator. *Am. J. Physiol.*, 252, R55–R62
 16. Hartwell, L. H. and Kastan, M. B. (1994) Cell cycle control and cancer. *Science*, 266, 1821–1828
 17. McDonald, E. R. 3rd and El-Deiry, W. S. (2000) Cell cycle control as a basis for cancer drug development (Review). *Int. J. Oncol.*, 16, 871–957
 18. Kastan, M. B. and Bartek, J. (2004) Cell-cycle checkpoints and cancer. *Nature*, 432, 316–323
 19. Gelens, L., Anderson, G. A. and Ferrell, J. E. Jr. (2014) Spatial trigger waves: positive feedback gets you a long way. *Mol. Biol. Cell*, 25, 3486–3493
 20. Novák, B. and Bentrup, F. W. (1972) An electrophysiological study of regeneration in *Acetabularia mediterranea*. *Planta*, 108, 227–244
 21. Gerisch, G. (1968) Cell aggregation and differentiation in *Dictyostelium*. *Curr. Top. Dev. Biol.*, 3, 157–197
 22. Chang, J. B. and Ferrell, J. E. Jr. (2013) Mitotic trigger waves and the spatial coordination of the *Xenopus* cell cycle. *Nature*, 500, 603–607
 23. Goodwin, B. C. and Cohen, M. H. (1969) A phase-shift model for the spatial and temporal organization of developing systems. *J. Theor. Biol.*, 25, 49–107
 24. Oates, A. C., Morelli, L. G. and Ares, S. (2012) Patterning embryos with oscillations: structure, function and dynamics of the vertebrate segmentation clock. *Development*, 139, 625–639
 25. Purvis, J. E., Karhohs, K. W., Mock, C., Batchelor, E., Loewer, A. and Lahav, G. (2012) p53 dynamics control cell fate. *Science*, 336, 1440–1444
 26. Isomura, A. and Kageyama, R. (2014) Ultradian oscillations and pulses: coordinating cellular responses and cell fate decisions. *Development*, 141, 3627–3636
 27. Ainsworth, M., Lee, S., Cunningham, M. O., Traub, R. D., Kopell, N. J. and Whittington, M. A. (2012) Rates and rhythms: a synergistic view of frequency and temporal coding in neuronal networks. *Neuron*, 75, 572–583
 28. Nelson, D. E., Ihekweba, A. E., Elliott, M., Johnson, J. R., Gibney, C. A., Foreman, B. E., Nelson, G., See, V., Horton, C. A., Spiller, D. G., *et al.* (2004) Oscillations in NF- κ B signaling control the dynamics of gene expression. *Science*, 306, 704–708
 29. Proctor, C. J. and Gray, D. A. (2008) Explaining oscillations and variability in the p53-Mdm2 system. *BMC Syst. Biol.*, 2, 75
 30. Salazar, C., Politi, A. Z. and Höfer, T. (2008) Decoding of calcium oscillations by phosphorylation cycles: analytic results. *Biophys. J.*, 94, 1203–1215
 31. Cai, L., Dalal, C. K. and Elowitz, M. B. (2008) Frequency-modulated nuclear localization bursts coordinate gene regulation. *Nature*, 455, 485–490
 32. Hodgkin, A. L., Huxley, A. F. and Katz, B. (1952) Measurement of current-voltage relations in the membrane of the giant axon of *Loligo*. *J. Physiol.*, 116, 424–448
 33. Hodgkin, A. L. and Katz, B. (1949) The effect of temperature on the electrical activity of the giant axon of the squid. *J. Physiol.*, 109, 240–249
 34. Hodgkin, A. L. and Huxley, A. F. (1952) The dual effect of membrane potential on sodium conductance in the giant axon of *Loligo*. *J. Physiol.*, 116, 497–506
 35. Hodgkin, A. L. and Huxley, A. F. (1952) The components of membrane conductance in the giant axon of *Loligo*. *J. Physiol.*, 116, 473–496
 36. Hodgkin, A. L. and Huxley, A. F. (1952) Currents carried by sodium and potassium ions through the membrane of the giant axon of *Loligo*. *J. Physiol.*, 116, 449–472
 37. Ghosh, A. and Chance, B. (1964) Oscillations of glycolytic intermediates in yeast cells. *Biochem. Biophys. Res. Commun.*, 16, 174–181
 38. Chance, B., Hess, B. and Betz, A. (1964) DPNH oscillations in a cell-free extract of *S. carlsbergensis*. *Biochem. Biophys. Res. Commun.*, 16, 182–187
 39. Chance, B., Schoener, B. and Elsaesser, S. (1965) Metabolic control phenomena involved in damped sinusoidal oscillations of reduced diphosphopyridine nucleotide in a cell-free extract of *Saccharomyces carlsbergensis*. *J. Biol. Chem.*, 240, 3170–3181
 40. Higgins, J. (1964) A chemical mechanism for oscillation of glycolytic intermediates in yeast cells. *Proc. Natl. Acad. Sci. USA*, 51, 989–994
 41. Pye, K. and Chance, B. (1966) Sustained sinusoidal oscillations of reduced pyridine nucleotide in a cell-free extract of *Saccharomyces carlsbergensis*. *Proc. Natl. Acad. Sci. USA*, 55, 888–894
 42. Gerisch, G., Fromm, H., Huesgen, A. and Wick, U. (1975) Control of cell-contact sites by cyclic AMP pulses in differentiating *Dictyostelium* cells. *Nature*, 255, 547–549
 43. Olsen, L. F. and Degn, H. (1978) Oscillatory kinetics of the peroxidase-oxidase reaction in an open system. *Experimental and theoretical studies. Biochim. Biophys. Acta*, 523, 321–334
 44. Goodwin, B. C. (1963) Temporal Organization in Cells. In *A Dynamic Theory of Cellular Control Processes*. New York: Academic Press
 45. Liu, Y., Tsinoremas, N. F., Johnson, C. H., Lebedeva, N. V., Golden, S. S., Ishiura, M. and Kondo, T. (1995) Circadian orchestration of gene expression in cyanobacteria. *Genes Dev.*, 9,

- 1469–1478
46. Crosthwaite, S. K., Dunlap, J. C. and Loros, J. J. (1997) *Neurospora wc-1* and *wc-2*: transcription, photoresponses, and the origins of circadian rhythmicity. *Science*, 276, 763–769
 47. Hamblen, M. J., White, N. E., Emery, P. T. J., Kaiser, K. and Hall, J. C. (1998) Molecular and behavioral analysis of four period mutants in *Drosophila melanogaster* encompassing extreme short, novel long, and unorthodox arrhythmic types. *Genetics*, 149, 165–178
 48. Antoch, M. P., Song, E. J., Chang, A. M., Vitaterna, M. H., Zhao, Y., Wilsbacher, L. D., Sangoram, A. M., King, D. P., Pinto, L. H. and Takahashi, J. S. (1997) Functional identification of the mouse circadian *Clock* gene by transgenic BAC rescue. *Cell*, 89, 655–667
 49. Millar, A.J. and Kay, S. A. (1997) The genetics of phototransduction and circadian rhythms in *Arabidopsis*. *BioEssays*, 19, 209–214
 50. Goldbeter, A. (1996) Biochemical Oscillations and Cellular Rhythms: The Molecular Bases of Periodic and Chaotic Behaviour. In *Physics Today*. New York: Cambridge University Press
 51. Goldbeter, A. (2007) Biological Rhythms as Temporal Dissipative Structures. Special Volume in Memory of Ilya Prigogine: *Advances in Chemical Physics*. pp. 253–295. New York: John Wiley & Sons, Inc.
 52. Yang, Q., Pando, B. F., Dong, G., Golden, S. S. and van Oudenaarden, A. (2010) Circadian gating of the cell cycle revealed in single cyanobacterial cells. *Science*, 327, 1522–1526
 53. Bieler, J., Cannavo, R., Gustafson, K., Gobet, C., Gatfield, D. and Naef, F. (2014) Robust synchronization of coupled circadian and cell cycle oscillators in single mammalian cells. *Mol. Syst. Biol.*, 10, 739
 54. Feillet, C., van der Horst, G. T., Levi, F., Rand, D. A. and Delaunay, F. (2015) Coupling between the circadian clock and cell cycle oscillators: implication for healthy cells and malignant growth. *Front. Neurol.*, 6, 96
 55. Sevim, V., Gong, X. and Socolar, J. E. S. (2010) Reliability of transcriptional cycles and the yeast cell-cycle oscillator. *PLoS Comput. Biol.*, 6, e1000842
 56. Longtin, A. (1993) Stochastic resonance in neuron models. *J. Stat. Phys.*, 70, 309–327
 57. Glass, L. (2001) Synchronization and rhythmic processes in physiology. *Nature*, 410, 277–284
 58. Prigogine, I., Lefever, R., Goldbeter, A. and Herschkowitz-Kaufman, M. (1969) Symmetry breaking instabilities in biological systems. *Nature*, 223, 913–916
 59. FitzHugh, R. (1961) Impulses and physiological states in theoretical models of nerve membrane. *Biophys. J.*, 1, 445–466
 60. Morris, C. and Lecar, H. (1981) Voltage oscillations in the barnacle giant muscle fiber. *Biophys. J.*, 35, 193–213
 61. Goodwin, B. C. (1965) Oscillatory behavior in enzymatic control processes. *Adv. Enzyme Regul.*, 3, 425–428
 62. Friesen, W. O. and Block, G. D. (1984) What is a biological oscillator? *Am. J. Physiol.*, 246, R847–R853
 63. Barkai, N. and Leibler, S. (2000) Circadian clocks limited by noise. *Nature*, 403, 267–268
 64. Novák, B. and Tyson, J. J. (2008) Design principles of biochemical oscillators. *Nat. Rev. Mol. Cell Biol.*, 9, 981–991
 65. Tsai, T. Y., Choi, Y. S., Ma, W., Pomerening, J. R., Tang, C. and Ferrell, J. E. Jr. (2008) Robust, tunable biological oscillations from interlinked positive and negative feedback loops. *Science*, 321, 126–129
 66. Elowitz, M. B. and Leibler, S. (2000) A synthetic oscillatory network of transcriptional regulators. *Nature*, 403, 335–338
 67. Atkinson, M. R., Savageau, M. A., Myers, J. T. and Ninfa, A. J. (2003) Development of genetic circuitry exhibiting toggle switch or oscillatory behavior in *Escherichia coli*. *Cell*, 113, 597–607
 68. Stricker, J., Cookson, S., Bennett, M. R., Mather, W. H., Tsimring, L. S. and Hasty, J. (2008) A fast, robust and tunable synthetic gene oscillator. *Nature*, 456, 516–519
 69. Fung, E., Wong, W. W., Suen, J. K., Bulter, T., Lee, S. G. and Liao, J. C. (2005) A synthetic gene-metabolic oscillator. *Nature*, 435, 118–122
 70. Tigges, M., Marquez-Lago, T. T., Stelling, J. and Fussenegger, M. (2009) A tunable synthetic mammalian oscillator. *Nature*, 457, 309–312
 71. Tigges, M., Dénervaud, N., Greber, D., Stelling, J. and Fussenegger, M. (2010) A synthetic low-frequency mammalian oscillator. *Nucleic Acids Res.*, 38, 2702–2711
 72. Mondragón-Palomino, O., Danino, T., Selimkhanov, J., Tsimring, L. and Hasty, J. (2011) Entrainment of a population of synthetic genetic oscillators. *Science*, 333, 1315–1319
 73. Butzin, N. C., Hochendoner, P., Ogle, C. T., Hill, P. and Mather, W. H. (2016) Marching along to an offbeat drum: entrainment of synthetic gene oscillators by a noisy stimulus. *ACS Synth. Biol.*, 5, 146–153
 74. Hussain, F., Gupta, C., Hirning, A. J., Ott, W., Matthews, K. S., Josic, K. and Bennett, M. R. (2014) Engineered temperature compensation in a synthetic genetic clock. *Proc. Natl. Acad. Sci. USA*, 111, 972–977
 75. Potvin-Trottier, L., Lord, N. D., Vinnicombe, G. and Paulsson, J. (2016) Synchronous long-term oscillations in a synthetic gene circuit. *Nature*, 538, 514–517
 76. Danino, T., Mondragón-Palomino, O., Tsimring, L. and Hasty, J. (2010) A synchronized quorum of genetic clocks. *Nature*, 463, 326–330
 77. Chen, Y., Kim, J. K., Hirning, A. J., Josić, K. and Bennett, M. R. (2015) Emergent genetic oscillations in a synthetic microbial consortium. *Science*, 349, 986–989
 78. Toettcher, J. E., Mock, C., Batchelor, E., Loewer, A. and Lahav, G. (2010) A synthetic-natural hybrid oscillator in human cells. *Proc. Natl. Acad. Sci. USA*, 107, 17047–17052
 79. Niederholtmeyer, H., Sun, Z. Z., Hori, Y., Yeung, E., Verpoorte, A., Murray, R. M. and Maerkl, S. J. (2015) Rapid cell-free forward engineering of novel genetic ring oscillators. *eLife*, 4, e09771
 80. Purcell, O., Savery, N. J., Grierson, C. S. and di Bernardo, M.

- (2010) A comparative analysis of synthetic genetic oscillators. *J. R. Soc. Interface*, 7, 1503–1524
81. Prindle, A., Samayoa, P., Razinkov, I., Danino, T., Tsimring, L. S. and Hasty, J. (2011) A sensing array of radically coupled genetic “biopixels”. *Nature*, 481, 39–44
 82. Scott, S. R. and Hasty, J. (2016) Quorum sensing communication modules for microbial consortia. *ACS Synth. Biol.*, 5, 969–977
 83. Marguet, P., Tanouchi, Y., Spitz, E., Smith, C. and You, L. (2010) Oscillations by minimal bacterial suicide circuits reveal hidden facets of host-circuit physiology. *PLoS One*, 5, e11909
 84. Dies, M., Galera-Laporta, L. and Garcia-Ojalvo, J. (2016) Mutual regulation causes co-entrainment between a synthetic oscillator and the bacterial cell cycle. *Integr. Biol.*, 8, 533–541
 85. Chen, A. H., Lubkowitz, D., Yeong, V., Chang, R. L. and Silver, P. A. (2015) Transplantability of a circadian clock to a noncircadian organism. *Sci. Adv.*, 1, e1500358
 86. Din, M. O., Danino, T., Prindle, A., Skalak, M., Selimkhanov, J., Allen, K., Julio, E., Atolia, E., Tsimring, L. S., Bhatia, S. N., *et al.* (2016) Synchronized cycles of bacterial lysis for *in vivo* delivery. *Nature*, 536, 81–85
 87. Murray, A. W., Solomon, M. J. and Kirschner, M. W. (1989) The role of cyclin synthesis and degradation in the control of maturation promoting factor activity. *Nature*, 339, 280–286
 88. Murray, A. W. (1991) Cell cycle extracts. *Methods Cell Biol.*, 36, 581–605
 89. Heald, R., Tournebise, R., Habermann, A., Karsenti, E. and Hyman, A. (1997) Spindle assembly in *Xenopus* egg extracts: respective roles of centrosomes and microtubule self-organization. *J. Cell Biol.*, 138, 615–628
 90. Kim, S. Y. and Ferrell, J. E. Jr. (2007) Substrate competition as a source of ultrasensitivity in the inactivation of Wee1. *Cell*, 128, 1133–1145
 91. Pomerening, J. R., Kim, S. Y. and Ferrell, J. E. Jr. (2005) Systems-level dissection of the cell-cycle oscillator: bypassing positive feedback produces damped oscillations. *Cell*, 122, 565–578
 92. Trunnell, N. B., Poon, A. C., Kim, S. Y. and Ferrell, J. E. Jr. (2011) Ultrasensitivity in the regulation of Cdc25C by Cdk1. *Mol. Cell*, 41, 263–274
 93. Lohka, M. J. and Maller, J. L. (1985) Induction of nuclear envelope breakdown, chromosome condensation, and spindle formation in cell-free extracts. *J. Cell Biol.*, 101, 518–523
 94. Lohka, M. J., Hayes, M. K. and Maller, J. L. (1988) Purification of maturation-promoting factor, an intracellular regulator of early mitotic events. *Proc. Natl. Acad. Sci. USA*, 85, 3009–3013
 95. Sudakin, V., Ganoth, D., Dahan, A., Heller, H., Hershko, J., Luca, F. C., Ruderman, J. V. and Hershko, A. (1995) The cyclosome, a large complex containing cyclin-selective ubiquitin ligase activity, targets cyclins for destruction at the end of mitosis. *Mol. Biol. Cell*, 6, 185–197
 96. Kumagai, A. and Dunphy, W. G. (1992) Regulation of the cdc25 protein during the cell cycle in *Xenopus* extracts. *Cell*, 70, 139–151
 97. Mueller, P. R., Coleman, T. R. and Dunphy, W. G. (1995) Cell cycle regulation of a *Xenopus* Wee1-like kinase. *Mol. Biol. Cell*, 6, 119–134
 98. Novak, B. and Tyson, J. J. (1993) Numerical analysis of a comprehensive model of M-phase control in *Xenopus* oocyte extracts and intact embryos. *J. Cell. Sci.*, 106, 1153–1168
 99. Thron, C. D. (1996) A model for a bistable biochemical trigger of mitosis. *Biophys. Chem.*, 57, 239–251
 100. Pomerening, J. R., Sontag, E. D. and Ferrell, J. E. Jr. (2003) Building a cell cycle oscillator: hysteresis and bistability in the activation of Cdc2. *Nat. Cell Biol.*, 5, 346–351
 101. Ferrell, J. E. Jr. (2008) Feedback regulation of opposing enzymes generates robust, all-or-none bistable responses. *Curr. Biol.*, 18, R244–R245
 102. Nakajima, M., Imai, K., Ito, H., Nishiwaki, T., Murayama, Y., Iwasaki, H., Oyama, T. and Kondo, T. (2005) Reconstitution of circadian oscillation of cyanobacterial KaiC phosphorylation *in vitro*. *Science*, 308, 414–415
 103. Rust, M. J., Markson, J. S., Lane, W. S., Fisher, D. S. and O’Shea, E. K. (2007) Ordered phosphorylation governs oscillation of a three-protein circadian clock. *Science*, 318, 809–812
 104. Forster, A. C. and Church, G. M. (2007) Synthetic biology projects *in vitro*. *Genome Res.*, 17, 1–6
 105. Nakano, M., Komatsu, J., Matsuura, S., Takashima, K., Katsura, S. and Mizuno, A. (2003) Single-molecule PCR using water-in-oil emulsion. *J. Biotechnol.*, 102, 117–124
 106. Tan, C., Saurabh, S., Bruchez, M. P., Schwartz, R. and Leduc, P. (2013) Molecular crowding shapes gene expression in synthetic cellular nanosystems. *Nat. Nanotechnol.*, 8, 602–608
 107. Shin, J. and Noireaux, V. (2012) An *E. coli* cell-free expression toolbox: application to synthetic gene circuits and artificial cells. *ACS Synth. Biol.*, 1, 29–41
 108. Noireaux, V. and Libchaber, A. (2004) A vesicle bioreactor as a step toward an artificial cell assembly. *Proc. Natl. Acad. Sci. USA*, 101, 17669–17674
 109. Kim, J., White, K. S. and Winfree, E. (2006) Construction of an *in vitro* bistable circuit from synthetic transcriptional switches. *Mol. Syst. Biol.*, 2, 68
 110. Ackermann, J., Wlotzka, B. and McCaskill, J. S. (1998) *In vitro* DNA-based predator-prey system with oscillatory kinetics. *Bull. Math. Biol.*, 60, 329–354
 111. Fujii, T. and Rondelez, Y. (2013) Predator-prey molecular ecosystems. *ACS Nano*, 7, 27–34
 112. Kim, J. and Winfree, E. (2011) Synthetic *in vitro* transcriptional oscillators. *Mol. Syst. Biol.*, 7, 465
 113. Montagne, K., Plasson, R., Sakai, Y., Fujii, T. and Rondelez, Y. (2011) Programming an *in vitro* DNA oscillator using a molecular networking strategy. *Mol. Syst. Biol.*, 7, 466
 114. Franco, E., Friedrichs, E., Kim, J., Jungmann, R., Murray, R., Winfree, E. and Simmel, F. C. (2011) Timing molecular motion and production with a synthetic transcriptional clock. *Proc. Natl. Acad. Sci. USA*, 108, E784–E793
 115. Weitz, M., Kim, J., Kapsner, K., Winfree, E., Franco, E. and Simmel, F. C. (2014) Diversity in the dynamical behaviour of a compartmentalized programmable biochemical oscillator. *Nat.*

- Chem., 6, 295–302
116. Hasatani, K., Leocmach, M., Genot, A. J., Estévez-Torres, A., Fujii, T. and Rondelez, Y. (2013) High-throughput and long-term observation of compartmentalized biochemical oscillators. *Chem. Commun. (Camb.)*, 49, 8090–8092
 117. Epstein, I. R., Vanag, V. K., Balazs, A. C., Kuksenok, O., Dayal, P. and Bhattacharya, A. (2012) Chemical oscillators in structured media. *Acc. Chem. Res.*, 45, 2160–2168
 118. Vanag, V. K. and Epstein, I. R. (2001) Pattern formation in a tunable medium: the Belousov-Zhabotinsky reaction in an aerosol OT microemulsion. *Phys. Rev. Lett.*, 87, 228301
 119. Good, M. C., Vahey, M. D., Skandarajah, A., Fletcher, D. A. and Heald, R. (2013) Cytoplasmic volume modulates spindle size during embryogenesis. *Science*, 342, 856–860
 120. Telley, I. A., Gáspár, I., Ephrussi, A. and Surrey, T. (2013) A single *Drosophila* embryo extract for the study of mitosis *ex vitro*. *Nat. Protoc.*, 8, 310–324
 121. Telley, I. A., Gáspár, I., Ephrussi, A. and Surrey, T. (2012) Aster migration determines the length scale of nuclear separation in the *Drosophila* syncytial embryo. *J. Cell Biol.*, 197, 887–895
 122. Ho, K. K., Lee, J. W., Durand, G., Majumder, S. and Liu, A. P. (2017) Protein aggregation with poly(vinyl) alcohol surfactant reduces double emulsion-encapsulated mammalian cell-free expression. *PLoS One*, 12, e0174689
 123. Zemella, A., Thoring, L., Hoffmeister, C. and Kubick, S. (2015) Cell-free protein synthesis: pros and cons of prokaryotic and eukaryotic systems. *ChemBioChem*, 16, 2420–2431
 124. Richard, P. (2003) The rhythm of yeast. *FEMS Microbiol. Rev.*, 27, 547–557
 125. Lu, Y., Lee, B. H., King, R. W., Finley, D. and Kirschner, M. W. (2015) Substrate degradation by the proteasome: a single-molecule kinetic analysis. *Science*, 348, 1250834
 126. Lu, Y., Wang, W. and Kirschner, M. W. (2015) Specificity of the anaphase-promoting complex: a single-molecule study. *Science*, 348, 1248737
 127. Shimizu, Y., Kanamori, T. and Ueda, T. (2005) Protein synthesis by pure translation systems. *Methods*, 36, 299–304
 128. Nevin, D. E. and Pratt, J. M. (1991) A coupled *in vitro* transcription-translation system for the exclusive synthesis of polypeptides expressed from the T7 promoter. *FEBS Lett.*, 291, 259–263
 129. Garamella, J., Marshall, R., Rustad, M. and Noireaux, V. (2016) The all *E. coli* TX-TL Toolbox 2.0: a platform for cell-free synthetic biology. *ACS Synth. Biol.*, 5, 344–355
 130. Genot, A. J., Baccouche, A., Sieskind, R., Aubert-Kato, N., Bredeche, N., Bartolo, J. F., Taly, V., Fujii, T. and Rondelez, Y. (2016) High-resolution mapping of bifurcations in nonlinear biochemical circuits. *Nat. Chem.*, 8, 760–767
 131. Wang, F. and Fan, C. (2016) DNA reaction networks: providing a panoramic view. *Nat. Chem.*, 8, 738–740
 132. Semenov, S. N., Kraft, L. J., Ainla, A., Zhao, M., Baghbanzadeh, M., Campbell, V. E., Kang, K., Fox, J. M. and Whitesides, G. M. (2016) Autocatalytic, bistable, oscillatory networks of biologically relevant organic reactions. *Nature*, 537, 656–660
 133. Tyson, J. J. and Novák, B. (2010) Functional motifs in biochemical reaction networks. *Annu. Rev. Phys. Chem.*, 61, 219–240
 134. Nguyen, L. K. (2012) Regulation of oscillation dynamics in biochemical systems with dual negative feedback loops. *J. R. Soc. Interface*, 9, 1998–2010
 135. Lomnitz, J. G. and Savageau, M. A. (2014) Strategy revealing phenotypic differences among synthetic oscillator designs. *ACS Synth. Biol.*, 3, 686–701
 136. Guantes, R. and Poyatos, J. F. (2006) Dynamical principles of two-component genetic oscillators. *PLoS Comput. Biol.*, 2, e30
 137. Wagner, A. (2005) Circuit topology and the evolution of robustness in two-gene circadian oscillators. *Proc. Natl. Acad. Sci. USA*, 102, 11775–11780
 138. Noman, N., Monjo, T., Moscato, P. and Iba, H. (2015) Evolving robust gene regulatory networks. *PLoS One*, 10, e0116258
 139. Castillo-Hair, S. M., Villota, E. R. and Coronado, A. M. (2015) Design principles for robust oscillatory behavior. *Syst. Synth. Biol.*, 9, 125–133
 140. Woods, M. L., Leon, M., Perez-Carrasco, R. and Barnes, C. P. (2016) A statistical approach reveals designs for the most robust stochastic gene oscillators. *ACS Synth. Biol.*, 5, 459–470
 141. Mukherji, S. and van Oudenaarden, A. (2009) Synthetic biology: understanding biological design from synthetic circuits. *Nat. Rev. Genet.*, 10, 859–871
 142. Anafi, R. C., Lee, Y., Sato, T. K., Venkataraman, A., Ramanathan, C., Kavakli, I. H., Hughes, M. E., Baggs, J. E., Growe, J., Liu, A. C., *et al.* (2014) Machine learning helps identify CHRONO as a circadian clock component. *PLoS Biol.*, 12, e1001840
 143. Caschera, F., Bedau, M. A., Buchanan, A., Cawse, J., de Lucrezia, D., Gazzola, G., Hanczyc, M. M. and Packard, N. H. (2011) Coping with complexity: machine learning optimization of cell-free protein synthesis. *Biotechnol. Bioeng.*, 108, 2218–2228
 144. Trejo Banos, D., Millar, A. J. and Sanguinetti, G. (2015) A Bayesian approach for structure learning in oscillating regulatory networks. *Bioinformatics*, 31, 3617–3624
 145. Malumbres, M. and Barbacid, M. (2009) Cell cycle, CDKs and cancer: a changing paradigm. *Nat. Rev. Cancer*, 9, 153–166
 146. Dart, A. (2016) Tumorigenesis: cancer goes tick tock. *Nat. Rev. Cancer*, 16, 409